



US 20160264972A1

(19) **United States**(12) **Patent Application Publication****Bavari et al.**(10) **Pub. No.: US 2016/0264972 A1**(43) **Pub. Date: Sep. 15, 2016**(54) **COMPOSITIONS AND METHODS FOR
INHIBITING EXPRESSION OF A GENE
FROM THE EBOLA VIRUS**(71) Applicants: **Alnylam Pharmaceuticals Inc.**,
Cambridge, MA (US); **The Government
of the United States, as Represented by
the Secretary of the Army**, Washington,
DC (US)(72) Inventors: **Sina Bavari**, Frederick, MD (US); **Kelly
Lyn Warfield**, Adamstown, MD (US);
Pamela Tan, Kulmbach (DE); **Anna
Borodovsky**, Cambridge, MA (US);
Tatiana Novobrantseva, Wellesley, MA
(US); **Antonin de Fougères**,
Brookline, MA (US)(21) Appl. No.: **14/882,388**(22) Filed: **Oct. 13, 2015****Related U.S. Application Data**(62) Division of application No. 14/279,156, filed on May
15, 2014, now Pat. No. 9,187,516, which is a divisionof application No. 13/627,508, filed on Sep. 26, 2012,
now Pat. No. 8,735,369, which is a division of appli-
cation No. 13/093,681, filed on Apr. 25, 2011, now Pat.
No. 8,354,390, which is a division of application No.
12/789,414, filed on May 27, 2010, now Pat. No.
7,973,020, which is a division of application No.
12/055,295, filed on Mar. 25, 2008, now Pat. No.
7,759,320.(60) Provisional application No. 60/908,793, filed on Mar.
29, 2007.**Publication Classification**(51) **Int. Cl.**
C12N 15/113 (2006.01)(52) **U.S. Cl.**
CPC **C12N 15/1131** (2013.01); **C12N 2310/14**
(2013.01); **C12N 2310/315** (2013.01); **C12N**
2310/321 (2013.01); **C12N 2310/3521**
(2013.01)(57) **ABSTRACT**The invention relates to a double-stranded ribonucleic acid
(dsRNA) for inhibiting the expression of a gene from the
Ebola virus.




FIG 1.




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8

COMPOSITIONS AND METHODS FOR INHIBITING EXPRESSION OF A GENE FROM THE EBOLA VIRUS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a divisional of U.S. patent application Ser. No. 14/279,156, filed May 15, 2014, which is a divisional of U.S. patent application Ser. No. 13/627,508, filed Sep. 26, 2012, issued as U.S. Pat. No. 8,735,369, which is a divisional of U.S. patent application Ser. No. 13/093,681, filed Apr. 25, 2011, issued as U.S. Pat. No. 8,354,390, which is a divisional of U.S. patent application Ser. No. 12/789,414, filed May 27, 2010, issued as U.S. Pat. No. 7,973,020, which is a divisional of U.S. patent application Ser. No. 12/055,295, filed Mar. 25, 2008, issued as U.S. Pat. No. 7,759,320, which claims priority to U.S. Provisional Application Ser. No. 60/908,793, filed Mar. 29, 2007, the entire contents of which are hereby incorporated by reference.

GOVERNMENT SUPPORT

[0002] This invention was made with government support under contract number HHSN266200600012C, awarded by the Department of Health and Human Services (DHHS). The government has certain rights in the invention.

REFERENCE TO A SEQUENCE LISTING

[0003] This application includes a Sequence Listing submitted electronically as a text file named 30589_US_Sequence_Listing.txt, created on Oct. 13, 2015, with a size of 760 KB and comprising 1108 sequences. The sequence listing is hereby incorporated by reference.

FIELD OF THE INVENTION

[0004] This invention relates to double-stranded ribonucleic acid (dsRNA), and its use in mediating RNA interference to inhibit the expression of one of the genes of the Ebola virus and the use of the dsRNA to treat pathological processes mediated by Ebola infection, such as systemic hemorrhage and multi-organ failure.

BACKGROUND OF THE INVENTION

[0005] Ebola Virus

[0006] Minus-strand (−) RNA viruses are major causes of human suffering that cause epidemics of serious human illness. In humans the diseases caused by these viruses include influenza (Orthomyxoviridae), mumps, measles, upper and lower respiratory tract disease (Paramyxoviridae), rabies (Rhabdoviridae), hemorrhagic fever (Filoviridae, Bunyaviridae and Arenaviridae), encephalitis (Bunyaviridae) and neurological illness (Bomaviridae). Virtually the entire human population is thought to be infected by many of these viruses.

[0007] The Ebola virus comes from the Filoviridae family, similar to the Marburg virus. It is named after the Ebola River in Zaire, Africa, near where the first outbreak was noted by Dr. Ngoy Mushola in 1976 after a significant outbreaks in both Yambuku, Zaire (now the Democratic Republic of the Congo), and Nzara, in western Sudan. Of 602 identified cases, there were 397 deaths.

[0008] The two strains identified in 1976 were named Ebola-Zaire (EBO-Z) and Ebola-Sudan (EBO-S). The outbreak in Sudan showed a lower fatality rate—50%—com-

pared to the 90% mortality rate of the Zaire strain. In 1990, a second, similar virus was identified in Reston, Virginia amongst monkeys imported from the Philippines, and was named Ebola-Reston.

[0009] Further outbreaks have occurred in Zaire/Congo (1995 and 2003), Gabon (1994, 1995 and 1996), and in Uganda (2000). A new subtype was identified from a single human case in the Côte d'Ivoire in 1994, EBO-CI.

[0010] Of around 1500 identified human Ebola infections, two-thirds of the patients have died. The animal (or other) reservoir which sustains the virus between outbreaks has not been identified.

[0011] Among humans, the Ebola virus is transmitted by direct contact with infected body fluids such as blood.

[0012] The incubation period of Ebola hemorrhagic fever varies from two days to four weeks. Symptoms are variable too, but the onset is usually sudden and characterised by high fever, prostration, myalgia, arthralgia, abdominal pains and headache. These symptoms progress to vomiting, diarrhea, oropharyngeal lesions, conjunctivitis, organ damage (notably the kidney and liver) by co-localized necrosis, proteinuria, and bleeding both internal and external, commonly through the gastrointestinal tract. Death or recovery to convalescence occurs within six to ten days of onset of symptomatology.

[0013] The development of a successful therapeutic for Ebola virus is a long-sought and seemingly difficult endeavor. Although they cause only a few hundred deaths worldwide each year, filoviruses are considered a significant world health threat and have many of the characteristics commonly associated with biological weapons since they can be grown in large quantities, can be fairly stable, are highly infectious as an aerosol, and are exceptionally deadly. Filoviruses are relatively simple viruses of 19 Kb genomes and consist of seven genes which encode nucleoprotein (NP), glycoprotein (GP), four smaller viral proteins (VP24, VP30, VP35 and VP40), and the RNA-dependent RNA polymerase (L protein) all in a single strand of negative-sensed RNA. Administration of type I interferons, therapeutic vaccines, immune globulins, ribavirin, and other nucleoside analogues have been somewhat successful in rodent Ebola virus models, but not in nonhuman primate infection models.

[0014] In view of the severity of the diseases caused by (−) RNA viruses, in particular members of the Filoviridae family of viruses, and the lack of effective prevention or therapies, it is therefore an object of the present invention to provide therapeutic compounds and methods for treating a host infected with a (−) RNA virus.

[0015] siRNA

[0016] Double-stranded RNA molecules (dsRNA) have been shown to block gene expression in a highly conserved regulatory mechanism known as RNA interference (RNAi). WO 99/32619 (Fire et al.) discloses the use of a dsRNA of at least 25 nucleotides in length to inhibit the expression of genes in *C. elegans*. dsRNA has also been shown to degrade target RNA in other organisms, including plants (see, e.g., WO 99/53050, Waterhouse et al.; and WO 99/61631, Heifetz et al.), *Drosophila* (see, e.g., Yang, D., et al., *Curr. Biol.* (2000) 10:1191-1200), and mammals (see WO 00/44895, Limmer; and DE 101 00 586.5, Kreutzer et al.). This natural mechanism has now become the focus for the development of a new class of pharmaceutical agents for treating disorders that are caused by the aberrant or unwanted regulation of a gene.

[0017] Recent reports have indicated that in vitro, RNAi may show some promising in reducing Ebola replication and providing protection in guinea pigs (Geisbert, et al., *The Journal of Infectious Diseases*, 193 (2006), 1650-1657). However, the RNAi agents examined were not designed against all known Ebola strains and were not selected for stability and other properties needed for in vivo therapeutic RNAi agents. Accordingly, despite significant advances in the field of RNAi, there remains a need for an agent that can selectively and efficiently silence a gene in the Ebola virus using the cell's own RNAi machinery that has both high biological activity and in vivo stability, and that can effectively inhibit replication of the Ebola virus for use in treating pathological processes mediated by Ebola infection.

SUMMARY OF THE INVENTION

[0018] The invention provides double-stranded ribonucleic acid (dsRNA), as well as compositions and methods for inhibiting the expression of the Ebola virus in a cell or mammal using such dsRNA. The invention also provides compositions and methods for treating pathological conditions and diseases caused by Ebola viral infection, such as systemic hemorrhage and multi-organ failure. The dsRNA featured in the invention includes an RNA strand (the antisense strand) having a region which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an mRNA transcript of a gene from the Ebola virus.

[0019] In one embodiment, the invention provides dsRNA molecules for inhibiting the expression of a gene of the Ebola virus and viral replication. The dsRNA comprises at least two sequences that are complementary to each other. The dsRNA comprises a sense strand comprising a first sequence and an antisense strand comprising a second sequence. The antisense strand comprises a nucleotide sequence which is substantially complementary to at least part of an mRNA encoded by a gene from the Ebola virus, and the region of complementarity is less than 30 nucleotides in length, generally 19-24 nucleotides in length. The dsRNA, upon contact with a cell infected with the Ebola virus, inhibits the expression of a gene from the Ebola virus by at least 40%.

[0020] For example, the dsRNA molecules of the invention can include a first sequence of the dsRNA that is selected from the group consisting of the sense sequences of Table 2 and the second sequence selected from the group consisting of the antisense sequences of Table 2. The dsRNA molecules featured in the invention can include naturally occurring nucleotides or can include at least one modified nucleotide, such as a 2'-O-methyl modified nucleotide, a nucleotide comprising a 5'-phosphorothioate group, and a terminal nucleotide linked to a cholesteryl derivative. Alternatively, the modified nucleotide may be chosen from the group of: a 2'-deoxy-2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a locked nucleotide, an abasic nucleotide, 2'-amino-modified nucleotide, 2'-alkyl-modified nucleotide, morpholino nucleotide, a phosphoramidate, and a non-natural base comprising nucleotide. Generally, such modified sequence will be based on a first sequence of said dsRNA selected from the group consisting of the sense sequences of Table 2 and a second sequence selected from the group consisting of the antisense sequences of Table 2.

[0021] In another embodiment, the invention provides a cell having a dsRNA of the invention. The cell is generally a mammalian cell, such as a human cell.

[0022] In another embodiment, the invention provides a pharmaceutical composition for inhibiting the replication of the Ebola virus in an organism, generally a human subject. The composition includes one or more of the dsRNA of the invention and a pharmaceutically acceptable carrier or delivery vehicle.

[0023] In another embodiment, the invention provides a method for inhibiting the expression of a gene in the Ebola virus in a cell, including the following steps:

[0024] (a) introducing into the cell a double-stranded ribonucleic acid (dsRNA), wherein the dsRNA includes at least two sequences that are complementary to each other. The dsRNA includes a sense strand having a first sequence and an antisense strand having a second sequence. The antisense strand includes a region of complementarity which is substantially complementary to at least a part of an mRNA encoded by the Ebola virus, and wherein the region of complementarity is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and optionally, wherein the dsRNA, upon contact with a cell infected with the Ebola virus, inhibits expression of a gene from the Ebola virus by at least 40%, such as in an assay described herein (e.g., a fluorescence-based assay); and

[0025] (b) maintaining the cell produced in step (a) for a time sufficient to obtain degradation of the mRNA transcript of a Ebola gene, thereby inhibiting expression of a gene from the Ebola virus in the cell.

[0026] In another embodiment, the invention provides methods for treating, preventing or managing pathological processes mediated by Ebola infection, such as systemic hemorrhage and multi-organ failure, comprising administering to a patient in need of such treatment, prevention or management a therapeutically or prophylactically effective amount of one or more of the dsRNAs of the invention.

[0027] In another embodiment, the invention provides vectors for inhibiting the expression of a gene of the Ebola virus in a cell, comprising a regulatory sequence operably linked to a nucleotide sequence that encodes at least one strand of a dsRNA of the invention.

[0028] In another embodiment, the invention provides a cell comprising a vector for inhibiting the expression of a gene of the Ebola virus in a cell. The vector comprises a regulatory sequence operably linked to a nucleotide sequence that encodes at least one strand of a dsRNA of the invention.

[0029] In one aspect, the invention provides for a method of increasing the life-span of a subject (e.g., a mammal, such as a human or nonhuman primate) infected with an Ebola virus. The method includes administering a dsRNA to the subject, where the dsRNA includes an antisense RNA strand having a region which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an mRNA transcript of a gene from the Ebola virus. The dsRNA is administered in an amount sufficient to increase the lifespan of the subject. In one embodiment, the dsRNA includes an antisense RNA strand having a region that is substantially complementary to at least part of an mRNA transcript of a gene selected from the VP30, VP35, NP, L, VP24, VP40 and GP genes. In a preferred embodiment, the dsRNA includes an antisense RNA strand having a region that is substantially complementary to at least part of an mRNA transcript of the VP35 gene. In some embodiments, the subject does not experience a decrease in one or both of lymphocyte or platelet count after administration of the

dsRNA. In other embodiments, the subject does not experience an increase in cytokine levels (e.g., IFN-alpha or TNF-alpha levels).

[0030] In another aspect, the invention features a method of decreasing viral titre in a subject (e.g., a mammal, such as a human or nonhuman primate) infected with an Ebola virus. The method includes administering a dsRNA to the subject, where the dsRNA includes an antisense RNA strand having a region which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an mRNA transcript of a gene from the Ebola virus. In one embodiment, the dsRNA includes an antisense RNA strand having a region that is substantially complementary to at least part of an mRNA transcript of the VP35 gene. In another embodiment, the subject does not experience a decrease in one or both of lymphocyte or platelet count after administration of the dsRNA. In other embodiments, the subject does not experience an increase in cytokine levels (e.g., IFN-alpha or TNF-alpha levels).

[0031] In another aspect, the invention features a method of sustaining lymphocyte or platelet count in a mammal (e.g., a human or nonhuman primate) infected with an Ebola virus. The method includes administering a dsRNA to the subject, where the dsRNA includes an antisense RNA strand having a region which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an mRNA transcript of a gene from the Ebola virus. In one embodiment, the dsRNA includes an antisense RNA strand having a region that is substantially complementary to at least part of an mRNA transcript of the VP35 gene. In other embodiments, the subject does not experience an increase in cytokine levels (e.g., IFN-alpha or TNF-alpha levels).

BRIEF DESCRIPTION OF THE FIGURES

[0032] FIG. 1 is a graph showing that siRNAs formulated with lipidoid LNP01 protected mice from a lethal Ebola virus challenge.

[0033] FIG. 2 is a graph showing that a single injection of a liposomally formulated siRNA delivered by ip or iv protected mice from a lethal Ebola challenge. VP35 siRNA was AD-11570

[0034] FIG. 3 is the structure of NP98 lipid.

[0035] FIG. 4 is a graph showing that siRNAs formulated with DODMA protected mice from a lethal Ebola virus challenge.

[0036] FIG. 5 is a graph showing that siRNAs formulated with DODMA were effective down to 0.04 mg/kg to protect mice injected with Ebola.

[0037] FIG. 6 is a graph showing that siRNAs formulated with DODMA were effective to protect guinea pigs from a lethal Ebola virus challenge.

[0038] FIG. 7 is a graph showing the efficacy of siRNAs against different Ebola genes formulated with DODMA in a guinea pig model of Ebola.

[0039] FIG. 8 is a graph presenting the observed decrease in viral titers in the serum of mice following administration of LNP01-formulated VP35 siRNA.

DETAILED DESCRIPTION OF THE INVENTION

[0040] The invention provides double-stranded ribonucleic acid (dsRNA), as well as compositions and methods for inhibiting the expression of a gene from the Ebola virus in a

cell or mammal using the dsRNA. The invention also provides compositions and methods for treating pathological conditions and diseases in a mammal caused by Ebola infection using dsRNA. dsRNA directs the sequence-specific degradation of mRNA through a process known as RNA interference (RNAi).

[0041] The dsRNA of the invention comprises an RNA strand (the antisense strand) having a region which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an mRNA transcript of a gene from the Ebola virus. The use of these dsRNAs enables the targeted degradation of mRNAs of genes that are implicated in replication and or maintenance of Ebola infection and the occurrence of systemic hemorrhage and multi-organ failure in a subject infected with the Ebola virus. Using cell-based and animal assays, the present inventors have demonstrated that very low dosages of these dsRNA can specifically and efficiently mediate RNAi, resulting in significant inhibition of expression of a gene from the Ebola virus. Thus, the methods and compositions of the invention comprising these dsRNAs are useful for treating pathological processes mediated by Ebolaviral infection by targeting a gene involved in Ebola replication and/or maintenance in a cell.

[0042] The following detailed description discloses how to make and use the dsRNA and compositions containing dsRNA to inhibit the expression of a gene from the Ebola virus, as well as compositions and methods for treating diseases and disorders caused by the infection with the Ebola virus, such as systemic hemorrhage and multi-organ failure. The pharmaceutical compositions of the invention comprise a dsRNA having an antisense strand comprising a region of complementarity which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an RNA transcript of a gene from the Ebola virus, together with a pharmaceutically acceptable carrier.

[0043] Accordingly, certain aspects of the invention provide pharmaceutical compositions comprising the dsRNA of the invention together with a pharmaceutically acceptable carrier, methods of using the compositions to inhibit expression of a gene in a gene from the Ebola virus, and methods of using the pharmaceutical compositions to treat diseases caused by infection with the Ebola virus.

[0044] I. Definitions

[0045] For convenience, the meaning of certain terms and phrases used in the specification, examples, and appended claims, are provided below. If there is an apparent discrepancy between the usage of a term in other parts of this specification and its definition provided in this section, the definition in this section shall prevail.

[0046] "G," "C," "A" and "U" each generally stand for a nucleotide that contains guanine, cytosine, adenine, and uracil as a base, respectively. However, it will be understood that the term "ribonucleotide" or "nucleotide" can also refer to a modified nucleotide, as further detailed below, or a surrogate replacement moiety. The skilled person is well aware that guanine, cytosine, adenine, and uracil may be replaced by other moieties without substantially altering the base pairing properties of an oligonucleotide comprising a nucleotide bearing such replacement moiety. For example, without limitation, a nucleotide comprising inosine as its base may base pair with nucleotides containing adenine, cytosine, or uracil. Hence, nucleotides containing uracil, guanine, or adenine

may be replaced in the nucleotide sequences of the invention by a nucleotide containing, for example, inosine. Sequences comprising such replacement moieties are embodiments of the invention.

[0047] As used herein, “Ebola viruses”, are members of the family Filoviridae, are associated with outbreaks of highly lethal hemorrhagic fever in humans and nonhuman primates. Human pathogens include Ebola Zaire, Ebola Sudan, and Ebola Ivory Coast. Ebola Reston is a monkey pathogen and is not considered a significant human pathogen. The natural reservoir of the virus is unknown and there are currently no available vaccines or effective therapeutic treatments for filovirus infections. The genome of Ebola virus consists of a single strand of negative sense RNA that is approximately 19 kb in length. This RNA contains seven sequentially arranged genes that produce 8 mRNAs upon infection. Ebola virions, like virions of other filoviruses, contain seven proteins: a surface glycoprotein (GP), a nucleoprotein (NP), four virion structural proteins (VP40, VP35, VP30, and VP24), and an RNA-dependent RNA polymerase (L) (Feldmann et al. (1992) *Virus Res.* 24, 1-19; Sanchez et al., (1993) *Virus Res.* 29, 215-240; reviewed in Peters et al. (1996) *In Fields Virology*, Third ed. pp. 1161-1176. Fields, B. N., Knipe, D. M., Howley, P. M., et al. eds. Lippincott-Raven Publishers, Philadelphia). The glycoprotein of Ebola virus is unusual in that it is encoded in two open reading frames. Transcriptional editing is needed to express the transmembrane form that is incorporated into the virion (Sanchez et al. (1996) *Proc. Natl. Acad. Sci. USA* 93, 3602-3607; Volchkov et al. (1995) *Virology* 214, 421-430).

[0048] As used herein, “target sequence” refers to a contiguous portion of the nucleotide sequence of an mRNA molecule formed during the transcription of a gene from the Ebola virus, including mRNA that is a product of RNA processing of a primary transcription product.

[0049] As used herein, the term “strand comprising a sequence” refers to an oligonucleotide comprising a chain of nucleotides that is described by the sequence referred to using the standard nucleotide nomenclature.

[0050] As used herein, and unless otherwise indicated, the term “complementary,” when used to describe a first nucleotide sequence in relation to a second nucleotide sequence, refers to the ability of an oligonucleotide or polynucleotide comprising the first nucleotide sequence to hybridize and form a duplex structure under certain conditions with an oligonucleotide or polynucleotide comprising the second nucleotide sequence, as will be understood by the skilled person. Such conditions can, for example, be stringent conditions, where stringent conditions may include: 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50° C. or 70° C. for 12-16 hours followed by washing. Other conditions, such as physiologically relevant conditions as may be encountered inside an organism, can apply. The skilled person will be able to determine the set of conditions most appropriate for a test of complementarity of two sequences in accordance with the ultimate application of the hybridized nucleotides.

[0051] This includes base-pairing of the oligonucleotide or polynucleotide comprising the first nucleotide sequence to the oligonucleotide or polynucleotide comprising the second nucleotide sequence over the entire length of the first and second nucleotide sequence. Such sequences can be referred to as “fully complementary” with respect to each other herein. However, where a first sequence is referred to as “substantially complementary” with respect to a second sequence

herein, the two sequences can be fully complementary, or they may form one or more, but generally not more than 4, 3 or 2 mismatched base pairs upon hybridization, while retaining the ability to hybridize under the conditions most relevant to their ultimate application. However, where two oligonucleotides are designed to form, upon hybridization, one or more single stranded overhangs, such overhangs shall not be regarded as mismatches with regard to the determination of complementarity. For example, a dsRNA comprising one oligonucleotide 21 nucleotides in length and another oligonucleotide 23 nucleotides in length, wherein the longer oligonucleotide comprises a sequence of 21 nucleotides that is fully complementary to the shorter oligonucleotide, may yet be referred to as “fully complementary” for the purposes of the invention.

[0052] “Complementary” sequences, as used herein, may also include, or be formed entirely from, non-Watson-Crick base pairs and/or base pairs formed from non-natural and modified nucleotides, in as far as the above requirements with respect to their ability to hybridize are fulfilled.

[0053] The terms “complementary”, “fully complementary” and “substantially complementary” herein may be used with respect to the base matching between the sense strand and the antisense strand of a dsRNA, or between the antisense strand of a dsRNA and a target sequence, as will be understood from the context of their use.

[0054] As used herein, a polynucleotide which is “substantially complementary to at least part of” a messenger RNA (mRNA) refers to a polynucleotide which is substantially complementary to a contiguous portion of the mRNA of interest (e.g., encoding Ebola). For example, a polynucleotide is complementary to at least a part of a Ebola mRNA if the sequence is substantially complementary to a non-interrupted portion of a mRNA encoding Ebola.

[0055] The term “double-stranded RNA” or “dsRNA,” as used herein, refers to a complex of ribonucleic acid molecules, having a duplex structure comprising two anti-parallel and substantially complementary, as defined above, nucleic acid strands. The two strands forming the duplex structure may be different portions of one larger RNA molecule, or they may be separate RNA molecules. Where the two strands are part of one larger molecule, and therefore are connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5' end of the respective other strand forming the duplex structure, the connecting RNA chain is referred to as a “hairpin loop”. Where the two strands are connected covalently by means other than an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5' end of the respective other strand forming the duplex structure, the connecting structure is referred to as a “linker”. The RNA strands may have the same or a different number of nucleotides. The maximum number of base pairs is the number of nucleotides in the shortest strand of the dsRNA minus any overhangs that are present in the duplex. In addition to the duplex structure, a dsRNA may comprise one or more nucleotide overhangs. dsRNAs as used herein are also referred to as “siRNAs” (short interfering RNAs).

[0056] As used herein, a “nucleotide overhang” refers to the unpaired nucleotide or nucleotides that protrude from the duplex structure of a dsRNA when a 3'-end of one strand of the dsRNA extends beyond the 5'-end of the other strand, or vice versa. “Blunt” or “blunt end” means that there are no unpaired nucleotides at that end of the dsRNA, i.e., no nucleotide overhang. A “blunt ended” dsRNA is a dsRNA that is

double-stranded over its entire length, i.e., no nucleotide overhang at either end of the molecule.

[0057] The term “antisense strand” refers to the strand of a dsRNA which includes a region that is substantially complementary to a target sequence. As used herein, the term “region of complementarity” refers to the region on the antisense strand that is substantially complementary to a sequence, for example a target sequence, as defined herein. Where the region of complementarity is not fully complementary to the target sequence, the mismatches are most tolerated in the terminal regions and, if present, are generally in a terminal region or regions, e.g., within 6, 5, 4, 3, or 2 nucleotides of the 5' and/or 3' terminus.

[0058] The term “sense strand,” as used herein, refers to the strand of a dsRNA that includes a region that is substantially complementary to a region of the antisense strand.

[0059] “Introducing into a cell”, when referring to a dsRNA, means facilitating uptake or absorption into the cell, as is understood by those skilled in the art. Absorption or uptake of dsRNA can occur through unaided diffusive or active cellular processes, or by auxiliary agents or devices. The meaning of this term is not limited to cells in vitro; a dsRNA may also be “introduced into a cell”, wherein the cell is part of a living organism. In such instance, introduction into the cell will include the delivery to the organism. For example, for in vivo delivery, dsRNA can be injected into a tissue site or administered systemically. In vitro introduction into a cell includes methods known in the art such as electroporation and lipofection.

[0060] The terms “silence” and “inhibit the expression of”, in as far as they refer to a gene from the Ebola virus, herein refer to the at least partial suppression of the expression of a gene from the Ebola virus, as manifested by a reduction of the amount of mRNA transcribed from a gene from the Ebola virus which may be isolated from a first cell or group of cells in which a gene from the Ebola virus is transcribed and which has or have been treated such that the expression of a gene from the Ebola virus is inhibited, as compared to a second cell or group of cells substantially identical to the first cell or group of cells but which has or have not been so treated (control cells). The degree of inhibition is usually expressed in terms of

$$\frac{(mRNA \text{ in control cells}) - (mRNA \text{ in treated cells})}{(mRNA \text{ in control cells})} \cdot 100\%$$

[0061] Alternatively, the degree of inhibition may be given in terms of a reduction of a parameter that is functionally linked to Ebola genome transcription, e.g. the amount of protein encoded by a gene from the Ebola virus, or the number of cells displaying a certain phenotype, e.g infection with the Ebola virus. In principle, Ebola genome silencing may be determined in any cell expressing the target, either constitutively or by genomic engineering, and by any appropriate assay. However, when a reference is needed in order to determine whether a given dsRNA inhibits the expression of a gene from the Ebola virus by a certain degree and therefore is encompassed by the instant invention, the assay provided in the Examples below shall serve as such reference.

[0062] For example, in certain instances, expression of a gene from the Ebola virus is suppressed by at least about 20%, 25%, 35%, or 50% by administration of the double-stranded oligonucleotide of the invention. In some embodiment, a gene

from the Ebola virus is suppressed by at least about 60%, 70%, or 80% by administration of the double-stranded oligonucleotide of the invention. In some embodiments, a gene from the Ebola virus is suppressed by at least about 85%, 90%, or 95% by administration of the double-stranded oligonucleotide of the invention.

[0063] As used herein in the context of Ebola expression, the terms “treat”, “treatment”, and the like, refer to relief from or alleviation of pathological processes mediated by Ebola infection. In the context of the present invention insofar as it relates to any of the other conditions recited herein below (other than pathological processes mediated by Ebola expression), the terms “treat”, “treatment”, and the like mean to relieve or alleviate at least one symptom associated with such condition, or to slow or reverse the progression of such condition, or to reduce the amount of virus present in the infected subject.

[0064] As used herein, the phrases “therapeutically effective amount” and “prophylactically effective amount” refer to an amount that provides a therapeutic benefit in the treatment, prevention, or management of pathological processes mediated by Ebola infection or an overt symptom of pathological processes mediated by Ebola expression or the amount virus present in the patient. The specific amount that is therapeutically effective can be readily determined by ordinary medical practitioner, and may vary depending on factors known in the art, such as, e.g. the type of pathological processes mediated by Ebola infection, the patient’s history and age, the stage of pathological processes mediated by Ebola infection, and the administration of other anti-pathological processes mediated by Ebola infection.

[0065] As used herein, a “pharmaceutical composition” comprises a pharmacologically effective amount of a dsRNA and a pharmaceutically acceptable carrier. As used herein, “pharmacologically effective amount,” “therapeutically effective amount” or simply “effective amount” refers to that amount of an RNA effective to produce the intended pharmacological, therapeutic or preventive result. For example, if a given clinical treatment is considered effective when there is at least a 25% reduction in a measurable parameter associated with a disease or disorder, a therapeutically effective amount of a drug for the treatment of that disease or disorder is the amount necessary to effect at least a 25% reduction in that parameter. Further, the pharmaceutical composition can be designed to enhance targeting cells involved in Ebola infection such as dendritic cells, macrophages, hepatocytes, and other parenchymal cells.

[0066] The term “pharmaceutically acceptable carrier” refers to a carrier for administration of a therapeutic agent. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The term specifically excludes cell culture medium. For drugs administered orally, pharmaceutically acceptable carriers include, but are not limited to pharmaceutically acceptable excipients such as inert diluents, disintegrating agents, binding agents, lubricating agents, sweetening agents, flavoring agents, coloring agents and preservatives. Suitable inert diluents include sodium and calcium carbonate, sodium and calcium phosphate, and lactose, while corn starch and alginic acid are suitable disintegrating agents. Binding agents may include starch and gelatin, while the lubricating agent, if present, will generally be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with

a material such as glyceryl monostearate or glyceryl distearate, to delay absorption in the gastrointestinal tract.

[0067] As used herein, a “transformed cell” is a cell into which a vector has been introduced from which a dsRNA molecule may be expressed.

[0068] II. Double-Stranded Ribonucleic Acid (dsRNA)

[0069] In one embodiment, the invention provides double-stranded ribonucleic acid (dsRNA) molecules for inhibiting the expression of a gene from the Ebola virus in a cell or mammal, wherein the dsRNA comprises an antisense strand comprising a region of complementarity which is complementary to at least a part of an mRNA formed in the expression of a gene from the Ebola virus, and wherein the region of complementarity is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and wherein said dsRNA, upon contact with a cell expressing the gene from the Ebola virus, inhibits the expression of the Ebola virus gene by at least 40%.

[0070] The dsRNA comprises two RNA strands that are sufficiently complementary to hybridize to form a duplex structure. One strand of the dsRNA (the antisense strand) comprises a region of complementarity that is substantially complementary, and generally fully complementary, to a target sequence, derived from the sequence of an mRNA formed during the expression of a gene from the Ebola virus, the other strand (the sense strand) comprises a region which is complementary to the antisense strand, such that the two strands hybridize and form a duplex structure when combined under suitable conditions. Generally, the duplex structure is between 15 and 30, more generally between 18 and 25, yet more generally between 19 and 24, and most generally between 19 and 21 base pairs in length. Similarly, the region of complementarity to the target sequence is between 15 and 30, more generally between 18 and 25, yet more generally between 19 and 24, and most generally between 19 and 21 nucleotides in length. The dsRNA of the invention may further comprise one or more single-stranded nucleotide overhang(s).

[0071] The dsRNA can be synthesized by standard methods known in the art as further discussed below, e.g., by use of an automated DNA synthesizer, such as are commercially available from, for example, Biosearch, Applied Biosystems, Inc. In one embodiment, a gene from the Ebola virus is the from human Ebola genome. In specific embodiments, the antisense strand of the dsRNA comprises the sense sequences of Table 2 and the second sequence is selected from the group consisting of the antisense sequences of Table 2. Alternative antisense agents that target elsewhere in the target sequence provided in Table 2 can readily be determined using the target sequence and the flanking Ebola sequence.

[0072] In further embodiments, the dsRNA comprises at least one nucleotide sequence selected from the groups of sequences provided in Table 2. In other embodiments, the dsRNA comprises at least two sequences selected from this group, wherein one of the at least two sequences is complementary to another of the at least two sequences, and one of the at least two sequences is substantially complementary to a sequence of an mRNA generated in the expression of a gene from the Ebola virus. Generally, the dsRNA comprises two oligonucleotides, wherein one oligonucleotide is described as the sense strand in Table 2 and the second oligonucleotide is described as the antisense strand in Table 2.

[0073] The skilled person is well aware that dsRNAs comprising a duplex structure of between 20 and 23, but specifi-

cally 21, base pairs have been hailed as particularly effective in inducing RNA interference (Elbashir et al., EMBO 2001, 20:6877-6888). However, others have found that shorter or longer dsRNAs can be effective as well. In the embodiments described above, by virtue of the nature of the oligonucleotide sequences provided in Table 2, the dsRNAs of the invention can comprise at least one strand of a length of minimally 21 nucleotides. It can be reasonably expected that shorter dsRNAs comprising one of the sequences of Table 2 minus only a few nucleotides on one or both ends may be similarly effective as compared to the dsRNAs described above. Hence, dsRNAs comprising a partial sequence of at least 15, 16, 17, 18, 19, 20, or more contiguous nucleotides from one of the sequences of Table 2, and differing in their ability to inhibit the expression of a gene from the Ebola virus in a FACS assay as described herein below by not more than 5, 10, 15, 20, 25, or 30% inhibition from a dsRNA comprising the full sequence, are contemplated by the invention. Further dsRNAs that cleave within the target sequence provided in Table 2 can readily be made using the Ebola virus sequence and the target sequence provided.

[0074] In addition, the RNAi agents provided in Table 2 identify a site in the Ebola virus mRNA that is susceptible to RNAi based cleavage. As such the present invention further includes RNAi agents that target within the sequence targeted by one of the agents of the present invention. As used herein a second RNAi agent is said to target within the sequence of a first RNAi agent if the second RNAi agent cleaves the message anywhere within the mRNA that is complementary to the antisense strand of the first RNAi agent. Such a second agent will generally consist of at least 15 contiguous nucleotides from one of the sequences provided in Table 2 coupled to additional nucleotide sequences taken from the region contiguous to the selected sequence in a gene from the Ebola virus. For example, the last 15 nucleotides of SEQ ID NO:1 combined with the next 6 nucleotides from the target Ebola genome produces a single strand agent of 21 nucleotides that is based on one of the sequences provided in Table 2.

[0075] The dsRNA of the invention can contain one or more mismatches to the target sequence. In one embodiment, the dsRNA of the invention contains no more than 3 mismatches. If the antisense strand of the dsRNA contains mismatches to a target sequence, it is preferable that the area of mismatch not be located in the center of the region of complementarity. If the antisense strand of the dsRNA contains mismatches to the target sequence, it is preferable that the mismatch be restricted to 5 nucleotides from either end, for example 5, 4, 3, 2, or 1 nucleotide from either the 5' or 3' end of the region of complementarity. For example, for a 23 nucleotide dsRNA strand which is complementary to a region of a gene from the Ebola virus, the dsRNA generally does not contain any mismatch within the central 13 nucleotides. The methods described within the invention can be used to determine whether a dsRNA containing a mismatch to a target sequence is effective in inhibiting the expression of a gene from the Ebola virus. Consideration of the efficacy of dsRNAs with mismatches in inhibiting expression of a gene from the Ebola virus is important, especially if the particular region of complementarity in a gene from the Ebola virus is known to have polymorphic sequence variation within the population.

[0076] In one embodiment, at least one end of the dsRNA has a single-stranded nucleotide overhang of 1 to 4, generally 1 or 2 nucleotides. dsRNAs having at least one nucleotide overhang have unexpectedly superior inhibitory properties

than their blunt-ended counterparts. Moreover, the present inventors have discovered that the presence of only one nucleotide overhang strengthens the interference activity of the dsRNA, without affecting its overall stability. dsRNA having only one overhang has proven particularly stable and effective in vivo, as well as in a variety of cells, cell culture mediums, blood, and serum. Generally, the single-stranded overhang is located at the 3'-terminal end of the antisense strand or, alternatively, at the 3'-terminal end of the sense strand. The dsRNA may also have a blunt end, generally located at the 5'-end of the antisense strand. Such dsRNAs have improved stability and inhibitory activity, thus allowing administration at low dosages, i.e., less than 5 mg/kg body weight of the recipient per day. Generally, the antisense strand of the dsRNA has a nucleotide overhang at the 3'-end, and the 5'-end is blunt. In another embodiment, one or more of the nucleotides in the overhang is replaced with a nucleoside thiophosphate.

[0077] In yet another embodiment, the dsRNA is chemically modified to enhance stability. The nucleic acids of the invention may be synthesized and/or modified by methods well established in the art, such as those described in "Current protocols in nucleic acid chemistry", Beaucage, S. L. et al. (Eds.), John Wiley & Sons, Inc., New York, N.Y., USA, which is hereby incorporated herein by reference. Specific examples of dsRNA compounds useful in this invention include dsRNAs containing modified backbones or no natural internucleoside linkages. As defined in this specification, dsRNAs having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified dsRNAs that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

[0078] Preferred modified dsRNA backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those) having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

[0079] Representative U.S. patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,195; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,316; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, each of which is herein incorporated by reference.

[0080] Preferred modified dsRNA backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatoms and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar

portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

[0081] Representative U.S. patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,64,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and, 5,677,439, each of which is herein incorporated by reference.

[0082] In other certain dsRNA mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an dsRNA mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar backbone of an dsRNA is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative U.S. patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., Science, 1991, 254, 1497-1500.

[0083] Other embodiments featured in the invention include dsRNAs with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular —CH.sub.2-NH—CH.sub.2—, —CH.sub.2-N(CH.sub.3)-O—CH.sub.2-[known as a methylene (methylimino) or MMI backbone], —CH.sub.2-O—N(CH.sub.3)-CH.sub.2—, —CH.sub.2-N(CH.sub.3)-N(CH.sub.3)-CH.sub.2— and —N(CH.sub.3)-CH.sub.2-CH.sub.2-[wherein the native phosphodiester backbone is represented as —O—P—O—CH.sub.2—] of the above-referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above-referenced U.S. Pat. No. 5,602,240. Also preferred are dsRNAs having morpholino backbone structures of the above-referenced U.S. Pat. No. 5,034,506.

[0084] Modified dsRNAs may also contain one or more substituted sugar moieties. In some embodiment, the dsRNAs include one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C.sub.1 to C.sub.10 alkyl or C.sub.2 to C.sub.10 alkenyl and alkynyl. Particularly preferred are O[(CH.sub.2).sub.nO].sub.mCH.sub.3, O(CH.sub.2).sub.2.sub.nOCH.sub.3, O(CH.sub.2).sub.2.sub.nNH.sub.2, O(CH.sub.2).sub.2.sub.nCH.sub.3, O(CH.sub.2).sub.2.sub.nONH.sub.2, and O(CH.sub.2).sub.2.sub.nON[(CH.sub.2).sub.nCH.sub.3].sub.2, where n and m are from 1 to about 10. Other preferred dsRNAs comprise one of the following at the 2' position: C.sub.1 to C.sub.10 lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH.sub.3, OCN, Cl, Br, CN, CF.sub.3, OCF.sub.3, SOCH.sub.3, SO.sub.2CH.sub.3, ONO.sub.2, NO.sub.2, N.sub.3, NH.sub.2, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an

intercalator, a group for improving the pharmacokinetic properties of an dsRNA, or a group for improving the pharmacodynamic properties of an dsRNA, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O—CH₂CH₂OCH₂CH₂, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxy-alkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, i.e., a O(CH₂CH₂)₂N(CH₂CH₂)₂ group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O—CH₂CH₂O—CH₂CH₂N(CH₂CH₂)₂, also described in examples hereinbelow.

[0085] Other preferred modifications include 2'-methoxy (2'-OCH₂CH₂), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the dsRNA, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked dsRNAs and the 5' position of 5' terminal nucleotide. DsRNAs may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative U.S. patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

[0086] dsRNAs may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo, particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-daazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J. L., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie, International Edition*, 1991, 30, 613, and those disclosed by Sanghvi, Y. S., Chapter 15, *DsRNA Research and Applications*, pages 289-302, Crooke, S. T. and Lebleu, B., Eds., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and

5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2. degree. C. (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., Eds., *DsRNA Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

[0087] Other nucleotide substitutions, such as "Universal" bases can be incorporated into siRNA duplexes to increase the number of target sequences (or in this case, number of different Ebola strains) any particular siRNA might have complementarity to and activity against. Universal bases are non-canonical synthetic molecules that mimic structures of traditional nucleotides (the genetic building blocks of DNA and RNA). However, instead of selectively pairing according to Watson/Crick rules (A with T or U, C with G), universal bases "stack" equally well with all natural bases. Incorporating universal bases into siRNAs may enable the siRNA to tolerate a mutation at that specific site in its target mRNA. Thus, by decreasing the need for absolute complementarity between siRNA and its mRNA target, universal-base containing siRNAs may be an approach to (1) prevent drug resistance caused by site-specific viral mutations and (2) create siRNAs able to be broadly reactive across viral species with similar, but not absolutely conserved, targets. Among the modifications that can be used as universal bases are: 3-Nitropyrrole, 5-Nitroindole, Imidazole-4-Carboxamide, 2,4-difluorotoluyl, and Inosine.

[0088] Representative U.S. patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Pat. No. 3,687,808, as well as U.S. Pat. Nos. 4,845,205; 5,130,30; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; and 5,681,941, each of which is herein incorporated by reference, and U.S. Pat. No. 5,750,692, also herein incorporated by reference.

[0089] Another modification of the dsRNAs of the invention involves chemically linking to the dsRNA one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the dsRNA. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 199, 86, 6553-6556), cholic acid (Manoharan et al., *Biorg. Med. Chem. Lett.*, 1994 4 1053-1060), a thioether, e.g., beryl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306-309; Manoharan et al., *Biorg. Med. Chem. Lett.*, 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J*, 1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, 1990, 259, 327-330; Svinarchuk et al., *Biochimie*, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycerol-3-Hphosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654), a palmitoyl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carboxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923-937). Preferred

conjugates will assist in targeting cells infected by Ebola virus such as dendritic cells and macrophages which are involved in early stages of infection and hepatocytes and other parenchymal cells which are involved in later phases of the infection. Such conjugates include, but are not limited to, mannose and folate conjugates.

[0090] Representative U.S. patents that teach the preparation of such dsRNA conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, each of which is herein incorporated by reference.

[0091] It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an dsRNA. The present invention also includes dsRNA compounds which are chimeric compounds. "Chimeric" dsRNA compounds or "chimeras," in the context of this invention, are dsRNA compounds, particularly dsRNAs, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an dsRNA compound. These dsRNAs typically contain at least one region wherein the dsRNA is modified so as to confer upon the dsRNA increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the dsRNA may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of dsRNA inhibition of gene expression. Consequently, comparable results can often be obtained with shorter dsRNAs when chimeric dsRNAs are used, compared to phosphorothioate deoxydsRNAs hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

[0092] In certain instances, the dsRNA may be modified by a non-ligand group. A number of non-ligand molecules have been conjugated to dsRNAs in order to enhance the activity, cellular distribution or cellular uptake of the dsRNA, and procedures for performing such conjugations are available in the scientific literature. Such non-ligand moieties have included lipid moieties, such as cholesterol (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86:6553), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4:1053), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660:306; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3:2765), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20:533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10:111; Kabanov et al., *FEBS Lett.*, 1990, 259:327; Svinarchuk et al., *Biochimie*, 1993,

75:49), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36:3651; Shea et al., *Nucl. Acids Res.*, 1990, 18:3777), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14:969), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36:3651), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264:229), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277:923). Representative United States patents that teach the preparation of such dsRNA conjugates have been listed above. Typical conjugation protocols involve the synthesis of dsRNAs bearing an aminolinker at one or more positions of the sequence. The amino group is then reacted with the molecule being conjugated using appropriate coupling or activating reagents. The conjugation reaction may be performed either with the dsRNA still bound to the solid support or following cleavage of the dsRNA in solution phase. Purification of the dsRNA conjugate by HPLC typically affords the pure conjugate.

[0093] Vector Encoded RNAi Agents

[0094] The dsRNA of the invention can also be expressed from recombinant viral vectors intracellularly in vivo. The recombinant viral vectors of the invention comprise sequences encoding the dsRNA of the invention and any suitable promoter for expressing the dsRNA sequences. Suitable promoters include, for example, the U6 or H1 RNA pol III promoter sequences and the cytomegalovirus promoter. Selection of other suitable promoters is within the skill in the art. The recombinant viral vectors of the invention can also comprise inducible or regulatable promoters for expression of the dsRNA in a particular tissue or in a particular intracellular environment. The use of recombinant viral vectors to deliver dsRNA of the invention to cells in vivo is discussed in more detail below.

[0095] dsRNA of the invention can be expressed from a recombinant viral vector either as two separate, complementary RNA molecules, or as a single RNA molecule with two complementary regions.

[0096] Any viral vector capable of accepting the coding sequences for the dsRNA molecule(s) to be expressed can be used, for example vectors derived from adenovirus (AV); adeno-associated virus (AAV); retroviruses (e.g., lentiviruses (LV), Rhabdoviruses, murine leukemia virus); herpes virus, and the like. The tropism of viral vectors can be modified by pseudotyping the vectors with envelope proteins or other surface antigens from other viruses, or by substituting different viral capsid proteins, as appropriate.

[0097] For example, lentiviral vectors of the invention can be pseudotyped with surface proteins from vesicular stomatitis virus (VSV), rabies, Ebola, Mokola, and the like. AAV vectors of the invention can be made to target different cells by engineering the vectors to express different capsid protein serotypes. For example, an AAV vector expressing a serotype 2 capsid on a serotype 2 genome is called AAV 2/2. This serotype 2 capsid gene in the AAV 2/2 vector can be replaced by a serotype 5 capsid gene to produce an AAV 2/5 vector. Techniques for constructing AAV vectors which express different capsid protein serotypes are within the skill in the art; see, e.g., Rabinowitz JE et al. (2002), *J Virol* 76:791-801, the entire disclosure of which is herein incorporated by reference.

[0098] Selection of recombinant viral vectors suitable for use in the invention, methods for inserting nucleic acid

sequences for expressing the dsRNA into the vector, and methods of delivering the viral vector to the cells of interest are within the skill in the art. See, for example, Dornburg R (1995), *Gene Therap.* 2: 301-310; Eglitis M A (1988), *Biotechniques* 6: 608-614; Miller A D (1990), *Hum Gene Therap.* 1: 5-14; Anderson W F (1998), *Nature* 392: 25-30; and Robinson D A et al., *Nat. Genet.* 33: 401-406, the entire disclosures of which are herein incorporated by reference.

[0099] Preferred viral vectors are those derived from AV and AAV. In a particularly preferred embodiment, the dsRNA of the invention is expressed as two separate, complementary single-stranded RNA molecules from a recombinant AAV vector comprising, for example, either the U6 or H1 RNA promoters, or the cytomegalovirus (CMV) promoter.

[0100] A suitable AV vector for expressing the dsRNA of the invention, a method for constructing the recombinant AV vector, and a method for delivering the vector into target cells, are described in Xia H et al. (2002), *Nat. Biotech.* 20: 1006-1010.

[0101] Suitable AAV vectors for expressing the dsRNA of the invention, methods for constructing the recombinant AV vector, and methods for delivering the vectors into target cells are described in Samulski R et al. (1987), *J. Virol.* 61: 3096-3101; Fisher K J et al. (1996), *J. Virol.* 70: 520-532; Samulski R et al. (1989), *J. Virol.* 63: 3822-3826; U.S. Pat. No. 5,252,479; U.S. Pat. No. 5,139,941; International Patent Application No. WO 94/13788; and International Patent Application No. WO 93/24641, the entire disclosures of which are herein incorporated by reference.

[0102] III. Pharmaceutical Compositions Comprising dsRNA

[0103] In one embodiment, the invention provides pharmaceutical compositions comprising a dsRNA, as described herein, and a pharmaceutically acceptable carrier. The pharmaceutical composition comprising the dsRNA is useful for treating a disease or disorder associated with the expression or activity of a gene from the Ebola virus and/or viral infection, such as systemic hemorrhage and multi-organ failure. Such pharmaceutical compositions are formulated based on the mode of delivery. One example is compositions that are formulated for systemic administration via parenteral delivery.

[0104] The pharmaceutical compositions of the invention are administered in dosages sufficient to inhibit expression of a gene from the Ebola virus. A maximum dosage of 5 mg dsRNA per kilogram body weight of recipient per day is sufficient to inhibit or completely suppress expression of a gene from the Ebola virus.

[0105] In general, a suitable dose of dsRNA will be in the range of 0.01 to 20.0 milligrams per kilogram body weight of the recipient per day, and optimally in the range of 0.01 to 3 mg per kilogram body weight per day. The pharmaceutical composition may be administered once daily, or the dsRNA may be administered as two, three, or more sub-doses at appropriate intervals throughout the day or even using continuous infusion or delivery through a controlled release formulation. In that case, the dsRNA contained in each sub-dose must be correspondingly smaller in order to achieve the total daily dosage. The dosage unit can also be compounded for delivery over several days, e.g., using a conventional sustained release formulation which provides sustained release of the dsRNA over a several day period. In this embodiment, the dosage unit contains a corresponding multiple of the daily dose.

[0106] The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a composition can include a single treatment or a series of treatments. Estimates of effective dosages and in vivo half-lives for the individual dsRNAs encompassed by the invention can be made using conventional methodologies or on the basis of in vivo testing using an appropriate animal model, as described elsewhere herein.

[0107] Advances in animal modeling has generated a number of Ebola infection models (mouse, guinea pig, and non-human primate) that reproduce all the major pathologies associated with human Ebola infection (reviewed in Warfield, K. L. et al. (2006) "Chapter 13: Viral Hemorrhagic Fevers" in *Biodefense: Research Methodology and Animal Models*, pages 227-258, J. R. Swearingen, Ed., Taylor & Francis, Boca Raton). Such models are used for in vivo testing of dsRNA, as well as for determining a therapeutically effective dose.

[0108] The present invention also includes pharmaceutical compositions and formulations which include the dsRNA compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical, pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion.

[0109] Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable.

[0110] Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

[0111] Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

[0112] The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

[0113] Liposomes

[0114] There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. Liposomal delivery systems have been used to effectively deliver siRNA in vivo and silence genes in hepatocytes [Zimmermann et al. (2006) *Nature*, 441:111-114]. Such siRNA-liposomal formulations have also been used for therapeutic benefit in animal models of dyslipidemias [Zimmermann et al. (2006) *Nature*, 441:111-114], HBV infection [Morrissey et al. (2005) *Nature Biotech* 23:1002-1007], Ebola infection [Geisbert, et al. (2006) *The Journal of Infectious Diseases*, 193:1650-1657], and rheumatoid arthritis [Khouri et al. (2006) *Arthritis & Rheumatism*, 54:1867-1877]. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

[0115] Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are phagocytosed by macrophages and other cells in vivo.

[0116] Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

[0117] Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes and as the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

[0118] Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

[0119] Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis

[0120] Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable

complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

[0121] Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).

[0122] One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

[0123] Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G_{sub}.M1, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al., *Cancer Research*, 1993, 53, 3765). Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside G_{sub}.M1, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Pat. No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G_{sub}.M1 or a galactocerebroside sulfate ester. U.S. Pat. No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

[0124] Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C_{sub}.sub.1215G, that contains a

PEG moiety. Illum et al. (FEBS Lett., 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Pat. Nos. 4,426,330 and 4,534,899). Klibanov et al. (FEBS Lett., 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (Biochimica et Biophysica Acta, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Pat. Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Pat. No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Pat. No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.). Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Pat. No. 5,540,935 (Miyazaki et al.) and U.S. Pat. No. 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

[0125] Liposomes and other nanoparticles have also been designed which contain specific targeting molecules. Targeting molecules used for siRNA delivery in vivo have included integrin-binding RGD peptides [Schiffelers et al. (2004) *Nucleic Acids Research* 32:e149], anisamide [Li and Huang (2006) *Molecular Pharmaceutics* 3:579-588] and folate [Hulieskovan et al. (2005) *Cancer Research* 65:8984-8992]. For delivery to myeloid and dendritic cells which are presumed to be important in early Ebola infection, incorporation of targeting agents such as mannose and folate into liposomes and nanoparticles may improve both siRNA delivery and therapeutic effect. Mannose-conjugated oligonucleotides have been shown to specifically improve delivery to myeloid cells [Rojanasakul et al. (1997) *Journal of Biological Chemistry* 272:3910-3914; Diebold et al. (2002) *Somatic Cell and Molecular Genetics* 27:65-73] and mannosylated liposomes are effective targeting agents in vivo [Diebold et al. (2002) *Somatic Cell and Molecular Genetics* 27:65-73; Hattori et al. (2006) *Journal Gene Medicine* 8:824-834; Hattori et al. (2006) *Journal of Pharmacology and Experimental Therapeutics* 318:828-834]. Folate conjugation has proven an effective delivery vehicle in a wide variety of contexts [reviewed in Hilgenbrink and Low (2005) *Journal Pharmaceutical Sciences* 94:2135-2146] and incorporation of folate into liposomes is possible using commercially available reagents such as DSPE-PEG (2000) Folate (Avanti Polar Lipids, Alabaster, Ala.). A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Pat. No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an dsRNA RNA. U.S. Pat. No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in

liposomes. WO 97/04787 to Love et al. discloses liposomes comprising dsRNA targeted to the raf gene.

[0126] Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

[0127] Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

[0128] If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

[0129] If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

[0130] If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

[0131] If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

[0132] The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

[0133] Agents that enhance uptake of dsRNAs at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Pat. No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of dsRNAs.

[0134] Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

[0135] Carriers

[0136] Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity per se) but is recognized as a nucleic acid by in vivo processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate dsRNA in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., DsRNA Res. Dev., 1995, 5, 115-121; Takakura et al., DsRNA & Nucl. Acid Drug Dev., 1996, 6, 177-183).

[0137] Excipients

[0138] In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

[0139] Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to

formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

[0140] Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

[0141] Other Components

[0142] The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

[0143] Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

[0144] Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycytosine, 4-hydroxyperoxycyclophosphoramide, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUDR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, The Merck Manual of Diagnosis and Therapy, 15th Ed. 1987, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents may be used individually

(e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, The Merck Manual of Diagnosis and Therapy, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

[0145] Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit high therapeutic indices are preferred.

[0146] The data obtained from cell culture assays and animal studies can be used in formulation a range of dosage for use in humans. The dosage of compositions of the invention lies generally within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range of the compound or, when appropriate, of the polypeptide product of a target sequence (e.g., achieving a decreased concentration of the polypeptide) that includes the IC50 (i.e., the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0147] In addition to their administration individually or as a plurality, as discussed above, the dsRNAs of the invention can be administered in combination with other known agents effective in treatment of pathological processes mediated by Ebola expression. In any event, the administering physician can adjust the amount and timing of dsRNA administration on the basis of results observed using standard measures of efficacy known in the art or described herein.

[0148] Methods for Treating Diseases Caused by Infection with the Ebola Virus

[0149] The invention relates in particular to the use of a dsRNA or a pharmaceutical composition prepared therefrom for the treatment or prevention of pathological conditions associated with Ebola infection, e.g., systemic hemorrhage and multi-organ failure.

[0150] The invention furthermore relates to the use of an dsRNA or a pharmaceutical composition thereof for treating systemic hemorrhage and multi-organ failure in combination with other pharmaceuticals and/or other therapeutic methods, e.g., with known pharmaceuticals and/or known therapeutic methods, such as, for example, those which are currently employed for treating viral infection and systemic hemorrhage. Preference is given to a combination with interferon or other antiviral agents.

[0151] Methods for Inhibiting Expression of a Gene from the Ebola Virus

[0152] In yet another aspect, the invention provides a method for inhibiting the expression of a gene from the Ebola virus in a mammal. The method comprises administering a composition of the invention to the mammal such that expression of the target Ebola genome is silenced. Because of their high specificity, the dsRNAs of the invention specifically target RNAs (primary or processed) of the target Ebola gene. Compositions and methods for inhibiting the expression of these Ebola genes using dsRNAs can be performed as described elsewhere herein.

[0153] In one embodiment, the method comprises administering a composition comprising a dsRNA, wherein the dsRNA comprises a nucleotide sequence which is complementary to at least a part of an RNA transcript of a gene from the Ebola virus, to the mammal to be treated. When the organism to be treated is a mammal such as a human, the composition may be administered by any means known in the art including, but not limited to oral or parenteral routes, including intravenous, intraperitoneal, intramuscular, subcutaneous, transdermal, airway (aerosol), nasal, administration. In some embodiments, the compositions are administered by intravenous infusion or injection.

[0154] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

EXAMPLES

Example 1

dsRNA Synthesis

[0155] Source of Reagents

[0156] Where the source of a reagent is not specifically given herein, such reagent may be obtained from any supplier of reagents for molecular biology at a quality/purity standard for application in molecular biology.

[0157] siRNA Synthesis

[0158] Single-stranded RNAs were produced by solid phase synthesis on a scale of 1 μ mole using an Expedite 8909


synthesizer (Applied Biosystems, Applera Deutschland GmbH, Darmstadt, Germany) and controlled pore glass (CPG, 500 Å, Proligo Biochemie GmbH, Hamburg, Germany) as solid support. RNA and RNA containing 2'-O-methyl nucleotides were generated by solid phase synthesis employing the corresponding phosphoramidites and 2'-O-methyl phosphoramidites, respectively (Proligo Biochemie GmbH, Hamburg, Germany). These building blocks were incorporated at selected sites within the sequence of the oligoribonucleotide chain using standard nucleoside phosphoramidite chemistry such as described in Current protocols in nucleic acid chemistry, Beaucage, S. L. et al. (Edrs.), John Wiley & Sons, Inc., New York, N.Y., USA. Phosphorothioate linkages were introduced by replacement of the iodine oxidizer solution with a solution of the Beaucage reagent (Chruachem Ltd, Glasgow, UK) in acetonitrile (1%). Further ancillary reagents were obtained from Mallinckrodt Baker (Griesheim, Germany).

[0159] Deprotection and purification of the crude oligoribonucleotides by anion exchange HPLC were carried out according to established procedures. Yields and concentrations were determined by UV absorption of a solution of the respective RNA at a wavelength of 260 nm using a spectral photometer (DU 640B, Beckman Coulter GmbH, Unterschleißheim, Germany). Double stranded RNA was generated by mixing an equimolar solution of complementary strands in annealing buffer (20 mM sodium phosphate, pH 6.8; 100 mM sodium chloride), heated in a water bath at 85-90° C. for 3 minutes and cooled to room temperature over a period of 3-4 hours. The annealed RNA solution was stored at -20° C. until use.

[0160] For the synthesis of 3'-cholesterol-conjugated siRNAs (herein referred to as -Chol-3'), an appropriately modified solid support was used for RNA synthesis. The modified solid support was prepared as follows:

Diethyl-2-azabutane-1,4-dicarboxylate AA


[0161]



[0162] A 4.7 M aqueous solution of sodium hydroxide (50 mL) was added into a stirred, ice-cooled solution of ethyl glycinate hydrochloride (32.19 g, 0.23 mole) in water (50 mL). Then, ethyl acrylate (23.1 g, 0.23 mole) was added and the mixture was stirred at room temperature until completion of the reaction was ascertained by TLC. After 19 h the solution was partitioned with dichloromethane (3×100 mL). The organic layer was dried with anhydrous sodium sulfate, filtered and evaporated. The residue was distilled to afford AA (28.8 g, 61%).

3-{[Ethoxycarbonylmethyl-[6-(9H-fluoren-9-yl-methoxycarbonylamino)-hexanoyl]-amino]-propionic acid ethyl ester AB


[0163]



[0164] Fmoc-6-amino-hexanoic acid (9.12 g, 25.83 mmol) was dissolved in dichloromethane (50 mL) and cooled with ice. Diisopropylcarbodiimide (3.25 g, 3.99 mL, 25.83 mmol) was added to the solution at 0° C. It was then followed by the addition of Diethyl-azabutane-1,4-dicarboxylate (5 g, 24.6 mmol) and dimethylamino pyridine (0.305 g, 2.5 mmol). The solution was brought to room temperature and stirred further for 6 h. Completion of the reaction was ascertained by TLC. The reaction mixture was concentrated under vacuum and ethyl acetate was added to precipitate diisopropyl urea. The suspension was filtered. The filtrate was washed with 5% aqueous hydrochloric acid, 5% sodium bicarbonate and water. The combined organic layer was dried over sodium sulfate and concentrated to give the crude product which was purified by column chromatography (50% EtOAc/Hexanes) to yield 11.87 g (88%) of AB.

3-[(6-Amino-hexanoyl)-ethoxycarbonylmethyl-amino]-propionic acid ethyl ester AC


[0165]



[0166] 3-{[Ethoxycarbonylmethyl-[6-(9H-fluoren-9-yl-methoxycarbonylamino)-hexanoyl]-amino]-propionic acid ethyl ester AB (11.5 g, 21.3 mmol) was dissolved in 20% piperidine in dimethylformamide at 0° C. The solution was continued stirring for 1 h. The reaction mixture was concentrated under vacuum, water was added to the residue, and the product was extracted with ethyl acetate. The crude product was purified by conversion into its hydrochloride salt.

3-({6-[17-(1,5-Dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-ylloxycarbonylamino]-hexanoyl}ethoxycarbonylmethyl-amino)-propionic acid ethyl ester AD

[0167]




[0168] The hydrochloride salt of 3-[(6-Amino-hexanoyl)-ethoxycarbonylmethyl-amino]-propionic acid ethyl ester AC (4.7 g, 14.8 mmol) was taken up in dichloromethane. The suspension was cooled to 0° C. on ice. To the suspension diisopropylethylamine (3.87 g, 5.2 mL, 30 mmol) was added. To the resulting solution cholesteryl chloroformate (6.675 g, 14.8 mmol) was added. The reaction mixture was stirred overnight. The reaction mixture was diluted with dichloromethane and washed with 10% hydrochloric acid. The product was purified by flash chromatography (10.3 g, 92%).

1-{6-[17-(1,5-Dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-ylloxycarbonylamino]-hexanoyl}-4-oxo-pyrrolidine-3-carboxylic acid ethyl ester AE


[0169]

[0170] Potassium t-butoxide (1.1 g, 9.8 mmol) was slurried in 30 mL of dry toluene. The mixture was cooled to 0° C. on ice and 5 g (6.6 mmol) of diester AD was added slowly with stirring within 20 mins. The temperature was kept below 5° C. during the addition. The stirring was continued for 30 mins at 0° C. and 1 mL of glacial acetic acid was added, immediately followed by 4 g of NaH₂PO₄·H₂O in 40 mL of water. The resultant mixture was extracted twice with 100 mL of dichloromethane each and the combined organic extracts were washed twice with 10 mL of phosphate buffer each, dried, and evaporated to dryness. The residue was dissolved in 60 mL of toluene, cooled to 0° C. and extracted with three 50 mL portions of cold pH 9.5 carbonate buffer. The aqueous extracts were adjusted to pH 3 with phosphoric acid, and extracted with five 40 mL portions of chloroform which were combined, dried and evaporated to dryness. The residue was purified by column chromatography using 25% ethylacetate/hexane to afford 1.9 g of b-ketoester (39%).



[6-(3-Hydroxy-4-hydroxymethyl-pyrrolidin-1-yl)-6-oxo-hexyl]-carbamic acid 17-(1,5-dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yl ester AF


[0171]



[0172] Methanol (2 mL) was added dropwise over a period of 1 h to a refluxing mixture of b-ketoester AE (1.5 g, 2.2 mmol) and sodium borohydride (0.226 g, 6 mmol) in tetrahydrofuran (10 mL). Stirring was continued at reflux temperature for 1 h. After cooling to room temperature, 1 N HCl (12.5 mL) was added, the mixture was extracted with ethylacetate (3×40 mL). The combined ethylacetate layer was dried over anhydrous sodium sulfate and concentrated under vacuum to yield the product which was purified by column chromatography (10% MeOH/CHCl₃) (89%).

(6-{3-[Bis-(4-methoxy-phenyl)-phenyl-methoxymethyl]-4-hydroxy-pyrrolidin-1-yl}-6-oxo-hexyl)-carbamic acid 17-(1,5-dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yl ester AG

[0173]




[0174] Diol AF (1.25 gm 1.994 mmol) was dried by evaporating with pyridine (2x5 mL) in vacuo. Anhydrous pyridine (10 mL) and 4,4'-dimethoxytritylchloride (0.724 g, 2.13 mmol) were added with stirring. The reaction was carried out at room temperature overnight. The reaction was quenched by the addition of methanol. The reaction mixture was concentrated under vacuum and to the residue dichloromethane (50 mL) was added. The organic layer was washed with 1M aqueous sodium bicarbonate. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The residual pyridine was removed by evaporating with toluene.

The crude product was purified by column chromatography (2% MeOH/Chloroform, Rf=0.5 in 5% MeOH/CHCl₃) (1.75 g, 95%).

Succinic acid mono-(4-[bis-(4-methoxy-phenyl)-phenyl-methoxymethyl]-1-{6-[17-(1,5 -dimethyl-hexyl)-10,13-dimethyl 2,3,4,7,8,9,10,11,12,13,14,15, 16,17-tetradecahydro-1H cyclopenta[a]phenanthren-3-yloxy-carbonylamino]-hexanoyl}-pyrrolidin-3-yl) ester AH

[0175]

AH




[0176] Compound AG (1.0 g, 1.05 mmol) was mixed with succinic anhydride (0.150 g, 1.5 mmol) and DMAP (0.073 g, 0.6 mmol) and dried in a vacuum at 40° C. overnight. The mixture was dissolved in anhydrous dichloroethane (3 mL), triethylamine (0.318 g, 0.440 mL, 3.15 mmol) was added and the solution was stirred at room temperature under argon atmosphere for 16 h. It was then diluted with dichloromethane (40 mL) and washed with ice cold aqueous citric acid (5 wt %, 30 mL) and water (2x20 mL). The organic phase was dried over anhydrous sodium sulfate and concentrated to dryness. The residue was used as such for the next step.

Cholesterol derivatised CPG AI

[0177]

AI



[0178] Succinate AH (0.254 g, 0.242 mmol) was dissolved in a mixture of dichloromethane/acetonitrile (3:2, 3 mL). To that solution DMAP (0.0296 g, 0.242 mmol) in acetonitrile (1.25 mL), 2,2'-Dithio-bis(5-nitropyridine) (0.075 g, 0.242 mmol) in acetonitrile/dichloroethane (3:1, 1.25 mL) were added successively. To the resulting solution triphenylphosphine (0.064 g, 0.242 mmol) in acetonitrile (0.6 mL) was added. The reaction mixture turned bright orange in color. The solution was agitated briefly using a wrist-action shaker (5 mins). Long chain alkyl amine-CPG (LCAA-CPG) (1.5 g, 61 mM) was added. The suspension was agitated for 2 h. The CPG was filtered through a sintered funnel and washed with acetonitrile, dichloromethane and ether successively. Unreacted amino groups were masked using acetic anhydride/pyridine. The achieved loading of the CPG was measured by taking UV measurement (37 mM/g).

[0179] The synthesis of siRNAs bearing a 5'-12-dodecanoic acid bisdecylamide group (herein referred to as "5'-C32-") or a 5'-cholesteryl derivative group (herein referred to as "5'-Chol-") was performed as described in WO 2004/065601, except that, for the cholesteryl derivative, the oxidation step was performed using the Beaucage reagent in order to introduce a phosphorothioate linkage at the 5'-end of the nucleic acid oligomer.

[0180] Nucleic acid sequences are represented below using standard nomenclature, and specifically the abbreviations of Table 1.

TABLE 1

Abbreviations of nucleotide monomers used in nucleic acid sequence representation. It will be understood that these monomers, when present in an oligonucleotide, are mutually linked by 5'-3'-phosphodiester bonds.	
Abbreviation	Nucleotide(s)
A	adenosine-5'-phosphate
C	cytidine-5'-phosphate
G	guanosine-5'-phosphate
T	2'-deoxy-thymidine-5'-phosphate
U	uridine-5'-phosphate
N	any nucleotide (G, A, C, or T)
a	2'-O-methyladenosine-5'-phosphate
c	2'-O-methylcytidine-5'-phosphate
g	2'-O-methylguanosine-5'-phosphate
u	2'-O-methyluridine-5'-phosphate
sT	2'-deoxy-thymidine-5'-phosphate-phosphorothioate

Example 2

Gene Walking of a Gene from the Ebola Virus

[0181] Design and in silico Selection of siRNAs Targeting Ebola Virus

[0182] siRNA design was carried out to identify siRNAs targeting Ebola virus mRNAs for genes VP30, VP35, NP, L, VP24, VP40 and GP with a focus on sequences isolated from the Zaire region (EBOV-Z), as well as sequences from Sudan (EBOV-S). The siRNA in silico selection resulted in 521 siRNAs satisfying our selection criteria (Table 2).

[0183] Ebola Zaire sequence AY354458 was downloaded from NCBI Nucleotide database and further on used as reference sequence for EBOV-Z. Ebola Sudan sequence AY729654 was used as reference sequence for EBOV-S, respectively.

[0184] Sequence regions encoding target genes VP30, VP35, NP, L, VP24, VP40 and GP according to the information in the Genbank file were extracted from both reference sequences, followed by extraction of all possible 19mers for each gene, resulting in the candidate siRNA target regions (and siRNA sense sequences) for each distinct gene.

[0185] In order to identify siRNAs targeting all available EBOV-Z and EBOV-S sequences, it was necessary to compile available Ebola sequences from all sequenced isolates. For this, each of the 7 target gene sequences extracted from the Ebola Zaire reference sequences was used in a blast search against viruses at NCBI with default parameters and resulting Ebola mRNA hits were downloaded.

[0186] Each candidate target region was tested for conservation across the Ebola sequences by searching the relevant target gene for presence of the 19mer target region. The percentage of conserved sequences across all downloaded sequences for the relevant gene was calculated for each candidate target region by dividing of the number of conserved sequences with the total number of downloaded sequences.

Example 3

Ebola siRNA in vitro Screening

[0187] Table 3 provides a summary of the screening results of the siRNAs described in Table 2. Following initial screening using a GFP-expressing Ebola-Zaire virus and immunofluorescence screening using Ebola-Sudan virus, siRNA showing activity were further tested by plaque assay for antiviral activity against Ebola-Zaire and Ebola-Sudan strains. Several siRNAs were identified that had significant activity against one or more Ebola strains. At a concentration of 100 nM many of the siRNA identified showed greater than a 1 log reduction (>90% inhibition) in Ebola virus titers. Negative control luciferase and GFP siRNA at the same concentration showed reductions in virus titer of between 10 and 35% (Table 3). Three previously identified Ebola siRNA (LS L#1, LS NP#1, LS VP35#1) were also tested in parallel and these inhibited Ebola virus by roughly 70%. The previously identified Ebola siRNA are 25 nucleotide blunt-ended duplexes with the following composition:

```

LS L#1, sense:                                     (SEQ ID NO: 1073)
5' CCAUCUCUGAGACACGACAUAUCUU 3'

anti-sense:                                       (SEQ ID NO: 1074)
5' AAGAU AUGUCGUGUCUCAGAGAUGG 3';

LS NP#1, sense:                                   (SEQ ID NO: 1075)
5' GGUUCAAGGCAAAUUAAGUACAU 3'

anti-sense                                       (SEQ ID NO: 1076)
5' AUGUACUUGAAUUUGCCUUUGAACC 3';

LS VP35#1, sense                                (SEQ ID NO: 1077)
5' CCCAAUGCAACAAACCAAGCCAAA 3'

```

-continued

anti-sense
(SEQ ID NO: 1078)
5' UUUGGCUUGGUUUGUUGCAUUUGGG 3'.

The siRNA sequences for AD-1955 and AD-5179
are as follows:
AD-1955, sense:
(SEQ ID NO: 1079)
5' CUUACGUCUGAGUACUUCGAdTsdT 3'

anti-sense:
(SEQ ID NO: 1080)
5' UCGAAGUACUCAGCGUAAGdTsdT 3';

AD-5179, sense:
(SEQ ID NO: 1081)
5' CcAcAuGAGcAGcACGACuS 3'

anti-sense:
(SEQ ID NO: 1082)
5' AAGUCGUGCUGCUUCAUGUGgsusC 3'.

[0188] Lead siRNAs were also screened for immunostimulatory activity (IFN α and TNF α). Immunostimulatory activity was assayed by transfecting siRNAs into human peripheral blood mononuclear cells and measuring cytokine release by ELISA as outlined in Hornung et al. Nature Medicine 2005. Cytokine levels were normalized to a positive control siRNA included in every assay. The lead candidates had no immunostimulatory activity.

[0189] The following procedures were used in generating the screening results.

[0190] GFP Ebola-Zaire Assay

[0191] VERO cells were transfected at $\sim 2 \times 10^4$ cells per well in a black-walled 96 well plate. Transfection was performed in EMEM with 10% FCS overnight at 100, 10 and 1 nM siRNA complexed with lipofectamine (1.2 μ l of lipofectamine per well in 50 μ l volume; complexation was performed at room temperature for 15-20 min).

[0192] Next day cells were infected with GFP-EBOLA virus (1/50 dilution of stock EBOLA-Zaire GFP, stock E6(4) from 11 Oct. 05, USAMRIID) in 50 μ l of EMEM with 10% FCS. 2 days later cells were fixed in 10% neutral-buffered formalin for >3 days. Formalin was changed before removing from BSL-4 suite. Next formalin was replaced with PBS.

[0193] To quantify infection level of cells in individual wells, cells were stained with 10-20 μ l/well of 10 μ g/ml Hoescht dye and read on Discovery 1 microscope. GFP signal normalized to Hoescht signal was read as a measure of infection level.

[0194] Immunofluorescence Ebola-Sudan Assay

[0195] VERO cells were transfected at $\sim 2 \times 10^4$ cell per well in a black-walled 96 well plate. Transfection was performed in EMEM with 10% FCS overnight at 100, 10 and 1 nM siRNA complexed with lipofectamine (1.2 μ l of lipofectamine per well in 50 μ l volume; complexation was performed at room temperature for 15-20 min).

[0196] Next day cells were infected with EBOLA-Sudan virus (1/100 dilution of EBOV-Sudan (Boniface), stock GP(1)V(2)E6(2) from 23 May 06, USAMRIID) in 50 μ l of EMEM with 10% FCS. Two days later cells were fixed in 10% neutral-buffered formalin for >3 days. Formalin was changed before removing from BSL-4 suite. Next formalin was replaced with PBS.

[0197] To detect infected cells, cells were stained for 4 h at room temperature with mouse anti-Sudan Boniface polyclonal sera (sera was collected from 20 animals and pooled at

day 30 post infection of C57BL/6 mice infected with ~ 1000 pfu of the EBOLA SUDAN-BONIFACE, stock GP(1) V(2) V(1) E6(2) from 23 May 06, USAMRIID) at 1:200 dilution in PBS. Then cells were washed with PBS 2 \times for 5 minutes. Goat anti-mouse IgG-AlexaFluor488 (Molecular Probes) was added at 1:500 dilution in PBS. Cells were washed again with PBS for 5 minutes, 100 μ l of PBS was added to each well. To quantify infection level of cells in individual wells, cell were stained with 10-20 μ l/well of 10 μ g/ml Hoescht dye and read on Discovery 1 microscope. AlexaFluor488 signal normalized to Hoescht signal was read as a measure of infection level.

[0198] Plaque Assay for Filoviruses for in vitro Assay

[0199] Vero cells were transfected in 24 well plates at the density of $\sim 1.5 \times 10^5$ /well density. Transfection was performed in EMEM with 10% FCS overnight at 100 nM siRNA complexed with lipofectamine (3 μ l of lipofectamine per well in 200 μ l volume; complexation was performed at room temperature for 15-20 min). Transfection was done in duplicates. 24 hours later duplicate plates were infected in 50 μ l/well with either 1/500 diluted Zaire-EBOV [(E6P2) stock from 20 Jun. 06, USAMRIID] or 1/1000 diluted EBOV-Sudan [(strain Boniface), stock GP(1)V(2)E6(2) from 23 May 06, USAMRIID]. After 1 hour at 37 $^\circ$ C. virus inoculum was replaced with 500 μ l of fresh 10% FCS/EMEM.

[0200] 48-72 h later supernatants were harvested from each well.

[0201] Plaque assay was performed with supernatants at 10^{-1} , 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} and 10^{-6} dilutions. Fresh Vero cells in 6-well plates were infected with diluted supernatants for 1 hour at 37 $^\circ$ C. with rocking plates every 15 minutes; overlaid with 2 ml/well of 0.5% agarose in EMEM, 5% FCS, Pen/Strep. Six days later plates were overlaid with 2 ml of overlay media+4% neutral red solution and read plates the following day.

[0202] siRNA Activity Determination Using the Plasmid System.

[0203] Consensus sequences of NP (SEQ ID NO:1043), GP (SEQ ID NO:1044), L, VP24 (SEQ ID NO:1045), VP30 (SEQ ID NO:1046), VP35 (SEQ ID NO:1047), VP40 (SEQ ID NO:1048) were synthesized by GENEART (Regensburg, Germany) and cloned into GENEART standard vector. The L gene was generated as 2 fragments (SEQ ID NO:1049 and SEQ ID NO:1050). All genes were subcloned into individual psiCheck-2 (Promega, Mannheim, Germany) vectors via XhoI and NotI sites, resulting in a construct with the flu gene between the stop-codon and the polyA-signal of *Renilla* luciferase. Correct cloning was confirmed by end sequencing performed by GATC Biotech (Konstanz, Germany).

[0204] Transfections:

[0205] Cos-7 cells were seeded at 1.5×10^4 cells/well on white 96-well plates with clear bottom (Greiner Bio-One GmbH, Frickenhausen, Germany) in 75 μ l of growth medium. Directly after seeding the cells, 50 ng of plasmid/well were transfected with Lipofectamine2000 (Invitrogen) as described below for the siRNAs, with the plasmid diluted in Opti-MEM to a final volume of 12.5 μ l/well, prepared as a mastermix for the whole plate.

[0206] siRNA transfections were performed in quadruplicates 4 h after plasmid transfection. For each well 0.5 μ l Lipofectamine2000 (Invitrogen GmbH, Karlsruhe, Germany) were mixed with 12 μ l Opti-MEM (Invitrogen) and incubated for 15 min at room temperature. For an siRNA concentration of 50 nM in the 100 μ l transfection volume, 1 μ l

of a 5 μ M siRNA were mixed with 11.5 μ l Opti-MEM per well, combined with the Lipofectamine2000-Opti-MEM mixture and again incubated for 15 minutes at room temperature. During incubation, the growth medium was removed from cells and replaced by 75 μ l/well of fresh medium. siRNA-Lipofectamine2000-complexes were applied completely (25 μ l each per well) to the cells and cells were incubated for 24 h at 37° C. and 5% CO₂ in a humidified incubator (Heraeus GmbH, Hanau, Germany).

[0207] Cells were harvested by removing growth medium and application of 150 μ l of a 1:1 mixture consisting of medium and Dual-Glo Luciferase substrate, from the Dual-Glo Luciferase Assay System (Promega, Mannheim, Germany). The luciferase assay was performed according to the manufacturer's protocol for Dual-Glo Luciferase assay and luminescence was measured in a Victor-Light 1420 Luminescence Counter (Perkin Elmer, Rodgau-Jügesheim, Germany). Values obtained with *Renilla* luciferase were normalized to the respective values obtained with Firefly luciferase. Values acquired with siRNAs directed against an Ebola gene were normalized to the value obtained with an unspecific siRNA (directed against neomycin resistance gene) set to 100%.

Example 4

In vivo Filovirus Infection Model

[0208] Liposome-formulated siRNAs targeting Ebola genes protected mice from a lethal Ebola virus challenge. Details on the liposome formulation are detailed in the next section. Mice were treated with liposome-formulated siRNA (described below) twice, at 2 hours prior to Ebola infection (5 mg/kg i.v.) and at 3 days after Ebola infection (3 mg/kg i.p.). Mice were infected intraperitoneally with 30,000 LD50 of Ebola-Zaire (LD50 is lethal dose of Ebola infection where 50% of animals die). Mice were monitored for survival with n=10 per treatment group. Negative controls included untreated mice and mice treated with liposome-formulated luciferase siRNA (AD-1955). EK1 is a previously published siRNA sequence targeting the Ebola L gene [Geisbert et al. (2006) *The Journal of Infectious Disease* 193:1950-1957] that was used as a positive control. The siRNA sequences for EK1 are as follows:

```
(sense)                               (SEQ ID NO: 1083)
5' GUACGAAGCUGUAUAUAAAdTdT 3'
and
(antisense)                           (SEQ ID NO: 1084)
5' UUUUAUAUACAGCUUCGUACdTdT 3'.
```

[0209] FIG. 1 provides the results. All the negative control-treated animals (untreated and liposomally-formulated luciferase siRNA-treated) died within 6-8 days following Ebola infection. Several of the liposomally-formulated Ebola siRNAs showed significant increases in survival rates compared to the negative controls. Multiple Ebola siRNA (AD-11691, AD-11710, AD-11588, AD-11599, AD-11570) showed more protection against lethal Ebola infection than the previously published EK1 siRNA (FIG. 1).

[0210] A further experiment was conducted utilizing one of the active Ebola siRNA (AD-11570) to investigate different dosing routes and treatment regimens. Mice were treated with

liposome-formulated Ebola VP35 siRNA (AD-11570) or negative control luciferase siRNA (AD-1955) 2 hours prior to Ebola infection (5 mg/kg i.p.). Mice were infected intraperitoneally with 30,000 LD50 of Ebola-Zaire (LD50 is lethal dose of Ebola infection where 50% of animals die). Mice were monitored for survival with n=10 per treatment group.

[0211] FIG. 2 provides the results. The animals treated with liposomally-formulated AD-11570 showed near complete protection against lethal Ebola infection as compared to the negative control-treated animals (untreated and liposomally-formulated luciferase siRNA-treated) (FIG. 2). These results indicate that a single siRNA administration either via intravenous or intraperitoneal route is able to have a significant impact on survival.

[0212] Formulation Procedure

[0213] The lipidoid ND98-4HCl (MW 1487) (FIG. 3), Cholesterol (Sigma-Aldrich), and PEG-Ceramide C16 (Avanti Polar Lipids) were used to prepare lipid-siRNA nanoparticles. Stock solutions of each in ethanol were prepared: ND98, 133 mg/mL; Cholesterol, 25 mg/mL, PEG-Ceramide C16, 100 mg/mL. ND98, Cholesterol, and PEG-Ceramide C16 stock solutions were then combined in a 42:48:10 molar ratio. Combined lipid solution was mixed rapidly with aqueous siRNA (in sodium acetate pH 5) such that the final ethanol concentration was 35-45% and the final sodium acetate concentration was 100-300 mM. Lipid-siRNA nanoparticles formed spontaneously upon mixing. Depending on the desired particle size distribution, the resultant nanoparticle mixture was in some cases extruded through a polycarbonate membrane (100 nm cut-off) using a thermobarrel extruder (Lipex Extruder, Northern Lipids, Inc). In other cases, the extrusion step was omitted. Ethanol removal and simultaneous buffer exchange was accomplished by either dialysis or tangential flow filtration. Buffer was exchanged to phosphate buffered saline (PBS) pH 7.2.

[0214] Characterization of Formulations

[0215] Formulations prepared by either the standard or extrusion-free method are characterized in a similar manner. Formulations are first characterized by visual inspection. They should be whitish translucent solutions free from aggregates or sediment. Particle size and particle size distribution of lipid-nanoparticles are measured by dynamic light scattering using a Malvern Zetasizer Nano ZS (Malvern, USA). Particles should be 20-300 nm, and ideally, 40-100 nm in size. The particle size distribution should be unimodal. The total siRNA concentration in the formulation, as well as the entrapped fraction, is estimated using a dye exclusion assay. A sample of the formulated siRNA is incubated with the RNA-binding dye Ribogreen (Molecular Probes) in the presence or absence of a formulation disrupting surfactant, 0.5% Triton-X100. The total siRNA in the formulation is determined by the signal from the sample containing the surfactant, relative to a standard curve. The entrapped fraction is determined by subtracting the "free" siRNA content (as measured by the signal in the absence of surfactant) from the total siRNA content. Percent entrapped siRNA is typically >85%.

Example 5

siRNAs Targeting Ebola Increased the Life-Span of Mice and Guinea Pigs Infected with Ebola.

[0216] siRNA directed against different Ebola genes were formulated in liposomes. A single dose of siRNA targeting the VP35 gene (AD-11570) protected both mice and guinea

pigs against lethal Ebola infection (1000 PFU; 30000× LD50). Protection was associated with reduction in viral titres and was seen when drug was administered either prophylactically or therapeutically. Irrelevant siRNA (targeting luciferase) similarly formulated showed no protective effect or impact on virus titer.

[0217] Studies were conducted in mouse, guinea pig, and nonhuman primate lethal Ebola infection models. The studies are summarized below.

[0218] Mouse study 1: Demonstrated efficacy in mouse model of Ebola for multiple siRNA sequences formulated in LNP01. siRNAs were administered by intravenous (i.v.) injection on day 0, followed by intraperitoneal (i.p.) injection on day 3. See FIG. 1.

[0219] Mouse study 3: Demonstrated efficacy of siRNA in LNP01 formulation in the mouse model of Ebola when given by either IV or IP route. See FIG. 2. Mice were monitored for survival with n=10 per treatment group.

[0220] Mouse study #14: Demonstrated efficacy of siRNA in DODMA in the mouse model of Ebola by the IP route. See FIG. 4. siRNAs were formulated in DODMA:DSPC:Chol:PEG-DMG. Mice were monitored for survival with n=10 per treatment group. Treatment with 10 mg/kg DODMA-formulated AD-11570 siRNAs was also effective to protect guinea pigs infected with Ebola.

[0221] Mouse study #15: Demonstrated that siRNA in DODMA formulation is effective down to 0.04 mg/kg in the mouse model of Ebola. AD-11570 consistently gave 25-50% protection, but no clear dose response was seen. The control siRNA AD-1955 gave 25-50% protection, but again, no dose response was observed. See FIG. 5.

[0222] Guinea pig study #6: Demonstrated efficacy of multiple doses of siRNA in DODMA formulation in the guinea pig model of Ebola. AD-11570 siRNAs formulated in DODMA:DSPC:Chol:PEG-DMG were effective to protect guinea pigs from Ebola. See FIG. 6.

[0223] Guinea pig study #11: Efficacy of siRNAs formulated with DODMA and targeting different Ebola genes in a guinea pig model of Ebola. See FIG. 7

[0224] A 95% decrease in viral titers was also observed following administration of LNP01 formulated VP35 siRNA to BALB/c mice (n=5 per group) (FIG. 8). Mice were dosed systemically with LNP01 formulated siRNA at 5 mg/kg i.v. at day 0, then 3 mg/kg i.p. at day 3. Two hours post-siRNA injection at day 0, mice were injected with 1,000 pfu Ebola-Zaire virus and monitored for survival. On day 6 post-infection, the mice were sacrificed and their blood viral titers were determined by plaque assay.

[0225] Table 3 shows the results of cell-based and plaque assays.

[0226] Table 4 shows the results of plaque assays for control siRNAs.

[0227] Table 5. shows the sequences of modified duplexes, and Table 6 shows the effect of the modified duplexes on plaque assay activity and IC₅₀ values in the plasmid-based system.

[0228] Table 7 shows the effect of siRNAs on cytokine levels (IFN-alpha and TNF-alpha).

[0229] Table 8 shows the siRNA silencing in the plasmid system and calculated IC₅₀ values.

[0230] Table 9 shows that nonhuman primates administered siRNAs targeting Ebola did not demonstrate a decrease in lymphocyte or platelet count.

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
3-21	VP35	A18480	1	GAUGAAGAUUAAAACCUUCTsT	A184B1	2	GAAGGUUUUAAUCUUCUAUCTsT	AD-11542
4-22	VP35	A18482	3	AUGAAGAUUAAAACCUUCATsT	A18483	4	UGAAGGUUUUAAUCUUCAUTsT	AD-11543
5-23	VP35	A18484	5	UGAAGAUUAAAACCUUCAUTsT	A18485	6	AUGAAGGUUUUAAUCUUCATsT	AD-11544
6-24	VP35	A18486	7	GAAGAUUAAAACCUUCAUCTsT	A18487	8	GAUGAAGGUUUUAAUCUUCTsT	AD-11545
1354-1372	VP35	A18488	9	UGAUGAAGAUUAAAGAAAATsT	A18489	10	UUUUUCUUAACUUCUUCATsT	AD-11546
7-25	VP35	A18490	11	AAGAUUAAAACCUUCAUCATsT	A18491	12	UGAUGAAGGUUUUAAUCUUTsT	AD-11547
8-26	VP35	A18492	13	AGAUUAAAACCUUCAUCAUTsT	A18493	14	AUGAUGAAGGUUUUAAUCUTsT	AD-11548
9-27	VP35	A18494	15	GAUUAAAACCUUCAUCAUCTsT	A18495	16	GAUGAUGAAGGUUUUAAUCTsT	AD-11549
10-28	VP35	A18496	17	AUUAAAACCUUCAUCUCCTsT	A18497	18	GGAUGAUGAAGGUUUUAAUTsT	AD-11550
11-29	VP35	A18498	19	UUAAAACCUUCAUCCUTsT	A18499	20	AGGAUGAUGAAGGUUUUAAATsT	AD-11551
12-30	VP35	A18500	21	UAAAACCUUCAUCCUUTsT	A18501	22	AAGGAUGAUGAAGGUUUUATsT	AD-11552

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1-19	VP30	A18502	23	GAUGAAGAUUAAAGAAAAGTsT	A18503	24	CUUUUUCUUAACUUCUAUCsT	AD-11553
3-21	VP35	A18504	151	GAuGAAGAuAAAACcuucTsT	A18505	152	GAAGGUUUUAACUUCUAUCsT	AD-11554
4-22	VP35	A18506	153	AuGAAGAuAAAACcuucATsT	A18507	154	UGAAGGUUUUAACUUCUAUCsT	AD-11555
5-23	VP35	A18508	155	uGAAGAuAAAACcuucAuTsT	A18509	156	AUGAAGGUUUUAACUUCATsT	AD-11556
6-24	VP35	A18510	157	GAAGAuAAAACcuucAucTsT	A18511	158	GAUGAAGGUUUUAACUUCsT	AD-11557
1354-1372	VP35	A18512	159	uGAuGAAGAuAAGAAAAATsT	A18513	160	UUUUUCUUAACUUCUAUCATsT	AD-11558
7-25	VP35	A18514	161	AAGAuAAAACcuucAucATsT	A18515	162	UGAUGAAGGUUUuAAUCUUsT	AD-11559
8-26	VP35	A18516	163	AGAuAAAACcuucAucAuTsT	A18517	164	AUGAUGAAGGUUUuAAUCUsT	AD-11560
9-27	VP35	A18518	165	GAuuAAAACcuucAucAucTsT	A18519	166	GAUGAUGAAGGUUUuAAUCsT	AD-11561
10-28	VP35	A18520	167	AuuAAAACcuucAucAuccTsT	A18521	168	GGAUGAUGAAGGUUUuAAUsT	AD-11562
11-29	VP35	A18522	169	uuAAAACcuucAucAuccuTsT	A18523	170	AGGAUGAUGAAGGUUUuAATsT	AD-11563
12-30	VP35	A18524	171	uAAAACCuCAuCAuCCuTsT	A18525	172	AAGGAUGAUGAAGGUUUuATsT	AD-11564
1-19	VP30	A18526	173	GAuGAAGAuAAGAAAAGTsT	A18527	174	CUUUUUCUuAAUCUUCUAUCsT	AD-11565
3-21	VP35	A18528	1019	GAuGAAGAuAAAACCUUCsT	A18529	1020	GAAGGuuuUAAUCuUCAUCsT	AD-11566
4-22	VP35	A18530	1021	AuGAAGAuAAAACcuucATsT	A18531	1022	UGAAGGUUUUAACUUCUAUCsT	AD-11567
5-23	VP35	A18532	1023	uGAAGAuAAAACcuucAuTsT	A18533	1024	AuGAAGGuuuuAAUCuUcATsT	AD-11568
6-24	VP35	A18534	1025	GAAGAuAAAACcuucAucTsT	A18535	1026	GAuGAAGGuuuuAAUCuUCsT	AD-11569
1354-1372	VP35	A18536	1027	uGAuGAAGAuAAGAAAAATsT	A18537	1028	UUUUUCuuAAUCuUcAUcATsT	AD-11570
7-25	VP35	A18538	1029	AAGAuAAAACcuucAucATsT	A18539	1030	UGAUGAAGGUUUuAAUCUUsT	AD-11571
8-26	VP35	A18540	1031	AGAuAAAACcuucAucAuTsT	A18541	1032	AuGAuGAAGGuUuAAUCUsT	AD-11572
9-27	VP35	A18542	1033	GAuuAAAACcuucAucAucTsT	A18543	1034	GAuGAuGAAGGuuuuAAUCsT	AD-11573
10-28	VP35	A18544	1035	AuuAAAACcuucAucAuccTsT	A18545	1036	GGAuGAuGAAGGuuuuAAUsT	AD-11574
11-29	VP35	A18546	1037	UuAAAACcuucAucAuccuTsT	A18547	1038	AGGAuGAuGAAGGuuuuAATsT	AD-11575

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
12-30	VP35	A18548	1039	uAAAAccuucAucAuccuuTsT	A18549	1040	AAGGAuGAuGAAGGuuuuATsT	AD-11576
1-19	VP30	A18550	1041	GAuGAAGAuAAGAAAAAGTsT	A18551	1042	CuuuuUCuuAAUCuUcAUCTsT	AD-11577
996-1014	NP	A18552	25	UGGACACAUGAUGGUGAUCTsT	A18553	26	GAUCACCAUCAUGUGUCCATsT	AD-11578
1467-1485	NP	A18554	27	AAGCAACUCCAACAUAUGTsT	A18555	28	CAUAUUGUUGGAGUUGCUUTsT	AD-11579
11-29	VP35	A18556	29	AAAACCUUCAUCAUCCUUUTsT	A18557	30	AAAGGAUGAUGAAGGUUUUTsT	AD-11580
1357-1375	VP35	A18558	31	UUGAUGAAGAUUAAGAAAAATsT	A18559	32	UUUUCUUAUCUUCUCAATsT	AD-11581
1-19	VP40	A18560	33	GAUGAAGAUUAAGAAAAAGTsT	A18561	34	CUUUUUCUUAUCUUCUCAUTsT	AD-11582
647-665	VP40	A18562	35	CCCUGCUGCAACAUGGACATsT	A18563	36	UGUCCAUGUUGCAGCAGGTSsT	AD-11583
2-20	VP30	A18564	37	AUGAAGAUUAAGAAAAAGUTsT	A18565	38	ACUUUUUCUUAUCUUCAUTsT	AD-11584
1-19	L	A18566	39	AAGAUAUAAGAAAAAGUCCATsT	A18567	40	UGGACUUUUUCUUAUCUUTsT	AD-11585
5-23	NP	A18568	41	AAGAUAUAUAUUUCCUCTsT	A18569	42	GAGGAAAAUUAUUAUCUUTsT	AD-11586
823-841	NP	A18570	43	AUGCCGGAAGAGGAGACAATsT	A18571	44	UUGUOICCUUCUCCGGCAUTsT	AD-11587
824-842	NP	A18572	45	UGCCGGAAGAGGAGACAATsT	A18573	46	GUUGUCUCCUUCUCCGGCATsT	AD-11588
987-1005	NP	A18574	47	GCAAUCAGUAGGACACAUGTsT	A18575	48	CAUGUGUCCUACUGAUUGCTsT	AD-11589
988-1006	NP	A18576	49	CAAUCAGUAGGACACAUGATsT	A18577	50	UCAUGUGUCCUACUGAUUGTsT	AD-11590
989-1007	NP	A18578	51	AAUCAGUAGGACACAUGATsT	A18579	52	AUCAUGUGUCCUACUGAUUTsT	AD-11591
990-1008	NP	A18580	53	AUCAGUAGGACACAUGAUGTsT	A18581	54	CAUCAUGUGUCCUACUGATsT	AD-11592
991-1009	NP	A18582	55	UCAGUAGGACACAUGAUGTsT	A18583	56	CCAUCAUGUGUCCUACUGATsT	AD-11593
992-1010	NP	A18584	57	CAGUAGGACACAUGAUGGUTsT	A18585	58	ACCAUCAUGUGUCCUACUGTsT	AD-11594
993-1011	NP	A18586	59	AGUAGGACACAUGAUGGUTsT	A18587	60	CACCAUCAUGUGUCCUACUTsT	AD-11595
994-1012	NP	A18588	61	GUAGGACACAUGAUGGUGATsT	A18589	62	UCACCAUCAUGUGUCCUACTsT	AD-11596
995-1013	NP	A18590	63	UAGGACACAUGAUGGUGATsT	A18591	64	AUCACCAUCAUGUGUCCUATsT	AD-11597
1005-1023	NP	A18592	65	GAUGGUGAUUUUCCGUUUGTsT	A18593	66	CAAACGGAAAAUCACCAUCTsT	AD-11598

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1006-1024	NP	A18594	67	AUGGUGAUUUUCCGUUGATsT	A18595	68	UCAAACGGAAAAUCACCAUTsT	AD-11599
1007-1025	NP	A18596	69	UGGUGAUUUUCCGUUGAUTsT	A18597	70	AUCAACGGAAAAUCACCATsT	AD-11600
1008-1026	NP	A18598	71	GGUGAUUUUCCGUUGAUGTsT	A18599	72	CAUCAACGGAAAAUCACCTsT	AD-11601
1462-1480	NP	A18600	73	GCUGAGAAGCAACUCCAACtsT	A18601	74	GUUGGAGUUGCUUCUCAGTsT	AD-11602
1463-1481	NP	A18602	75	CUGAGAAGCAACUCCAACATsT	A18603	76	UGUUGGAGUUGCUUCUCAGTsT	AD-11603
1464-1482	NP	A18604	77	UGAGAAGCAACUCCAACAATsT	A18605	78	UUGUUGGAGUUGCUUCUCATsT	AD-11604
1465-1483	NP	A18606	79	GAGAAGCAACUCCAACAUTsT	A18607	80	AUUGUUGGAGUUGCUUCUCTsT	AD-11605
1466-1484	NP	A18608	81	AGAAGCAACUCCAACAUAUsT	A18609	82	UAUUGUUGGAGUUGCUUCUTsT	AD-11606
1353-1371	VP35	A18610	83	AAAAGUGAUGAAGAUUAAGTsT	A18611	84	CUUAAUCUUCACACUUUTsT	AD-11607
1354-1372	VP35	A18612	85	AAAGUGAUGAAGAUUAAGATsT	A18613	86	UCUUAUCUUCACACUUUTsT	AD-11608
1355-1373	VP35	A18614	87	AAGUGAUGAAGAUUAAGAATsT	A18615	88	UUCUUAUCUUCACACUUTsT	AD-11609
1356-1374	VP35	A18616	89	AGUGAUGAAGAUUAAGAAATsT	A18617	90	UUUCUUAUCUUCACACUTsT	AD-11610
645-663	VP40	A18618	91	CUGCCUGCUGCAACAUGGATsT	A18619	92	UCCAUGUUGCAGCAGGCAGTsT	AD-11611
646-664	VP40	A18620	93	UGCCUGCUGCAACAUGGACTsT	A18621	94	GUCCAUGUUGCAGCAGGCATsT	AD-11612
451-469	QP	A18622	95	GGCUGAAAACUGCUACAAUsT	A18623	96	AUUGUAGCAGUUUUCAGCCTsT	AD-11613
452-470	GP	A18624	97	GCUGAAAACUGCUACAAUCTsT	A18625	98	GAUUGUAGCAGUUUUCAGTsT	AD-11614
453-471	GP	A18626	99	CUGAAAACUGCUACAAUCUTsT	A18627	100	AGAUGUAGCAGUUUUCAGTsT	AD-11615
454-472	GP	A18628	101	UGAAAACUGCUACAAUCUUTsT	A18629	102	AAGAUUGUAGCAGUUUUCATsT	AD-11616
455-473	GP	A18630	103	GAAAACUGCUACAAUCUUGTsT	A18631	104	CAAGAUUGUAGCAGUUUUCTsT	AD-11617
456-474	GP	A18632	105	AAAACUGCUACAAUCUUGATsT	A18633	106	UCAAGAUUGUAGCAGUUUTsT	AD-11618
457-475	GP	A18634	107	AAACUGCUACAAUCUUGAATsT	A18635	108	UUCAAGAUUGUAGCAGUUUTsT	AD-11619
458-476	GP	A18636	109	AACUGCUACAAUCUUGAAATsT	A18637	110	UUUCAAGAUUGUAGCAGUUTsT	AD-11620
459-477	GP	A18638	111	ACUGCUACAAUCUUGAAATsT	A18639	112	AUUUCAAGAUUGUAGCAGUTsT	AD-11621

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
599-617	VP30	A18640	113	AGCAAUCCAACGGCUGAUTsT	A18641	114	AUCAGCCGUUGGAUUUGCTsT	AD-11622
600-618	VP30	A18642	115	GCAAUCCAACGGCUGAUGTsT	A18643	116	CAUCAGCCGUUGGAUUUGCTsT	AD-11623
601-619	VP30	A18644	117	CAAAUCCAACGGCUGAUGATsT	A18645	118	UCAUCAGCCGUUGGAUUUGTsT	AD-11624
135-153	L	A18646	119	UUGGACCAUGUGACCUAGTsT	A18647	120	CUAGGUCACAUUGGUCCAATsT	AD-11625
136-154	L	A18648	121	UGGACCAUGUGACCUAGUTsT	A18649	122	ACUAGGUCACAUUGGUCCAATsT	AD-11626
2100-2118	L	A18650	123	AUGCAUGUCAGUGAUUAUUTsT	A18651	124	AAUAAUCACUGACAUGCAUTsT	AD-11627
2101-2119	L	A18652	125	UGCAUGUCAGUGAUUAUUTsT	A18653	126	UAAUAAUCACUGACAUGCATsT	AD-11628
2102-2120	L	A18654	127	GCAUGUCAGUGAUUAUUTsT	A18655	128	AUAAUAAUCACUGACAUGCTsT	AD-11629
2103-2121	L	A18656	129	CAUGUCAGUGAUUAUUTsT	A18657	130	UAUAAUAAUCACUGACAUGTsT	AD-11630
2104-2122	L	A18658	131	AUGUCAGUGAUUAUUAATsT	A18659	132	UUAUAAUAAUCACUGACAUTsT	AD-11631
2114-2132	L	A18660	133	UUAUUAUAAUCCACCACAUTsT	A18661	134	AUGUGGUGGAUUAUAAUATsT	AD-11632
2115-2133	L	A18662	135	UAUUAUAAUCCACCACAUATsT	A18663	136	UAUGUGGUGGAUUAUAAUTsT	AD-11633
2116-2134	L	A18664	137	AUUUAUAAUCCACCACAUATsT	A18665	138	UUAUGUGGUGGAUUAUAAUTsT	AD-11634
2412-2430	L	A18666	139	AAAGUUACAAGUGCCUGUGTsT	A18667	140	CACAGGCACUUGUAACUUTsT	AD-11635
2413-2431	L	A18668	141	AAGUUACAAGUGCCUGUGTsT	A18669	142	CCACAGGCACUUGUAACUUTsT	AD-11636
2466-2464	L	A18670	143	UCAGGUUUUAUCUAUUUUGTsT	A18671	144	CAAAUAGAUAAAACUGATsT	AD-11637
2467-2485	L	A18672	145	CAGGUUUUAUCUAUUUUGTsT	A18673	146	CCAAAUAGAUAAAACUGTsT	AD-11638
2556-2574	L	A18674	147	UCUGAUGCAAUUUUUGAUGTsT	A18675	148	CAUCAAAAAUUGCAUCAGATsT	AD-11639
2557-2575	L	A18676	149	CUGAUGCAAUUUUUGAUGATsT	A18677	150	UCAUCAAAAAUUGCAUCAGTsT	AD-11640
1825-1843	NP	A18678	1085	AGuuAcucGGAAAacGGcATsT	A18679	1086	UGCCGUUUUCCGAGuAACUTsT	AD-11641
1588-1606	NP	A18680	1087	AAcGcuAuGGuAAcucuAATsT	A18681	1088	UuAGAGUuACcAuAGCGUUTsT	AD-11642
1827-1845	NP	A18682	1089	uuACUCGGAAAAcGGcAuGTsT	A18683	1090	CAUGCCGUUUUCCGAGuAATsT	AD-11643
1583-1601	NP	A18684	1091	AAAcAAAcGcuAuGGuAAcTsT	A18685	1092	GUuACcAuAGCGUUUGUUTsT	AD-11644

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1488-1506	NP	A18686	1093	AGAGuCuCGcGAAcuuGAcTsT	A18687	1094	GUcAAGUUCGCGAGACUCUTsT	AD-11645
1489-1507	NP	A18688	1095	GAGucucGcGAAcuuGAccTsT	A18689	1096	GGUcAAGUUCGCGAGACUCTsT	AD-11646
1585-1603	NP	A18690	1097	AcAAAcGcuAuGGuAAcucTsT	A18691	1098	GAGUuACcAuAGCGUUUGUTsT	AD-11647
1586-1604	NP	A18692	1099	cAAAcGcuAuGGuAAcucuTsT	A18693	1100	AGAGUuACcAuAGCGUUUGTsT	AD-11648
2231-2249	NP	A18694	1101	cAccGGcucGcGuAuAcAGTsT	A18695	1102	CUGuAuACGGGAGCCGGUGTsT	AD-11649
2873-2891	NP	A18696	1103	CuAACuAGcGAuuuAuCuATsT	A18697	1104	uAGAuAAAUCGCUAGUuAGTsT	AD-11650
1172-1190	VP35	A18698	1105	GcuGAACuAuAGGGuAcGuTsT	A18699	1106	ACGuACCCuAuAGUUCAGCTsT	AD-11651
1176-1194	VP35	A18700	1107	AACuAuAGGGuAcGuuACATsT	A18701	1108	UGuAACGuACCCuAuAGUUTsT	AD-11652
1174-1192	VP35	A18702	175	uGAACuAuAGGGuAcGuuATsT	A18703	176	uAACGuACCCuAuAGUUCATsT	AD-11653
1178-1196	VP35	A18706	177	cuAuAGGGuAcGuuAcAuTsT	A18707	178	AAUGuAACGuACCCuAuAGTsT	AD-11654
251-269	VP35	A18704	179	GGAuAuGcuAcGcAucccTsT	A18705	180	GGGAUGCGuAGcAuAAUCCTsT	AD-11655
416-434	VP35	A18708	181	UUAGAAcAAcGcAuAcGATsT	A18709	182	UCGuAAUGCGUUGUUCuAATsT	AD-11656
421-439	VP35	A18710	183	AcAAcGcAuAcGAGucuuTsT	A18711	184	AAGACUCGuAAUGCGUUGUTsT	AD-11657
1057-1075	VP35	A18712	185	uGAucGAGGuuGGGuAuGuTsT	A18713	186	AcAuACCcAACCCUGAUcATsT	AD-11658
167-185	GP	A18714	187	CCUCGuGAucGAuucAAGATsT	A18715	188	UCUUGAAUCGAUcACGAGGTsT	AD-11659
163-181	GP	A18716	189	GuuAccucGuGAucGAuucTsT	A18717	190	GAAUCGAUcACGAGGuAACTsT	AD-11660
658-676	GP	A18720	191	AAcGAcuuucGcuGAAGGuTsT	A18721	192	ACCUUCAGCGAAAGUCGUUTsT	AD-11662
755-773	GP	A18722	193	AcGGAGGAcccGucuAGuGTsT	A18723	194	cACuAGACGGGUCCUCCGUTsT	AD-11663
966-984	GP	A18724	195	AGGucAAcCCCCAAuuGATsT	A18725	196	UCAAUUUCGGGGUUGACCUTsT	AD-11664
978-996	GP	A18726	197	AAAuAuGauAcAAcAAucGGTsT	A18727	198	CCGAUUGUUGuAuCAAUUUTsT	AD-11665
985-1003	GP	A18728	199	uAcAAcAAucGGGGAGuGGTsT	A18729	200	CcACUCCCCGAUUGUUGuATsT	AD-11666
1101-1119	GP	A18730	201	AGAGuccGGcGcGAAcuucTsT	A18731	202	GAAGuUCGCGCCGGACUCUTsT	AD-11667
1730-1748	GP	A18718	203	uGGAuAccAuAuucGGGcTsT	A18719	204	GCCCCGAAuAUGGuAUCcATsT	AD-11661

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1820-1838	GP	A18732	205	cuGGccAAcGAGAcGAcucTsT	A18733	206	GAGUCGUCUCGUUGGCcAGTsT	AD-11668
1298-1316	VP30	A18734	207	uAucGcucGuAAuAuAAccTsT	A18735	208	GGUuAuAUuACGAGCGAuATsT	AD-11669
295-313	VP30	A18736	209	uucGAGcAcGAucAucAucTsT	A18737	210	GAuGAuGAUCGuGCUCGAATsT	AD-11670
590-608	VP30	A18738	211	CUCGcGcuuAGcAAAuccATsT	A18739	212	UGGAUUUGCuAAGCGCGAGTsT	AD-11671
519-537	VP30	A18740	213	uuAcuccuAcuAAucGcccTsT	A18741	214	GGGCGAUuAGuAGGAGuAATsT	AD-11672
126-144	VP30	A18742	215	cuGcGAAccGGuAGAGuuuTsT	A18743	216	AAACUCuACCGGUUCGcAGTsT	AD-11673
133-151	VP30	A18744	217	ccGGuAGAGuuuAGuuGcATsT	A18745	218	UGcAACuAAACUCuACCGGTsT	AD-11674
292-310	VP30	A18746	219	AuGuucGAGcAcGAucAucTsT	A18747	220	GAUGAUCGUGCUCGAAcAUTsT	AD-11675
321-339	VP30	A18748	221	AAuuAucGAGGuGAGuAccTsT	A18749	222	GGuACUcACCUCGAuAAUUTsT	AD-11676
910-928	VP30	A18750	223	GGGAccGAcAAucCCuAAuTsT	A18751	224	AUuAGGGAUUUGUCGGUCCCTsT	AD-11677
1295-1313	VP30	A18752	225	ucGuAucGcucGuAAuAuATsT	A18753	226	uAuAUuACGAGCGAuACGATsT	AD-11678
331-349	VP30	A18754	227	GuGAGuAccGucAAucAAGTsT	A18755	228	CUUGAUUGACGGuACUcACTsT	AD-11679
123-141	VP30	A18756	229	GAucUGcGAAccGGuAGAGTsT	A18757	230	CUCuACCGGUUCGcAGAUCTsT	AD-11680
124-142	VP30	A18758	231	AucUGcGAACCGGuAGAGuTsT	A18759	232	ACUCuACCGGUUCGcAGAUTsT	AD-11681
1293-1311	VP30	A18760	233	ucucGuAucGcucGuAAuATsT	A18761	234	uAUuACGAGCGAuACGAGATsT	AD-11682
145-163	VP30	A18762	235	AGuuGcAAccuAACACACATsT	A18763	236	UGUGUGUuAGGUUGcAACUTsT	AD-11683
293-311	VP30	A18764	237	uGuucGAGcAcGAucAucATsT	A18765	238	UGAUGAUCGUGCUCGAAcATsT	AD-11684
358-376	VP30	A18766	239	cAcAAGuGcGcGuuccuAcTsT	A18767	240	GuAGGAACGCGcACUUGUGTsT	AD-11685
359-377	VP30	A18768	241	AcAAGuGcGcGuuccuAcuTsT	A18769	242	AGUAGGAACGCGCACUUGUTsT	AD-11686
518-536	VP30	A18770	243	AuuAcuCCuAuuAAucGccTsT	A18771	244	GGCGAUuAGuAGGAGuAAUTsT	AD-11687
520-538	VP30	A18772	245	uAcuccuAuuAAucGCCCGTsT	A18773	246	CGGGCGAUuAGuAGGAGuATsT	AD-11688
524-542	VP30	A18774	247	CCuAuuAAucGcccGuAAGTsT	A18775	248	CUuACGGGCGAUuAGuAGGTsT	AD-11689
525-543	VP30	A18776	249	CuACuAAucGcccGuAAGATsT	A18777	250	UCUuACGGGCGAUuAGuAGTsT	AD-11690

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
584-602	VP30	A18778	251	cAAGGAcucGcGcuuAGcATsT	A18779	252	UGCcAAGCGCGAGUCCUUGTsT	AD-11691
469-487	VP24	A18784	253	AGcuAcGGGAcGAuAcAAuTsT	A18785	254	AUUGuAUCGUCCCGuAGCUTsT	AD-11694
910-928	VP24	A18786	255	GuCGuuGAuuCGAuCCAAuTsT	A18787	256	AUUGGAUCGAAUCAACGACTsT	AD-11695
467-485	VP24	A18788	257	AAAGcuAcGGGAcGAuAcATsT	A18789	258	UGuAUCGUCCCGuAGCUUUTsT	AD-11696
862-880	VP24	A18792	259	CAACAuGcGAACACAAcGuTsT	A18793	260	ACGUUGUGUUCGCAUGUUGTsT	AD-11698
466-484	VP24	A18796	261	uAAAGcuAcGGGAcGAuAcTsT	A18797	262	GuAUCGUCCCGuAGCUUuATsT	AD-11700
523-541	VP24	A18804	263	uGucuuAAGcGAccucuGuTsT	A18805	264	AcAGAGGUCGCUuAAGAcATsT	AD-11704
958-976	VP24	A18806	265	ucuAcAuGucGuGAACuAcTsT	A18807	266	GuAGUUCACGAcAUGuAGATsT	AD-11705
959-977	VP24	A18808	267	cuAcAuGucGuGAACuAcATsT	A18809	268	UGuAGUUCACGAcAUGuAGTsT	AD-11706
971-989	VP24	A18810	269	AAcuAcAAcGGAuuGuuGATsT	A18811	270	UcAAcAAUCCGUUGuAGUUTsT	AD-11707
1071-1089	VP24	A18812	271	ccGAcAAAucGGcAAuGAATsT	A18813	272	UUCAUUGCCGAUUUGUCGGTsT	AD-11708
5886-5904	L	A18816	273	AGAuGAAuuGuAcGAAGTsT	A18817	274	CUUCGuAcAAUUUCGAUCUTsT	AD-11710
192-210	L	A18818	275	AAuccGcAAcuAcGcAAcuTsT	A18819	276	AGUUGCGuAGUUGCGGAUUTsT	AD-11711
5395-5413	L	A18820	277	cAcGccAAuuAAcGucAucTsT	A18821	278	GAUGACGUuAAUUGGCGUGTsT	AD-11712
193-211	L	A18822	279	AuccGcAAcuAcGcAAcuGTsT	A18823	280	cAGUUGCGuAGUUGCGGAUTsT	AD-11713
219-237	L	A18824	281	CCGAAAcAuAuCuAccGuuTsT	A18825	282	AACGGuAGAuAUGUUUCGGTsT	AD-11714
2840-2858	L	A18826	283	uuucuAccGGAAucuAGGATsT	A18827	284	UCCuAGAUAUCCGGuAGAAATsT	AD-11715
4779-4797	L	A18828	285	AuuAAucGcGGAAcAAuuGTsT	A18829	286	cAAUUGUUCGCGAUuAAUTsT	AD-11716
5275-5293	L	A18830	287	AuuucGAucGAucGAGAcATsT	A18831	288	uGUCUCGAUCGAUCGAAAUTsT	AD-11717
5391-5409	L	A18832	289	GGGAcAcGccAAuuAAcGuTsT	A18833	290	ACGUuAAUUGGCGUGUCCCTsT	AD-11718
191-209	L	A18834	291	uAAuccGcAAcuAcGcAAcTsT	A18835	292	GUUGCGuAGUUGCGGAUuATsT	AD-11719
1614-1632	L	A18836	293	AGuAcuAAAcGuGuAccGGTsT	A18837	294	CCGGuAcACGUUuAGuACUTsT	AD-11720
4588-4606	L	A18838	295	cAcAucGcucAuuGcGGAuTsT	A18839	296	AUUCGcAAUGAGCGAUGUGTsT	AD-11721

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
4590-4608	L	A18840	297	cAucGcucAuuGcGAuAcTsT	A18841	298	GuAUUCGcAAUGAGCGAUGTsT	AD-11722
5884-5902	L	A18842	299	AcAGAUcGAAuuGuAcGATsT	A18843	300	UCGuAcAAUUUCGAUCUGUTsT	AD-11723
161-179	L	A18844	301	AGcuuGcGGGuuAuAuucATsT	A18845	302	UGAAuAuAACCCGcAAGCUTsT	AD-11724
778-796	L	A18846	303	cuGccGAcGucuuGAuAAuTsT	A18847	304	AUuAUcAAGACGUCGGcAGTsT	AD-11725
5446-5464	L	A18848	305	AGuAcuuAcGGcAAuuGAGTsT	A18849	306	CUCAAUUGCCGuAAGuACUTsT	AD-11726
6297-6315	L	A18850	307	AAccucGucGAuucAAAAATsT	A18851	308	UUUUUGAAUCGACGAGGuUTsT	AD-11727
5269-5287	L	A18852	309	AAcuAAuuucGAucGAucTsT	A18853	310	GAUCGAUCGAAUUuAGUUTsT	AD-11728
1778-1796	L	A18854	311	GccuuAuccGAcucGcAAuTsT	A18855	312	AUUGCGAGUCGGAuAAGCTsT	AD-11729
1780-1798	L	A18856	313	CuuAuccGAcucGcAAuGuTsT	A18857	314	AcAUUGCGAGUCGGAuAAGTsT	AD-11730
3163-3181	L	A18858	315	GucGuuuuGcGGccGAuAuTsT	A18859	316	AuAUCGGCCGcAAAACGACTsT	AD-11731
3164-3182	L	A18860	317	ucGuuuuGcGGccGAuAuCTsT	A18861	318	GAuAUCGGCCGcAAAACGATsT	AD-11732
5273-5291	L	A18862	319	AAAuuuGcAucGAucGAGATsT	A18863	320	UCUCGAUCGAUCGAAuuUTsT	AD-11733
6295-6313	L	A18864	321	AuAAcCucGucGAuucAAATsT	A18865	322	UUUGAAUCGACGAGGUuAUTsT	AD-11734
1702-1720	L	A18866	323	uAcuAccAcAAuAucGGAATsT	A18867	324	UUCCGAuAUUGUGGuAGuATsT	AD-11735
1781-1799	L	A18868	325	uuAuccGAcucGcAAuGuuTsT	A18869	326	AAcAUUGCGAGUCGGAuAATsT	AD-11736
5270-5288	L	A18870	327	AcuAAAuuuGcAucGAucGTsT	A18871	328	CGAUCGAUCGAAUUuAGUTsT	AD-11737
5276-5294	L	A18872	329	uuucGAucGAucGAGAcAcTsT	A18873	330	GuGUCUCGAUCGAUCGAAATsT	AD-11738
5394-5412	L	A18874	331	AcAcGccAAuuAAcGucAuTsT	A18875	332	AUGACGUuAAUUGGCGUGUTsT	AD-11739
6242-6260	L	A18876	333	AAGuuAuAuccGccuuGGuTsT	A18877	334	ACcAAGGCGGAuAuAACUUTsT	AD-11740
182-200	L	A18878	335	AuAcuCCCUuAAuccGcAATsT	A18879	336	UUGCGGAUuAAGGGAGuAUTsT	AD-11741
194-212	L	A18880	337	uCCGcAAcuAcGcAAcuGuTsT	A18881	338	AcAGUUGCGuAGUUGCGGATsT	AD-11742
575-593	L	A18882	339	ucGAGGAAAcuCuAGAuCATsT	A18883	340	UGAUCuAGAGUUUCCUCGATsT	AD-11743
1565-1583	L	A18884	341	uGcAGuAuucGAGCCuAAuTsT	A18885	342	AUuAGGCUCGAAuACUGcATsT	AD-11744

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1566-1584	L	A18886	343	GcAGuAuucGAGccuAAuGTsT	A18887	344	cAUuAGGCUCGAAuACUGCTsT	AD-11745
1567-1585	L	A18888	345	CAGuAuucGAGcCuAAuGTsT	A18889	346	ACAuUAGGCUCGAAuACUGTsT	AD-11746
2779-2797	L	A18890	347	cAuuGGcAcuAGcGGuAccTsT	A18891	348	GGuACCGCuAGUGCcAAUGTsT	AD-11747
2838-2856	L	A18892	349	uGuuucuAccGGAuAuAGTsT	A18893	350	CuAGAUAUCCGGuAGAAAcATsT	AD-11748
2892-2910	L	A18894	351	AcuuAucuccGAAuGAuuGTsT	A18895	352	cAAUcAUUCGAGAuAAGUTsT	AD-11749
2981-2999	L	A18896	353	AAAuccuAGcGGAuuAAATsT	A18897	354	AUUuAAUCCGCuAGGAUUUTsT	AD-11750
2982-3000	L	A18898	355	AAuccuAGcGGAuuAAuGTsT	A18899	356	CAUUUAAUCCGCuAGGAUUUTsT	AD-11751
3038-3056	L	A18900	357	GAuuGuAcGcAGGAccAucTsT	A18901	358	GAUGGUCCUGCGuAcAAUCTsT	AD-11752
3149-3167	L	A18902	359	AAcuccuGuuAuGAGucGuTsT	A18903	360	ACGACUcAuAAcAGGAGUUTsT	AD-11753
3168-3186	L	A18904	361	uuuGcGGccGAuAucuuuuTsT	A18905	362	AAAAGAuAUCGGCCGcAAATsT	AD-11754
3889-3907	L	A18906	363	GGuAcAAcGAucAAuAcAGTsT	A18907	364	CUGuAUUGAUCGUUGuACCTsT	AD-11755
3922-3940	L	A18908	365	uGGccAAucGuAuGAGuAATsT	A18909	366	UuACUcAuACGAUUGGCcATsT	AD-11756
4001-4019	L	A18910	367	GucuGcAcGcGAcAGcAAuTsT	A18911	368	AUUGCUGUCGCGUGcAGACTsT	AD-11757
4584-4602	L	A18912	369	cuAccAcAucGcucAuuGcTsT	A18913	370	GcAAUGAGCGAUGUGGuAGTsT	AD-11758
4593-4611	L	A18914	371	cGcucAuuGcGAAuAcuuATsT	A18915	372	uAAGuAUUCGcAAUGAGCGTsT	AD-11759
4598-4616	L	A18916	373	AuuGcGAAuAcuuAAGccATsT	A18917	374	UGGCUuAAGuAUUCGcAAUTsT	AD-11760
4601-4619	L	A18918	375	GcGAAuAcuuAAGccAAcATsT	A18919	376	UGUUGGCUuAAGuAUUCGCTsT	AD-11761
4638-4656	L	A18920	377	AuGucAcGGuuAAuGAGuATsT	A18921	378	UACUCAUUAACCGUGACAUTsT	AD-11762
4778-4796	L	A18922	379	AAuuAAucGcGGAACAAuTsT	A18923	380	AAUUGUUCGCGAUuAAUUTsT	AD-11763
5274-5292	L	A18924	381	AAuuucGAucGAucGAGAcTsT	A18925	382	GUCUCGAUCGAUCGAAuUTsT	AD-11764
5392-5410	L	A18926	383	GGAcAcGccAAuuAAcGucTsT	A18927	384	GACGUuAAUUGGCGUGUCTsT	AD-11765
5649-5667	L	A18928	385	AcGcuAGcuAcuGAGuccATsT	A18929	386	UGGACUcAGuAGCuAGCGUTsT	AD-11766
5833-5851	L	A18930	387	cuAAGcAAGucGAGGuuAuTsT	A18931	388	AuAACCUCGACUUGCUuAGTsT	AD-11767

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
6243-6261	L	A18932	389	AGuuAuAuccGccuuGGuuTsT	A18933	390	AACcAAGGCGGAuAuAACUTsT	AD-11768
6290-6308	L	A18934	391	cAGGuAuAAccucGucGAuTsT	A18935	392	AUCGACGAGGUuAuACCUGTsT	AD-11769
6291-6309	L	A18936	393	AGGuAuAAccucGucGAuTsT	A18937	394	AAUCGACGAGGUuAuACCUTsT	AD-11770
1816-1834	NP	A18938	395	AcuAcGAGGAuucGGcuGATsT	A18939	396	UcAGCCGAAUCCUCGuAGUTsT	AD-11771
875-893	NP	A18940	397	uCuAcccAAAcuuGucGuuTsT	A18941	398	AACGAcAAGUUUGGGuAGATsT	AD-11772
1817-1835	NP	A18942	399	CuAcGAGGAuucGGcuGAATsT	A18943	400	UUcAGCCGAAUCCUCGuAGTsT	AD-11773
1812-1830	NP	A18944	401	CCuGAcuAcGAGGAuucGGTsT	A18945	402	CCGAAUCCUCGuAGUcAGGTsT	AD-11774
1819-1837	NP	A18946	403	AcGAGGAuucGGCuGAAGTsT	A18947	404	CCUUCAGCCGAAUCCUCGUTsT	AD-11775
2140-2158	NP	A18948	405	AcGAGAGucucAcAucccuTsT	A18949	406	AGGGAuGuGAGACUCUCGUTsT	AD-11776
730-748	VP35	A18950	407	AAAuuuCGGGcGACCuuACTsT	A18951	408	GuAAGGUCGCCCGAAAuUTsT	AD-11777
735-753	VP35	A18952	409	uCGGGcGAccuuAcAuucTsT	A18953	410	GAAAUGuAAGGUCGCCCGATsT	AD-11778
195-213	VP35	A18954	411	uGAccGGcAAAAuAccGcuTsT	A18955	412	AGCGGuAUUUUGCCGGuATsT	AD-11779
198-216	VP35	A18956	413	CCGGcAAAAuACCGcuAACTsT	A18957	414	GUuAGCGGuAUUUUGCCGGTsT	AD-11780
379-397	VP35	A18958	415	AGcuGuGcGucGGcAAAccTsT	A18959	416	GGUUUGCCGACGcAcAGCUTsT	AD-11781
646-664	VP35	A18960	417	AuuGAAAGAuucGAAcGGTsT	A18961	418	CCCGUUCGGAUCUUUcAAUTsT	AD-11782
731-749	VP35	A18962	419	AAuuucGGGcGAccuuAcATsT	A18963	420	UGuAAGGUCGCCCGAAAUTsT	AD-11783
732-750	VP35	A18964	421	AuuucGGGcGAccuuAcAuTsT	A18965	422	AUGuAAGGUCGCCCGAAAUTsT	AD-11784
1193-1211	VP35	A18966	423	GucuAuGuGucAuAAGcuTsT	A18967	424	AGCUuAUGAcAcAAuAGACTsT	AD-11785
438-456	VP40	A18968	425	cucGcAucuuAuAcGAucATsT	A18969	426	UGAUCGuAuAAGAUCGAGTsT	AD-11786
1301-1319	VP40	A18970	427	uGcAuAAGcGAuCCAuAcuTsT	A18971	428	AGuAUGGAUCGCUuAUGcATsT	AD-11787
1191-1209	VP40	A18972	429	AAuGuAcuAAucGGGuCAATsT	A18973	430	UUGACCCGAUuAGuAcAUUTsT	AD-11788
442-460	VP40	A18974	431	CAuCUuAuAcGAuCACCCATsT	A18975	432	UGGGUGAUCGuAuAAGAUGTsT	AD-11789
443-461	VP40	A18976	433	AUCUUuAuAcGAucAcccAuTsT	A18977	434	AUGGGUGAUCGuAuAAGAUTsT	AD-11790

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
478-496	VP40	A18978	435	AcccccucGuuAGAGuGAATsT	A18979	436	UUCACUCuAACGAGGGGUTsT	AD-11791
834-852	VP40	A18980	437	AucGuGccAAuuGAuccAGTsT	A18981	438	CUGGAUcAAUUGGcACGAUTsT	AD-11792
1192-1210	VP40	A18982	439	AuGuAcuAAucGGGucAAGTsT	A18983	440	CUUGACCCGAUuAGuAcAUTsT	AD-11793
1194-1212	VP40	A18984	441	GuAcuAAucGGGucAAGATsT	A18985	442	UCCUUGACCCGAUuAGuACTsT	AD-11794
1300-1318	VP40	A18986	443	AuGcAuAAGcGAuccAuAcTsT	A18987	444	GuAUGGAUCGCUuAUGcAUTsT	AD-11795
465-483	GP	A18988	445	AcGGGAGcGAAuGcuuAccTsT	A18989	446	GGuAAGcAUUCGCUCCCGUTsT	AD-11796
358-376	VP30	A18990	447	AGuuAGAGucccuAcGGuuTsT	A18991	448	AACCGuAGGGACUCuAACUTsT	AD-11797
331-349	VP30	A18992	449	cuAccGuAGuAGucGAAGuTsT	A18993	450	ACUUCGACuACuACGGuAGTsT	AD-11798
250-268	VP30	A18994	451	GAAuuCAcGuGccGAccAGTsT	A18995	452	CUGGUCGGcACGUGAAUUCTsT	AD-11799
1009-1027	VP30	A18996	453	uGcccccccAAGcGuuAAuTsT	A18997	454	AUuAACGCUUGGGGGGcATsT	AD-11800
1318-1336	VP30	A18998	455	AGAGuGuuAGGAucGuuAuTsT	A18999	456	AuAACGAUCCuAAcACUCUTsT	AD-11801
126-144	VP30	A19000	457	AAucccGAGGcGGcAAuucTsT	A19001	458	GAAuuGCCGCCUCGGGAuUTsT	AD-11802
354-372	VP30	A19002	459	cGcAAGuuAGAGucccuAcTsT	A19003	460	GuAGGGACUCuAACUUGCGTsT	AD-11803
553-571	VP30	A19004	461	uGAuucAucGcuuAAuAuTsT	A19005	462	uAuAUuAAGCGAUGAAUcATsT	AD-11804
583-601	VP30	A19006	463	AGAccuAAGAcuAGcAAuTsT	A19007	464	AUUUGCuAGUCUuAGGUCUTsT	AD-11805
652-670	VP30	A19008	465	AuuAcuAGucGAGAcuGcuTsT	A19009	466	AGcAGUCUCGACuAGuAAUTsT	AD-11806
992-1010	VP30	A19010	467	ucAGGccuAcGcuuAcuuGTsT	A19011	468	CAAGuAAGCGuAGGCCUGATsT	AD-11807
1013-1031	VP30	A19012	469	CCCCAAGcGuuAAuGAAGTsT	A19013	470	CUUCAUuAACGCUUGGGGGTsT	AD-11808
404-422	VP24	A19014	471	AuuAuAcGGGuccAuAAuTsT	A19015	472	AUuAAUGGACCCGuAuAAUTsT	AD-11809
888-906	VP24	A19016	473	cucAAcGAGuAAAGGAccATsT	A19017	474	UGGUCCUuAcUCGUUGAGTsT	AD-11810
1247-1265	VP24	A19018	475	uuGuAcGAuAGGGcuAAcATsT	A19019	476	UGUuAGCCCuAUCGuAcAATsT	AD-11811
536-554	VP24	A19020	477	GuuGuGuuuAGcGAccuAuTsT	A19021	478	AuAGGUCGCUAAAcAcAACTsT	AD-11812
1050-1068	VP24	A19022	479	GGAcuAAuAuGGGuuAucuTsT	A19023	480	AGAuAACCcAuAUuAGUCCTsT	AD-11813

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1095-1113	VP24	A19024	481	CuGcGAuGGAuAuAcGAcATsT	A19025	482	UGUCGuAuAUCcAUCGcAGTsT	AD-11814
535-553	VP24	A19026	483	AGuuGuGuuuAGcGAccuATsT	A19027	484	uAGGUCGCUAAAcAcAACUTsT	AD-11815
196-214	VP24	A19028	485	uuGAAcuAGucuAcucGcATsT	A19029	486	UGCGAGuAGAcuAGUUCaATsT	AD-11816
215-233	VP24	A19030	487	GAAuCCuAccGGGAAuAGATsT	A19031	488	UCuAUUCCCGGuAGGAUUCTsT	AD-11817
403-421	VP24	A19032	489	uAuuAuAcGGGuCCAuAATsT	A19033	490	UuAAUGGACCCGuAuAAuATsT	AD-11818
406-424	VP24	A19034	491	uAuAcGGGuccAuuAAuuTsT	A19035	492	AAAUuAAUGGACCCGuAuATsT	AD-11819
1140-1158	VP24	A19036	493	uACAuGAAucGACACuuAATsT	A19037	494	UuAAGUGUCGAUUCAGuATsT	AD-11820
1243-1261	VP24	A19038	495	AAAAuuGuAcGAuAGGGcuTsT	A19039	496	AGCCCuAUCGuAcAAUUUUTsT	AD-11821
1249-1267	VP24	A19040	497	GuAcGAuAGGGcuAAcAuuTsT	A19041	498	AAUGUuAGCCCuAUCGUACTsT	AD-11822
1590-1608	VP24	A19042	499	GAGcccAAAuuAACAcGGuTsT	A19043	500	ACCGUGUuAAUUUGGGCUCTsT	AD-11823
3688-3706	L	A19044	501	cccGcuAuuAAGccGAGGuTsT	A19045	502	ACCUCGGCUuAAuAGCGGGTsT	AD-11824
3687-3705	L	A19046	503	GcccGcuAuuAAGccGAGGTsT	A19047	504	CCUCGGCUuAAuAGCGGGTsT	AD-11825
2956-2974	L	A19048	505	AAuuGuAGcGcAAuuGAcuTsT	A19049	506	AGUcAAUUGCGCUAcAAUUTsT	AD-11826
2615-2633	L	A19050	507	AGcGAucAAucuccGAAAcTsT	A19051	508	GuuUCGGAGAuGAUUCGCUTsT	AD-11827
2612-2630	L	A19052	509	uuGAGcGAucAAucuccGATsT	A19053	510	UCGGAGAUUGAUUCGCUcAATsT	AD-11828
4595-4613	L	A19054	511	uucGAAucuuCAAacGAcTsT	A19055	512	GUCGGuuuGAAGAuUCGAATsT	AD-11829
2613-2631	L	A19056	513	uGAGcGAuCAAuCuCCGAATsT	A19057	514	UUCGGAGAUUGAUCGCUCATsT	AD-11833
2614-2632	L	A19058	515	GAGcGAucAAucuccGAAATsT	A19059	516	uuUCGGAGAuGAUUCGCUCTsT	AD-11831
3941-3959	L	A19060	517	CAAcGcGcuuGAuGGuAuCTsT	A19061	518	GAuACcAUcAAGCGCGUUGTsT	AD-11832
3942-3960	L	A19062	519	AAcGcGcuuGAuGGuAucTsT	A19063	520	AGAuACcAUcAAGCGCGUUTsT	AD-11833
1680-1698	L	A19064	521	AuAcGcCCAAGAAcuuAGGTsT	A19065	522	CCuAAGUUCUUGGGCGuAUTsT	AD-11834
3686-3704	L	A19066	523	AGcccGcuAuuAAGccGAGTsT	A19067	524	CUCGGCUuAAuAGCGGGCUTsT	AD-11835
4255-4273	L	A19068	525	UUAucGAuuGAcAGuccuuTsT	A19069	526	AAGGACUGUcAAUCGAuAATsT	AD-11836

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1374-1392	L	A19070	527	AGAccGAuGuuuAAcGCcGTsT	A19071	528	CGGCGUuAAAcAUCGGUCUTsT	AD-11837
5470-5488	L	A19072	529	AccAuAuAuuGucGcuucATsT	A19073	530	UGAAGCGAcAAuAuAUGGUTsT	AD-11838
3872-3890	L	A19074	531	AuAuuGuGcAucGGuAuAATsT	A19075	532	UuAuACCGAUGcAcAAuAUTsT	AD-11839
1384-1402	L	A19076	533	uuAAcGccGGGAuuGAAuTsT	A19077	534	AAUUCAAUCCCGGCGUuAATsT	AD-11840
4519-4537	L	A19078	535	uGcAcGAAAAAGAcGGAcTsT	A19079	536	GUCCGAUCUUUUUCUGGcATsT	AD-11841
3682-3700	L	A19080	537	GGucAGcccGcuAuuAAGcTsT	A19081	538	GCUuAAuAGCGGGCUGACCTsT	AD-11842
2954-2972	L	A19082	539	GGAAuuGuAGcGcAAuuGATsT	A19083	540	UCAAUUGCGCUACAAUUCCTsT	AD-11843
5467-5485	L	A19084	541	AcuAccAuAuAuuGucGcuTsT	A19085	542	AGCGAcAAuAuAUGGuAGUTsT	AD-11844
1376-1394	L	A19086	543	AccGAuGuuuAAcGccGGGTsT	A19087	544	CCCGGCGUuAAAcAUCGGUTsT	AD-11845
2448-2466	L	A19088	545	uGAuGAGAcuuucGuAcAcTsT	A19089	546	GUGuACGAAAGUCUcAUcATsT	AD-11846
1023-1041	L	A19090	547	AcGAAAAGGGcGGuuuuuATsT	A19091	548	uAAAAACCGCCUUUUCGUTsT	AD-11847
1377-1395	L	A19092	549	CCGAuGuuuAAcGccGGGATsT	A19093	550	UCCCGGCGUuAAAcAUCGGTsT	AD-11848
2619-2637	L	A19094	551	AucAAucuccGAAAcuAGATsT	A19095	552	UCuAGUUUCGGAGAUGAUTsT	AD-11849
5608-5626	L	A19096	553	AAAuAcGGcGuuAAGAAGTsT	A19097	554	ACUUCUuAACGCCGuAUUUTsT	AD-11850
5607-5625	L	A19098	555	AAAAuAcGGcGuuAAGAAGTsT	A19099	556	CUUCUuAACGCCGuAUUUTsT	AD-11851
6396-6414	L	A19100	557	ucGAAcccAGAcuuAucAuTsT	A19101	558	AUGAuAAGUCUGGGUUCGATsT	AD-11852
4165-4183	L	A19102	559	AcAAcCacGcuAAAcuAGTsT	A19103	560	CuAGAUUuAGCGUGGUUGUTsT	AD-11853
4250-4268	L	A19104	561	GcAAcuuAucGAuuGAcAGTsT	A19105	562	CUGUcAAUCGAuAAGUUGCTsT	AD-11854
6434-6452	L	A19106	563	GAcGGAuAAcuAAAcuAGuTsT	A19107	564	ACuAGUUuAGUuAUCCGUCTsT	AD-11855
2959-2977	L	A19108	565	uGuAGcGcAAuuGAcuuuGTsT	A19109	566	cAAAGUcAAUUGCGCuAcATsT	AD-11856
6433-6451	L	A19110	567	GGAcGGAuAAcuAAAcuAGTsT	A19111	568	CuAGUUuAGUuAUCCGUCTsT	AD-11857
83-101	L	A19112	569	uGGcuAcccAAcAuAcAcATsT	A19113	570	UGUGuAUGUUGGGuAGCcATsT	AD-11858
1382-1400	L	A19114	571	GuuuAAcGccGGGAuuGAATsT	A19115	572	UUcAAUCCCGGCGUuAAACTsT	AD-11859

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1014-1032	NP	A19116	573	uuuCCGuuuGAuGcGAACATsT	A19117	574	UGUUCGcAUcAAACGGAAATsT	AD-11860
1805-1823	NP	A19118	575	GAAGCuAcGGcGAAuACCATsT	A19119	576	UGGuAUUCGCCGuAGCUUCTsT	AD-11861
1862-1880	NP	A19120	577	uGGuccuAuucGAucAGATsT	A19121	578	UCuAGAUCGAAuAGGACcATsT	AD-11862
1016-1034	NP	A19122	579	UCCGuuuGAuGcGAACAAATsT	A19123	580	UUUGUUCGCAUCAACGGATsT	AD-11863
2230-2248	NP	A19124	581	CCAccGGcucccGuAuAcATsT	A19125	582	UGuAuACGGGAGCCGGUGGTsT	AD-11864
2233-2251	NP	A19126	583	CCGGcucccGuAuAcAGAGTsT	A19127	584	CUCUGuAuACGGGAGCCGGTsT	AD-11865
959-977	NP	A19136	585	AAGGACuGAuACAuAuCCTsT	A19137	586	GGAuAUUGuAUCAGUCCUUTsT	AD-11870
1017-1035	NP	A19138	587	ccGuuuGAuGcGAACAAATsT	A19139	588	AUUUGUUCGCAUCAACGGTsT	AD-11871
2124-2142	NP	A19140	589	CCCACuGGAcGAuGccGAcTsT	A19141	590	GUCGGcAUCGUCcAGUGGGTsT	AD-11872
745-763	NP	A19142	591	cGuGAuGGAGuGAAGcGccTsT	A19143	592	GGCGCUUCACUCCAUCACGTsT	AD-11873
2229-2247	NP	A19144	593	CCCAccGGcucccGuAuAcTsT	A19145	594	GuAuACGGGAGCCGGUGGGTsT	AD-11874
2119-2137	NP	A20118	595	GcAGAcCCAcuGGAcGAuGTsT	A20119	596	CAUCGUCCAGUGGGUCUGCTsT	AD-12462
1587-1605	NP	A20120	597	AAAcGcuAuGGuAAcucuATsT	A20121	598	uAGAGUuACcAuAGCGUUUTsT	AD-12463
1300-1318	NP	A20122	599	UUCGCCCGACuuuuGAACCTsT	A20123	600	GGUUCAAAAGUCGGGCGAATsT	AD-12464
1808-1826	NP	A20124	601	GcuAcGGcGAAuAccAGAGTsT	A20125	602	CUCUGGuAUUCGCCGuAGCTsT	AD-12465
1813-1831	NP	A20126	603	GGcGAAuACCAGAuAcuTsT	A20127	604	AGuAACUCUGGuAUUCGCCTsT	AD-12466
532-550	VP35	A19146	605	GGuGAuGAcAAccGGucGGTsT	A19147	606	CCGACCGGUUGUcAUcACCTsT	AD-11875
417-435	VP35	A19148	607	uAGAAcAAcGcAuAcGAGTsT	A19149	608	CUCGuAAUGCGUUGUUCuATsT	AD-11876
741-759	VP35	A19152	609	GGAAAcCUGAcAuucGGcTsT	A19153	610	GCCGAAUUGUcAGGUUUCCTsT	AD-11878
1049-1067	VP35	A19154	611	cccAAGAuGaucGAGGuTsT	A19155	612	AACCUCGAUCAUCUUGGGTsT	AD-11879
206-224	VP35	A19160	613	AGAAuuccuGuAAGcGAcATsT	A19161	614	UGUCGCUuAcAGGAAUUCUTsT	AD-11882
246-264	VP35	A19162	615	AuccAGGAuuAuGcuAcGcTsT	A19163	616	GCGuAGcAuAAUCCUGGAUTsT	AD-11883
247-265	VP35	A19164	617	UGCAGGAuuAuGcuAcGcATsT	A19165	618	UGCGuAGcAuAAUCCUGGATsT	AD-11884

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
287-305	VP35	A19166	619	ccAAAcccGAAGAcGcGcATsT	A19167	620	uGCGCGUCuUCGGGuuuGGTsT	AD-11885
314-332	VP35	A19168	621	AcccAAAcGGAcccAAuuuTsT	A19169	622	AAAuuggGUCCGuuuGGGUTsT	AD-11886
319-337	VP35	A19170	623	AAcGGAcCcAAuuuGcAAuTsT	A19171	624	AUUGcAAAUUGGGUCCGUUTsT	AD-11887
414-432	VP35	A19172	625	cAuuAGAAcAAcGCAuuAcTsT	A19173	626	GuAAUGCGUUGUUCuAAUGTsT	AD-11888
415-433	VP35	A19174	627	AuuAGAAcAAcGcAuuAcGTsT	A19175	628	CGuAAUGCGUUGUUCuAAUTsT	AD-11889
439-457	VP35	A19176	629	uGAGAAuGGucAAAGCcATsT	A19177	630	UGGCUUuAGAcAUUCUcATsT	AD-11890
576-594	VP35	A19178	631	AGGcuuAuuGGGccGAACATsT	A19179	632	UGUUCGGCCcAAuAAGCCUTsT	AD-11891
413-431	VP35	A20128	633	ucAuuAGAAcAAcGcAuuATsT	A20129	634	uAAUGCGUUGUUCuAAUGATsT	AD-12467
583-601	VP35	A20130	635	uuGGGccGAACuAGGuCAATsT	A20131	636	UUGACcAUGUUCGGCCcAATsT	AD-12468
983-1001	VP35	A20132	637	cAcAuccGcucucGAGGuGTsT	A20133	638	CACCUCGAGAGCGGAUGUGTsT	AD-12469
318-336	VP35	A20134	639	AAAcGGAcccAAuuuGcAATsT	A20135	640	UUGcAAAUUGGGUCCGUUUTsT	AD-12470
420-438	VP35	A20136	641	AAcAAcGcAuuAcGAGucUTsT	A20137	642	AGACUCGuAAUGCGUUGUUTsT	AD-12471
419-437	VP35	A20138	643	GAAcAAcGcAuuAcGAGucTsT	A20139	644	GACUCGuAAUGCGUUGUUTsT	AD-12472
134-152	VP35	A20140	645	GccAcGAcucAAAACGAcATsT	A20141	646	UGUCGUUUUGAGUCGUGGCTsT	AD-12473
893-911	VP40	A19180	647	ccAcAAGcuGAccGGuAAGTsT	A19181	648	CUuACCGGUcAGCUUGUGTsT	AD-11892
892-910	VP40	A19182	649	uCCAcAAGCuGAccGGuAATsT	A19183	650	UuACCGGUcAGCUUGUGATsT	AD-11893
325-343	VP40	A19188	651	uGAAuGucAuAucGGGcccTsT	A19189	652	GGGCCCGAuAUGAcAUUcATsT	AD-11896
450-468	VP40	A19190	653	AcuAucAcccAuuucGGcATsT	A19191	654	UGCCGAAAUGGGUGAuAGUTsT	AD-11897
662-680	VP40	A19194	655	GAccGAuGAcAcuccAAcATsT	A19195	656	UGUUGGAGUGUcAUCGGUCTsT	AD-11899
200-218	VP40	A19198	657	GAcAccGGAGucAGucAAuTsT	A19199	658	AuuGACuGACUCCGGUGUCTsT	AD-11901
203-221	VP40	A19200	659	AccGGAGucAGucAAuGGGTsT	A19201	660	CCcAUUGACUGACUCCGGUTsT	AD-11902
204-222	VP40	A19202	661	CCGGAGuCAGuCAAuGGGGTsT	A19203	662	CCCCAUUGACUGACUCCGGTsT	AD-11903
225-243	VP40	A19204	663	AcuccAucGAuccAcucATsT	A19205	664	uGAGuGGAuUCGAuGGAGUTsT	AD-11904

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
386-404	VP40	A19206	665	uGucGCuGAucAAAAGACCTsT	A19207	666	GGUCUUUGAUCAGCGACATsT	AD-11905
584-602	VP40	A19208	667	AGuccAAcAcccccAGuAuTsT	A19209	668	AuACUGGGGuAGUUGGACUTsT	AD-11906
631-649	VP40	A19210	669	uGAucAcccAAccAcuGccTsT	A19211	670	GGcAGUGGUUGGGUGAUcATsT	AD-11907
660-678	VP40	A19212	671	uGGACcGAuGACACUCCAATsT	A19213	672	UUGGAGUGUCAUCGGUCCATsT	AD-11908
663-681	VP40	A19214	673	AccGAuGAcAcuCCAACAGTsT	A19215	674	CUGUUGGAGUGUCAUCGGUTsT	AD-11909
929-947	VP40	A19218	675	uGGAcAAccAAucAuCCCuTsT	A19219	676	AGGGAUGAUUGGUUGUCCATsT	AD-11911
1019-1037	VP40	A19220	677	uuGuGAcAcGuGucAuucTsT	A19221	678	AGAAUGAcACGUGUcAcAATsT	AD-11912
243-261	VP40	A19224	679	AGGCcAAuuGccGAuGAcATsT	A19225	680	UGUCAUCGGcAAUUGGCCUTsT	AD-11914
140-158	VP40	A20142	681	AuAcccuGucAGGucAAAuTsT	A20143	682	AUUUGACCUGAcAGGGuAUTsT	AD-12474
141-159	VP40	A20144	683	uAcccuGucAGGucAAAuTsT	A20145	684	AAUUGACCUGAcAGGGuATsT	AD-12475
378-396	VP40	A20146	685	CCUCUAGGuGuCGCuGAuCTsT	A20147	686	GAUCAGCGACACCuAGAGGTsT	AD-12476
427-445	VP40	A20148	687	CCGccAucAuGcuuGcuucTsT	A20149	688	GAAGcAAGcAUGAUGGCGGTsT	AD-12477
898-916	VP40	A20150	689	AGCuGACcGGuAAGAAGGuTsT	A20151	690	ACCUUCUuACCGGUCAGCUTsT	AD-12478
199-217	VP40	A20152	691	uGAcAccGGAGucAGucAATsT	A20153	692	UUGACUGACUCCGGUGUCATsT	AD-12479
568-586	VP40	A20154	693	AGuucGuucuuuccGccAGuTsT	A20155	694	ACUGGCGGAAGAACGAACUTsT	AD-12480
569-587	VP40	A20156	695	GuucGuucuuuccGccAGucTsT	A20157	696	GACUGGCGGAAGAACGAACTsT	AD-12481
1728-1746	GP	A19232	697	ccuGGAuAccAuAuuuuGGTsT	A19233	698	CCGAAuAUGGuAUCcAGGTsT	AD-11918
1729-1747	GP	A19234	699	cuGGAuAccAuAuuuuGGTsT	A19235	700	CCCGAAuAUGGuAUCcAGTsT	AD-11919
1818-1836	GP	A19246	701	AGcuGGCCAAcGAGAcGAcTsT	A19247	702	GUCGUCUCGUUGGCcAGCUTsT	AD-11925
1821-1839	GP	A19248	703	uGGccAAcGAGAcGAcucATsT	A19249	704	UGAGUCGUCUCGUUGGCcATsT	AD-11926
1732-1750	GP	A19250	705	GAuAccAuAuuuuGGGccATsT	A19251	706	UGGCCCGAAuAUGGuAUCTsT	AD-11927
1956-1974	GP	A20158	707	cGGAcuGcuGuAucGAAccTsT	A20159	708	GGUUCGAuAcAGcAGUCCGTsT	AD-12482
2107-2125	GP	A20160	709	uGGAGuuAcAGGcGuuAuATsT	A20161	710	uAuAACGCCUGuAACUCcATsT	AD-12483

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
2124-2142	GP	A20162	711	uAAuGcAGuuAucGCUUUTsT	A20163	712	AAAGCGAuAACUGcAAUuATsT	AD-12484
2109-2127	GP	A20164	713	GAGuuAcAGGcGuuAuAAuTsT	A20165	714	AUuAuAACGCCUGuAACUCTsT	AD-12485
1958-1976	GP	A20166	715	GAcuGcuGuAucGAAccAcTsT	A20167	716	GUGGUUCGAuAcAGcAGUCTsT	AD-12486
1890-1908	GP	A20168	717	uccucAAccGuAAGGcAAuTsT	A20169	718	AUUGCCUuACGGUUGAGGATsT	AD-11487
1891-1909	GP	A20170	719	CCUCAAccGuAAGGcAAuTsT	A20171	720	AAUUGCCUuACGGUUGAGGATsT	AD-12488
1307-1325	GP	A20172	721	AAuAcAcCcGuGuAuAAAcTsT	A20173	722	GUUuAuAcACGGGUGuAUUTsT	AD-12489
1823-1841	GP	A20174	723	GccAAcGAGAcGAcucAAGTsT	A20175	724	CUUGAGUCGUCUCGUUGGCTsT	AD-12490
2110-2128	GP	A20176	725	AGuuAcAGGcGuuAuAAuTsT	A20177	726	AAUuAuAACGCCUGuAACUTsT	AD-12491
1308-1326	GP	A20178	727	AuAcAcccGuGuAuAAAcuTsT	A20179	728	AGUuAuAcACGGGUGuAUTsT	AD-12492
2113-2131	GP	A20180	729	uAcAGGcGuuAuAAuGcATsT	A20181	730	UGcAAUuAuAACGCCUGuATsT	AD-12493
1654-1672	GP	A20182	731	cAAuGcucAAcccAAuGcTsT	A20183	732	GcAUUUGGGUUGAGcAUUGTsT	AD-12494
1824-1842	GP	A20184	733	ccAAcGAGAcGAcucAAGcTsT	A20185	734	GCUUGAGUCGUCUCGUUGGTsT	AD-12495
1313-1331	GP	A20186	735	CCCGuGuAuAAAcuugAcATsT	A20187	736	UGUcAAGUuAuAcACGGGTsT	AD-12496
1873-1891	GP	A20188	737	GCuAcGcAccuuuucAAucTsT	A20189	738	GAUUGAAAAGGUGCGuAGCTsT	AD-12497
1953-1971	GP	A20190	739	GAccGGAcuGcuGuAucGATsT	A20191	740	UCGAuAcAGcAGUCCGGUCTsT	AD-12498
1964-1982	GP	A20192	741	uGuAucGAAccAcAuGAuTsT	A20193	742	AAUcAUGUGGUUCGAuAcATsT	AD-12499
329-347	VP30	A20194	743	AGGuGAGuAccGucAAucATsT	A20195	744	UGAUUGACGGuACUcACCUTsT	AD-12500
426-444	VP30	A20196	745	AAAGAcAuAuGuCCGAccuTsT	A20197	746	AGGUCGGAcAuAUGUCUUUTsT	AD-12501
842-860	VP30	A20198	747	uCUCGAAGuAuAuCAAcGATsT	A20199	748	UCGUUGAuAuACUUCGAGATsT	AD-12502
909-927	VP30	A20200	749	uGGGAccGAcAAucccuAATsT	A20201	750	UuAGGGAUUGUCGGUCCcATsT	AD-12503
523-541	VP30	A20202	751	UCCuACuAAucGCCCGuAATST	A20203	752	UuACGGGCGAUuAGuAGGATsT	AD-12504
429-447	VP30	A20204	753	GAcAuAuGuccGAccuuGATsT	A20205	754	UCAAGGUCGGAcAuAUGUCTsT	AD-12505
521-539	VP30	A20206	755	AcuccuAcuAAucGcccGuTsT	A20207	756	ACGGGCGAUuAGuAGGAGUTsT	AD-12506

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
903-921	VP30	A20208	757	cAACAAuGGGACCGACAAuTsT	A20209	758	AUUGUCGGUCCcAUUGUUGTsT	AD-12507
355-373	VP30	A20210	759	ccucAcAAGuGcGcGuuccTsT	A20211	760	GGAACGCGcACUUGUGAGGTsT	AD-12508
337-355	VP30	A20212	761	AccGuCAAucAAGGAGcGcTsT	A20213	762	GCGCUCCUUGAUUGACGGUTsT	AD-12509
908-926	VP24	A19262	763	CuGucGuuGAuucGAuccATsT	A19263	764	UGGAUCGAAUcAACGAcAGTsT	AD-11933
522-540	VP24	A19272	765	uuGucuuAAGcGAccucUGTsT	A19273	766	cAGAGGUCGCUuAAGAcAATsT	AD-11938
790-808	VP24	A19274	767	uuuGAuuGAAccuuAGcATsT	A19275	768	UGCuuAAGGGUUCcAAUcAAATsT	AD-11939
863-881	VP24	A20214	769	AAcAuGcGAACAcAAcGuGTsT	A20215	770	CACGUUGUGUUCGCAUGUUTsT	AD-12510
1102-1120	VP24	A20216	771	uGGGCCGGcGAAuuuuCCTsT	A20217	772	GGAAAAUUUCGCCGGCCCATsT	AD-12511
912-930	VP24	A20218	773	cGuuGAuucGAuccAAuAuTsT	A20219	774	AuAUUGGAUCGAAUcAACGTsT	AD-12512
954-972	VP24	A20220	775	AuGcucuACAuGucGuGAATsT	A20221	776	UUCACGAcAUGuAGAGcAUTsT	AD-12513
475-493	VP24	A20222	777	GGGAcGAuAcAAuCuAAuATsT	A20223	778	uAUuAGAUUGuAUCGUCCCTsT	AD-12414
1069-1087	VP24	A20224	779	ACCCGAcAAAucGGcAAuGTsT	A20225	780	CAUUGCCGAUUUGUCGGGUTsT	AD-12515
486-504	VP24	A20226	781	AucuAAuAucGcccAAAAATsT	A20227	782	UUUUUGGGCGAuAUuAGAUTsT	AD-12516
525-543	VP24	A20228	783	UCUuAAGcGAccucGuAATsT	A20229	784	UuAcAGAGGUCGCUuAAGATsT	AD-12517
867-885	VP24	A20230	785	uGcGAACAcAAcGuGucAATsT	A20231	786	UUGAcACGUUGUGUUCGcATsT	AD-12518
1028-1046	VP24	A20232	787	AuAAcucGAAcuAAcAuGGTsT	A20233	788	CcAUGUuAGUUCGAGUuAUTsT	AD-12519
471-489	VP24	A20234	789	CUAcGGGAcGAuAcAAucUTsT	A20235	790	AGAUGuAUCGUCCCGuAGTsT	AD-12520
1029-1047	VP24	A20236	791	uAAcucGAAcuAAcAuGGTsT	A20237	792	CCcAUGUuAGUUCGAGUuATsT	AD-12521
1948-1966	L	A20238	793	cAGuuAGAGGGAGuAGcuTsT	A20239	794	AAGCuACUCCCUcAACUGTsT	AD-12522
2003-2021	L	A20240	795	AuAuGAGuuuAcAGcAccuTsT	A20241	796	AGGUGCUGuAAACUcAuAUTsT	AD-12523
2005-2023	L	A20242	797	AuGAGuuuAcAGcAccuuuTsT	A20243	798	AAAGGUGCUGuAAACUcAUTsT	AD-12524
2070-2088	L	A20244	799	uGGAuGcAuuAuAcAAuccTsT	A20245	800	GGAUUGuAuAAUGcAUCcATsT	AD-12525
1959-1977	NP	A19278	801	AccuuGGAcGGAGcGAAAAATsT	A19279	802	UUUUCGCUCCGUCCAAGGUTsT	AD-11941

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1687-1705	NP	A19280	803	cAuuucccGGGccGAucuATsT	A19281	804	uAGAUCGGCCCGGAAUGTsT	AD-11942
1775-1793	NP	A19282	805	uGuuGuuGAcccGuAuGAuTsT	A19283	806	AUcAuACGGGUcAAcAAcATsT	AD-11943
384-402	NP	A19284	807	GuuuAccuGAGAGccuAcATsT	A19285	808	UGuAGGCUCUcAGGuAAACTsT	AD-11944
400-418	NP	A19286	809	AcAAcAuGGAuAAAcGGGuTsT	A19287	810	ACCCGUUuAUcAuGUUGUTsT	AD-11945
1773-1791	NP	A19288	811	GGuGuuGuuGAcccGuAuGTsT	A19289	812	CAuACGGGUcAAcAAcACCTsT	AD-11946
1964-1982	NP	A19290	813	GGAcGGAGcGAAAAGGuGTsT	A19291	814	cACCUUUUUCGUCUCGUCTsT	AD-11947
411-429	NP	A19292	815	AAAcGGGuGAGAGGuuAuTsT	A19293	816	AUGAACCUcUcACCCGUUUTsT	AD-11948
1815-1833	NP	A19294	817	GAcuAcGAGGAuucGGcuGTsT	A19295	818	cAGCCGAAUCCUCGuAGUCTsT	AD-11919
407-425	NP	A19296	819	GGAuAAAcGGGuGAGAGGuTsT	A19297	820	ACCUCUcACCCGUUuAUCCTsT	AD-11950
2405-2423	NP	A19298	821	uuAucAccuAAuGAGuGAuTsT	A19299	822	AUcACUcAUuAGGUGAuAATsT	AD-11951
409-427	NP	A19300	823	AuAAAcGGGuGAGAGGuucTsT	A19301	824	GAACCUCUcACCCGUUuAUTsT	AD-11952
1811-1829	NP	A19302	825	uccuGAcuAcGAGGAuucGTsT	A19303	826	CGAAUCCUCGuAGUcAGGATsT	AD-11953
408-426	NP	A19304	827	GAuAAAcGGGuGAGAGGuuTsT	A19305	828	AACCUCUcACCCGUUuAUCTsT	AD-11954
1958-1976	NP	A19306	829	GAccuuGGAcGGAGcGAAATsT	A19307	830	UUUCGUCUCGUcAAGGUCTsT	AD-11955
1973-1991	NP	A19308	831	GAAAAAGGuGccGGAGuuGTsT	A19309	832	CAACUCCGGCACCUUUUUCTsT	AD-11956
1810-1828	NP	A19310	833	AuCCuGACuAcGAGGAuuCTsT	A19311	834	GAAUCCUCGuAGUCAGGAUTsT	AD-11957
1953-1971	NP	A19312	835	GAuccGAccuuGGAcGGAGTsT	A19313	836	CUCCGUcAAGGUCGGAUCTsT	AD-11958
1692-1710	NP	A19314	837	CCCGGGccGAucuAuGAuGTsT	A19315	838	CAUCAuAGAUCGGCCCGGTsT	AD-11959
197-215	VP35	A19316	839	AccGGcAAAAuAccGcuAATsT	A19317	840	UuAGCGGuAUUUUGCCGGUTsT	AD-11960
196-214	VP35	A19318	841	GAccGGcAAAAuAccGcuATsT	A19319	842	uAGCGGuAUUUUGCCGGUCTsT	AD-11961
409-427	VP35	A19320	843	AucACuAGAAGGucGAGuATsT	A19321	844	uACUCGACCUUCuAGUGAUTsT	AD-11962
476-494	VP35	A19322	845	AuAucAuCCCuGAAuGcATsT	A19323	846	UGCGAUUcAGGGAUGAuAUTsT	AD-11963
611-629	VP35	A19324	847	CCAucAuuGuAcGAGGAuGTsT	A19325	848	cAUCCUCGuAcAAUGAUGGTsT	AD-11964

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
645-663	VP35	A19326	849	AAuuGAAAGAUccGAacGGTsT	A19327	850	CCGUUCGGAUCUUUcAAUUTsT	AD-11965
726-744	VP35	A19328	851	AGGAAAuuucGGGcGACCTsT	A19329	852	GGUCGCCCCGAAuuuUCCUTsT	AD-11966
1130-1148	VP35	A19330	853	GuAAGcucAuuuuGcGAuGTsT	A19331	854	cAUCGcAAAAUGAGCUuACTsT	AD-11967
729-747	VP35	A19332	855	AAAuuucGGGcGAccuuATsT	A19333	856	uAAGGUCGCCCCGAAUUUUTsT	AD-1168
606-624	VP35	A19334	857	CAGGCCCAuCAuuGuAcGATsT	A19335	858	UCGuACAAUGAUGGGCCUGTsT	AD-11969
256-274	VP35	A19336	859	cccGAuAAccAuAuAuAGuTsT	A19337	860	ACuAAuAAUGGUuAUCGGGTsT	AD-11970
478-496	VP35	A19338	861	AuCAuCCCuGAuCGcAGcTsT	A19339	862	GCUGCGAUUcAGGGAUGAUTsT	AD-11971
724-742	VP35	A19340	863	uGAGGAAAuuucGGGcGATsT	A19341	864	UCGCCCCGAAUUUCCUCATsT	AD-11972
644-662	VP35	A19342	865	AAAuuGAAAGAUccGAacGTsT	A19343	866	CGUUCGGAUCUUUcAAUUTsT	AD-11973
1239-1257	VP35	A19344	867	AuccuAAucAAuuGAuAAuTsT	A19345	868	AUuAUcAAUUGAUuAGGAUTsT	AD-11974
1052-1070	VP35	A19346	869	AucGAuAGAGGuuGGGucTsT	A19347	870	AGACCCaACCUCuAUCGAUTsT	AD-11975
429-447	VP40	A19348	871	GcAAuuAuGcucGcAucuuTsT	A19349	872	AAGAUGCGAGcAuAAUUGCTsT	AD-11976
1189-1207	VP40	A19350	873	AAAAuGuAcuAAucGGGucTsT	A19351	874	GACCCGAUuAGuAcAUUUUTsT	AD-11977
1190-1208	VP40	A19352	875	AAAuGuAcuAAucGGGucATsT	A19353	876	UGACCCGAUuAGuAcAUUUUTsT	AD-11978
373-391	VP40	A19354	877	GGuuGccAcucGGAAuuGcTsT	A19355	878	GcAAUUCGAGUGGcAACCTsT	AD-11979
439-457	VP40	A19356	879	ucGcAucuuAuAcGAucAcTsT	A19357	880	GUGAUCGuAuAAGAUGCGATsT	AD-11980
441-459	VP40	A19358	881	GcAucuuAuAcGAucAcccTsT	A19359	882	GGGUGAUCGuAuAAGAUGCTsT	AD-11981
1121-1139	VP40	A19360	883	AuAGcAAcucAAucGAcuuTsT	A19361	884	AAGUCGAUUGAGUUGCuAUTsT	AD-11982
1127-1145	VP40	A19362	885	AcucAAucGAcuuuuAGGATsT	A19363	886	UCCuAAAAGUCGAUUGAGUTsT	AD-11983
1193-1211	VP40	A19364	887	uGuAcuAAucGGGucAAGGTsT	A19365	888	CCUUGACCCGAUuAGuAcATsT	AD-11984
1298-1316	VP40	A19366	889	AcAuGcAuAAGcGAuccAuTsT	A19367	890	AUGGAUCGCUuAUGcAUGUTsT	AD-11985
1307-1325	VP40	A19368	891	AGcGAuccAuAcuucGcccTsT	A19369	892	GGGCGAAGuAUGGAUCGCUTsT	AD-11986
361-379	VP40	A19370	893	AAuucccuAuuuGGuuGccTsT	A19371	894	GGCAACCAAAuAGGGAUUUTsT	AD-11987

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
437-455	VP40	A19372	895	GcucGcAucuuAuAcGAucTsT	A19373	896	GAUCGuAuAAGAUGCGAGCTsT	AD-11988
857-875	VP40	A19374	897	GAGuAuCAuuGGGAucGAGTsT	A19375	898	CUCGAUCCcAAUGAuACUCTsT	AD-11989
484-502	VP40	A19376	899	ucGuuAGAGuGAAucGAcuTsT	A19377	900	AGUCGAUUCACUCuAACGATsT	AD-11990
1845-1863	GP	A19378	901	GGAGcuGcGGAcAuAuAccTsT	A19379	902	GGuAuAUGUCCGcAGCUCCTsT	AD-11991
254-272	GP	A19380	903	GAGAuGAccAGcuAGucuTsT	A19381	904	AGAcuAGCUGGUcAAUCUCTsT	AD-11992
461-479	GP	A19382	905	ccGGAcGGGAGcGAAuGcuTsT	A19383	906	AGcAUUCGCUCcCGGCTsT	AD-11993
466-484	GP	A19384	907	cGGGAGcGAAuGcuuAcccTsT	A19385	908	GGGuAAGcAUUCGCUCcCGTtsT	AD-11994
933-951	GP	A19386	909	uAAuuuGGAcAcuAGAuGcTsT	A19387	910	GcAUCuAGUGUCcAAAUuATsT	AD-11995
1045-1063	GP	A19388	911	uuAucGcucAAcGAGAcAGTsT	A19389	912	CUGUCUCGUUGAGCGAuAATsT	AD-11996
1100-1118	GP	A19390	913	GAAGAAucuccGAccGGGcTsT	A19391	914	GCCCGGUCGGAGAuUCuUCTsT	AD-11997
1102-1120	GP	A19392	915	AGAAucuccGAccGGGccATsT	A19393	916	UGGCCCGGUCGGAGAUUCUTsT	AD-11998
1191-1209	GP	A19394	917	AAcAAcAuGccGuCUCAGTsT	A19395	918	CUGAGACGGcAAUGUUGUUTsT	AD-11999
1203-1221	GP	A19396	919	GuCuCAGAAuuCGACAGAAtsT	A19397	920	uUCuGUCGAAuUCuGAGACTsT	AD-12000
1844-1862	GP	A19398	921	cGGAGcuGcGGAcAuAuAcTsT	A19399	922	GuAuAUGUCCGcAGCUCcGTsT	AD-12001
255-273	GP	A19400	923	AGAuugACCAGcuAGuCuGTsT	A19401	924	cAGAcuAGCUGGUCAAUCUTsT	AD-12002
1212-1230	GP	A19402	925	uucGAcAGAAGGucGAAGATsT	A19403	926	UCUUCGACCUUCUGUCGAATsT	AD-12003
1706-1724	GP	A19404	927	GGAucccGuAcuuuGGAccTsT	A19405	928	GGUCcAAAGuACGGGAUCCTsT	AD-12004
125-143	GP	A19406	929	CUuAGccuAcuccAAuuGcTsT	A19407	930	GcAAUUGGAGuAGGCuAAGTsT	AD-12005
264-282	GP	A19408	931	AGcuAGucuGcAAGGAucATsT	A19409	932	UGAUCCUUGcAGAcuAGCUTsT	AD-12006
332-350	GP	A19410	933	AGcGGAGuAucuAcuGAuATsT	A19411	934	uAUcAGuAGAuACUCCGCUTsT	AD-12007
464-482	GP	A19412	935	GAcGGGAGcGAAuGcuuAcTsT	A19413	936	GuAAGcAUUCGCUCcCGUCTsT	AD-12008
1210-1228	GP	A19414	937	AAuucGAcAGAAGGuCGAATsT	A19415	938	uUCGACCuUCuGUCGAAUUTsT	AD-12009
1213-1231	GP	A19416	939	ucGAcAGAAGGucGAAGAGTsT	A19417	940	CUCuUCGACCuUCuGUCGATsT	AD-12010

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1850-1868	GP	A19418	941	uGcGGAcAuAuAccAuAcuTsT	A19419	942	AGuAUGGuAuAUGUCCGcATsT	AD-12011
124-142	GP	A19420	943	UCUuAGCCuACuCCAuuGTsT	A19421	944	CAAUUGGAGuAGGCuAAGATsT	AD-12012
1044-1062	GP	A19422	945	uuuAucGcucAAcGAGAcATsT	A19423	946	UGUCUCGUUGAGCGAuAAATsT	AD-12013
265-283	GP	A19424	947	GcuAGuCuGcAAGGAuCAuTsT	A19425	948	AUGAUCCUUGcAGACuAGCTsT	AD-12014
361-379	VP30	A19426	949	uAGAGucccuAcGGuuuucTsT	A19427	950	GAAAACCGuAGGGACUCuATsT	AD-12015
324-342	VP30	A19428	951	cAAcAGAcuAccGuAGuAGTsT	A19429	952	CuACuACGGuAGUCUGUUGTsT	AD-12016
994-1012	VP30	A19430	953	AGGccuAcGcuuAcuuGccTsT	A19431	954	GGcAAGuAAGCGuAGGCCUTsT	AD-12017
248-266	VP30	A19432	955	AGGAuuAcGuGccGAcCTsT	A19433	956	GGUCGGcACGUGAAUCCUTsT	AD-12018
491-509	VP30	A19434	957	cuuGAAAGccuAAccGAccTsT	A19435	958	GGUCGGUuAGGCUUUCaAGTsT	AD-12019
322-340	VP30	A19436	959	AAcAAcAGAcuAccGuAGuTsT	A19437	960	ACuACGGuAGUCUGUUGUTsT	AD-12020
323-341	VP30	A19438	961	AcAAcAGAcuAccGuAGuATsT	A19439	962	uACuACGGuAGUCUGUUGUTsT	AD-12021
517-535	VP30	A19440	963	ccuAcuucuuAuAGcAcGGTsT	A19441	964	CCGUGCuAuAAGAAGuAGGTsT	AD-12022
295-313	VP30	A19442	965	GAcAAGAuccAuuucccGGTsT	A19443	966	CCGGGAAuAGGAUCuuGUCTsT	AD-12023
229-247	VP30	A19444	967	ucGuGAGcGcGGGAGAuATsT	A19445	968	UGAUCUCCGCGCUcACGATsT	AD-12024
251-269	VP30	A19446	969	AAuucAcGuGCCGAccAGcTsT	A19447	970	GCUGGUCGGcACGUGAAUUTsT	AD-12025
340-358	VP30	A19448	971	uAGucGAAGuACuucGcAATsT	A19449	972	UUGCGAAGuACUUCGACuATsT	AD-12026
1350-1368	VP30	A19450	973	ucccuAGAAGcGuuGAAucTsT	A19451	974	GAUUCaACGCUUCuAGGGATsT	AD-12027
1057-1075	VP24	A19452	975	uAuGGGuuAucuuGucGAGTsT	A19453	976	CUCGAcAAGAuAACCCcAuATsT	AD-12028
878-896	VP24	A19454	977	AAcAuGAGAAcucAAcGAGTsT	A19455	978	CUCGUUGAGUUCUcAUGUUTsT	AD-12029
1056-1074	VP24	A19456	979	AuAuGGGuuAucuuGucGATsT	A19457	980	UCGAcAAGAuAACCCcAuAUTsT	AD-12030
1137-1155	VP24	A19458	981	uAcuAcAuGAuAcGAcACuTsT	A19459	982	AGUGUCGAUUCaUGuAGuATsT	AD-12031
1099-1117	VP24	A19460	983	GAuGGAuAuAcGAcAcCCuTsT	A19461	984	AGGGUGUCGuAuAUCcAUCTsT	AD-12032
1591-1609	VP24	A19462	985	AGcccAAAuuAAcAcGGuATsT	A19463	986	uACCGUGUuAAUUUGGGCUTsT	AD-12033

-continued

position		double overhang design sense strand			antisense strand			duplex name
in target	Target	name	Seq ID	sequence (5'-3')	name	Seq ID	sequence (5'-3')	
1094-1112	VP24	A19464	987	ucuGcGAuGGGAuAuAcGAcTsT	A19465	988	GUCGuAuAUCcAUCGcAGATsT	AD-12034
1135-1153	VP24	A19466	989	cuuAcuAcAuGAAucGAcATsT	A19467	990	UGUCGAUUCaUGuAGuAAGTsT	AD-12035
152-170	VP24	A19468	991	cuAGGCUAGGGuuuAuAGuTsT	A19469	992	ACuAuAAACCCuAGCCuAGTsT	AD-12036
624-642	VP24	A19470	993	AccAAAAGGGuAuuAcccuTsT	A19471	994	AGGGuAAuACCCUUUUGGUTsT	AD-12037

TABLE 3

duplex name	Ebola Zaire GFP assay % reduction			Ebola Zaire plaque % reduction vs no siRNA	Ebola Sudan IF assay % reduction			Ebola Sudan plaque % reduction vs no siRNA
	100 nM	10 nM	1 nM		100 nM	10 nM	1 nM	
AD-11542	90.7	80.1	55.2	55.24	-5.3	-1.5	-11.8	-103.39
AD-11543	62.7	55.9	47.8	55.83	-11.9	10.2	-9.0	-7.12
AD-11544	63.5	32.8	41.0	42.06	22.5	34.6	61.1	-16.95
AD-11545	73.0	71.8	61.4	47.23	61.5	71.9	71.8	39.41
AD-11546	94.5	88.3	72.5	86.12	-8.8	-73.8	5.5	55.93
AD-11547	79.8	62.8	48.6	36.44	-20.7	15.8	-4.0	18.64
AD-11548	-15.4	-13.0	-15.6		19.3	29.2	49.3	
AD-11549	3.8	13.8	16.1	44.12	63.7	51.5	90.4	94.38
AD-11550	17.5	-1.2	-10.5		-41.1	-39.8	-29.4	
AD-11551	-14.2	-20.4	-21.3		-32.6	-23.9	-30.6	
AD-11552	-14.7	-11.9	-6.5		-22.7	-9.5	11.9	
AD-11553	4.7	7.4	13.9	27.67	44.5	40.9	44.6	80.33
AD-11554	-12.6	-20.5	-32.1		-12.6	-17.7	-8.0	
AD-11555	-20.8	-22.2	-36.7		-9.8	-8.9	0.8	
AD-11556	6.4	-11.6	-21.7	29.03	11.1	37.1	34.7	-17.37
AD-11557	30.6	22.0	-8.7	-175.73	41.5	48.3	78.3	82.20
AD-11558	3.0	1.2	-33.7	72.97	-23.5	-18.2	-30.8	82.49
AD-11559	-29.1	-38.0	-42.9		-2.4	-37.3	-26.5	
AD-11560	-42.0	-46.1	-54.6		19.8	0.7	19.1	
AD-11561	-16.4	-16.6	-15.9	37.38	50.5	61.2	57.4	-129.83
AD-11562	12.0	-13.9	-63.1	61.33	-35.3	-46.2	-42.4	63.56
AD-11563	-45.3	-51.6	-61.7		-45.3	-54.7	-42.9	
AD-11564	-19.0	-31.3	-44.0		-38.5	-32.3	-0.1	
AD-11565	-27.0	-4.4	-10.7	50.32	6.7	47.4	52.9	26.95
AD-11566	6.8	6.0	-0.5		32.0	18.2	0.7	
AD-11567	-1.1	-5.7	0.3		3.1	-4.6	14.4	
AD-11568	10.3	7.9	11.4		-0.7	27.6	25.6	
AD-11569	17.6	22.0	15.4	33.01	47.1	51.3	63.7	-85.01
AD-11570	23.8	5.9	-15.4	98.14	9.3	-5.5	-15.3	-217.80
AD-11571	-27.5	-23.9	-32.4		-8.5	-13.0	10.8	
AD-11572	-35.2	-25.7	-26.1		4.3	8.8	18.6	
AD-11573	-8.3	16.7	15.7	46.60	55.9	76.5	37.8	-22.05
AD-11574	1.1	-12.0	-27.6		12.5	-2.9	-11.3	
AD-11575	-29.9	-30.0	-35.9		1.6	27.3	-11.4	
AD-11576	-9.4	-8.2	-20.8		0.1	34.6	3.1	
AD-11577	-8.5	-0.1	13.7		45.2	31.5	23.2	
AD-11578	9.8	17.0	4.1		-40.1	-56.3	-53.3	
AD-11579	15.1	12.2	-7.8		-47.6	-37.3	-18.5	
AD-11580	-4.4	2.3	4.1	76.38	14.2	4.9	33.6	12.97
AD-11581	10.6	2.5	3.4	-26.97	57.4	65.1	81.7	-109.94
AD-11582	10.6	4.3	-33.2	-1.17	-99.7	-93.3	-90.8	-533.33
AD-11583	-16.6	-18.4	-28.3		-85.8	-72.4	-55.3	
AD-11584	-24.1	-12.0	-18.7	59.87	-1.0	54.0	16.5	1.61
AD-11585	-7.2	-9.6	9.1	51.46	37.5	19.1	45.7	82.71
AD-11586	-6.8	-0.7	-9.1		-81.1	-87.9	-86.9	
AD-11587	6.1	-0.4	0.9		-80.7	-6.2	-51.8	
AD-11588	20.9	10.4	-9.5	99.40	-12.9	-0.1	20.5	-117.23
AD-11589	28.7	24.3	16.1	35.75	18.2	38.7	54.2	88.14
AD-11590	20.7	28.8	3.0	98.81	-38.5	-55.2	20.9	44.92
AD-11591	-18.3	-29.6	-27.0		-3.1	-40.1	-30.1	

TABLE 3-continued

duplex name	Ebola Zaire GFP assay % reduction			Ebola Zaire plaque % reduction vs no siRNA	Ebola Sudan IF assay % reduction			Ebola Sudan plaque % reduction vs no siRNA
	100 nM	10 nM	1 nM		100 nM	10 nM	1 nM	
AD-11592	-11.3	-14.7	-12.3		-36.3	-38.7	-46.1	
AD-11593	3.8	5.0	-12.3		1.8	-15.8	4.6	
AD-11594	33.4	4.9	-3.2	99.07	-25.2	-31.3	-21.7	7.63
AD-11595	-30.9	-37.7	-54.2		-11.6	-2.7	-2.7	
AD-11596	-8.2	-14.8	-3.3		-7.8	26.8	3.5	
AD-11597	13.0	13.0	11.7		21.7	28.2	25.2	83.00
AD-11598	40.0	-1.6	-10.4	98.91	-30.2	-33.7	-17.7	0.00
AD-11599	13.0	24.6	-18.3	98.79	-20.9	-22.8	-50.8	34.32
AD-11600	16.7	2.8	-8.1	98.52	-4.3	-28.5	6.8	-120.34
AD-11601	10.4	25.3	2.0	71.20	23.0	6.3	7.9	-652.16
AD-11602	39.7	45.0	24.4	38.73	47.4	37.7	54.8	48.13
AD-11603	41.2	42.4	36.0	54.85	38.2	35.7	43.8	73.04
AD-11604	39.6	36.3	31.6	-1.51	37.0	56.6	52.8	84.65
AD-11605	50.2	38.8	21.2	60.19	48.1	58.6	77.6	66.95
AD-11606	41.5	36.8	1.4	89.23	53.2	33.2	37.0	79.32
AD-11607	-12.8	2.7	-9.9	38.83	29.0	8.3	21.4	-13.98
AD-11608	10.9	-6.7	6.1	14.24	39.5	49.3	42.9	-75.07
AD-11609	27.3	32.9	29.0	51.46	36.8	40.7	50.2	20.81
AD-11610	23.9	19.7	16.0	23.95	37.7	70.3	49.6	97.93
AD-11611	5.6	11.9	16.1		30.7	8.5	15.6	
AD-11612	-2.9	-4.0	-14.4		9.9	21.9	10.0	
AD-11613	13.4	32.7	23.7		-2.6	17.5	44.4	
AD-11614	9.0	14.3	10.3		21.9	9.9	-29.6	
AD-11615	-5.9	22.3	-0.9		-24.8	-16.9	-21.2	
AD-11616	2.3	5.5	11.0		-10.4	-7.6	26.4	
AD-11617	14.3	5.3	-1.9	-617.15	39.8	62.5	91.5	14.55
AD-11618	-2.5	-4.3	-1.2		-11.0	-26.8	-28.6	
AD-11619	-8.3	-37.8	-30.2		-33.9	-43.8	-23.9	
AD-11620	11.0	-19.2	-37.5		-30.5	-12.0	0.7	
AD-11621	-29.1	-4.5	4.7	-255.66	60.9	64.7	78.8	74.06
AD-11622	11.9	15.0	4.2		9.8	-29.1	-15.8	
AD-11623	7.7	-17.8	-24.6		-27.3	-48.1	-19.4	
AD-11624	-16.3	-36.6	-45.8		-36.1	-17.2	-8.3	
AD-11625	45.0	11.0	0.3	37.98	19.5	48.2	43.1	-23.73
AD-11626	-3.6	13.2	11.9		-25.4	-81.4	-80.3	
AD-11627	34.1	24.7	21.1	48.69	-46.8	-73.0	-54.8	12.71
AD-11628	-8.9	-1.4	-13.8	9.60	-9.5	6.1	35.2	59.80
AD-11629	22.9	23.0	11.8	-73.11	65.3	71.2	75.1	3.39
AD-11630	12.2	19.0	-2.0		-99.9	-123.8	-85.0	
AD-11631	-10.4	-15.2	-18.3		-127.0	-87.2	-59.2	
AD-11632	-50.7	-49.9	-49.4	-128.91	-32.4	1.3	29.8	51.76
AD-11633	-39.4	-10.2	1.8	-19.31	20.6	42.9	57.4	49.57
AD-11634	2.2	12.3	-14.1		-55.8	-129.3	-136.9	
AD-11635	-44.3	-46.0	-29.5		-60.2	-89.0	-66.2	
AD-11636	-48.8	-44.7	-41.1		-35.4	-17.4	7.3	
AD-11637	-36.8	-27.2	-17.3	59.55	34.3	33.8	52.8	80.40
AD-11638	-8.4	8.5	-3.9		-9.4	12.4	28.9	
AD-11639	-47.6	-1.7	36.0	23.31	2.9	20.2	12.6	-97.46
AD-11640	69.4	39.4	34.6	75.14	66.0	71.5	82.5	49.15
AD-11641	42.6	15.2	15.9	65.14	73.1	81.9	87.7	96.50
AD-11642	24.8	21.4	32.7	-42.93	89.8	87.4	87.4	4.03
AD-11643	37.7	53.8	52.7	89.14	94.3	94.2	97.5	-22.03
AD-11644	54.1	17.7	5.5	94.29	26.7	38.5	51.2	31.78
AD-11645	5.9	2.5	-2.5	51.97	64.7	68.1	80.1	93.34
AD-11646	-1.7	2.1	7.9	59.55	72.6	76.8	79.0	92.72
AD-11647	8.4	8.8	37.5	57.28	75.6	81.7	93.7	74.71
AD-11648	56.9	19.2	8.1	66.10	-15.4	-9.9	13.0	71.19
AD-11649	-3.3	-6.9	-3.5	65.47	36.0	54.2	64.9	82.84
AD-11650	1.9	-2.5	-1.5	17.28	58.7	65.3	59.6	2.31
AD-11651	6.7	9.4	30.9	65.70	70.3	87.1	91.2	83.96
AD-11652	55.8	27.7	10.3		-56.8	59.6	68.4	33.90
AD-11653	12.4	13.8	10.9	86.84	66.4	70.4	75.8	2.02
AD-11654	13.8	7.2	10.7		76.1	73.8	73.8	-123.92
AD-11655	9.1	14.6	40.5	-7.01	82.4	83.9	92.6	-146.11
AD-11656	39.1	9.6	-1.0	81.71	-21.9	30.7	19.0	20.34
AD-11657	-1.1	1.1	0.2		-3.1	20.7	48.9	
AD-11658	7.3	0.9	1.5	86.84	55.5	50.8	41.3	93.52
AD-11659	2.9	4.1	28.0	78.79	46.4	51.1	66.6	79.47
AD-11660	23.8	13.0	0.9	80.57	11.0	20.8	-57.0	90.51
AD-11661	-6.4	-8.0	-6.1		-37.1	-16.9	19.1	
AD-11662	-1.6	-7.9	-7.9		10.9	10.7	27.7	
AD-11663	1.6	5.6	28.1		25.4	20.4	30.8	
AD-11664	11.9	7.6	3.9	50.16	54.4	63.1	66.8	77.95

TABLE 3-continued

duplex name	Ebola Zaire GFP assay % reduction			Ebola Zaire plaque % reduction vs no siRNA	Ebola Sudan IF assay % reduction			Ebola Sudan plaque % reduction vs no siRNA
	100 nM	10 nM	1 nM		100 nM	10 nM	1 nM	
AD-11665	5.1	1.2	0.2	15.10	52.7	65.2	62.3	-13.26
AD-11666	0.0	-3.5	5.3	5.72	67.9	80.3	80.3	79.54
AD-11667	3.7	2.9	8.0	47.73	83.6	80.4	90.6	84.70
AD-11668	16.3	-0.3	-1.3		-91.6	-4.4	2.8	
AD-11669	-1.0	3.8	0.3		3.4	33.5	43.3	
AD-11670	-0.1	-4.6	-2.6	35.28	42.7	43.6	62.6	-5.48
AD-11671	0.1	1.1	16.0	80.58	61.0	65.1	72.7	13.54
AD-11672	8.2	-0.1	-2.2		-69.9	25.3	-57.8	
AD-11673	1.7	0.9	3.2		-46.1	-22.7	1.5	
AD-11674	-4.2	-3.0	-6.0		34.4	14.8	32.2	
AD-11675	-0.6	0.6	18.4		11.1	45.1	59.7	
AD-11676	22.1	11.5	7.7	78.86	75.7	86.4	78.6	16.95
AD-11677	5.6	-3.1	3.2		86.1	78.0	87.1	1.27
AD-11678	-2.3	-7.8	5.0	87.06	86.4	86.6	86.6	-141.35
AD-11679	5.6	7.3	34.4	28.26	88.3	88.6	90.9	-79.25
AD-11680	30.4	7.9	-0.5	75.62	-38.9	13.4	-88.3	72.03
AD-11681	-3.7	-10.1	-0.2		14.4	25.0	24.2	
AD-11682	-10.0	-11.0	-7.7		-39.9	53.8	57.2	
AD-11683	-6.4	6.6	32.3	92.99	59.6	53.3	48.2	57.85
AD-11684	31.5	14.4	2.9	33.26	-156.0	-119.6	-198.2	87.80
AD-11685	0.0	-6.5	-9.5	-6.15	-43.0	-107.3	-93.7	-20.61
AD-11686	-12.0	-7.9	-4.7	88.12	-134.6	-69.2	-151.8	43.11
AD-11687	-4.8	6.3	29.0		-77.4	-41.4	-79.0	
AD-11688	40.0	28.5	26.7	-15.66	73.4	79.8	88.0	66.95
AD-11689	34.4	27.1	32.4	57.33	83.4	74.8	86.5	85.59
AD-11690	24.0	30.2	42.1	71.71	82.5	89.4	89.4	83.05
AD-11691	47.9	44.1	55.3	87.43	92.0	93.9	97.0	70.96
AD-11694	44.3	8.5	5.7		2.1	4.8	13.6	
AD-11695	1.5	0.5	-5.3		42.1	36.7	46.0	
AD-11696	5.1	4.8	10.8		49.2	56.9	76.3	
AD-11698	18.4	20.4	45.5	59.09	69.5	79.5	82.0	54.66
AD-11700	30.4	8.5	-2.1	84.00	-151.2	-108.0	-67.5	4.52
AD-11704	-7.2	-6.1	-9.9		-37.2	-56.5	-32.4	
AD-11705	-2.9	-2.4	-4.7		-7.3	0.9	-14.8	
AD-11706	8.5	21.1	99.9		21.5	-8.9	33.9	
AD-11707	41.8	25.9	21.9	90.69	79.3	84.2	85.1	63.14
AD-11708	27.2	27.4	28.1	83.92	80.0	81.1	83.7	24.58
AD-11710	31.0	37.7	43.0	77.54	86.9	89.6	89.6	99.70
AD-11711	33.8	36.5	48.2	73.43	91.1	89.8	93.9	-34.75
AD-11712	81.1	20.2	8.0	79.71	22.6	-4.3	27.5	-13.56
AD-11713	-8.6	-4.0	-2.1		39.1	39.1	51.8	
AD-11714	-8.7	-8.3	-4.3	63.27	45.0	66.1	74.5	39.48
AD-11715	0.8	14.9	40.7	-92.23	79.0	82.0	86.9	1.15
AD-11716	32.4	8.3	1.8	83.05	-46.0	-51.4	-11.3	-577.97
AD-11717	2.2	-0.5	-0.6		-1.5	-13.0	-14.3	
AD-11718	-8.1	-11.9	-19.8		2.4	18.8	29.2	
AD-11719	4.2	11.6	33.8	57.93	55.3	56.6	74.0	-2.02
AD-11720	29.8	17.3	10.4	77.05	86.4	91.8	95.6	-373.73
AD-11721	14.6	13.7	12.4	82.56	90.8	91.1	95.1	-31.70
AD-11722	17.0	15.1	23.7	82.96	94.1	91.7	91.7	-112.39
AD-11723	16.0	13.5	30.5	90.99	94.1	95.5	95.7	-55.62
AD-11724	33.9	10.4	5.1	99.98	63.2	73.2	73.7	-1.27
AD-11725	6.4	-2.0	0.0	93.62	85.5	92.4	89.5	-172.62
AD-11726	2.4	0.2	0.2	58.58	90.3	91.9	94.4	-2.54
AD-11727	-1.0	2.8	20.2	4.72	93.9	94.5	98.3	49.86
AD-11728	30.7	12.9	4.0	73.43	26.9	27.5	41.0	30.51
AD-11729	5.1	1.5	-4.3	26.86	51.9	65.4	58.9	-20.46
AD-11730	2.6	1.3	0.8	24.70	60.2	67.7	71.6	52.02
AD-11731	4.9	3.3	24.1	79.94	66.8	67.3	89.0	96.16
AD-11732	33.6	9.3	6.0	63.43	63.0	79.9	87.5	70.34
AD-11733	2.2	2.8	-0.7	75.57	84.2	89.7	85.1	-9.51
AD-11734	8.6	6.7	10.8	90.52	90.8	90.6	90.6	-119.16
AD-11735	6.9	9.3	31.4	-22.87	90.8	88.5	94.5	-77.23
AD-11736	31.5	9.9	3.0	21.14	29.9	39.5	45.3	-611.86
AD-11737	-8.0	-3.4	-1.7	-183.17	78.3	88.4	82.5	14.41
AD-11738	-0.2	1.5	-4.7	-102.05	85.3	86.4	80.4	10.52
AD-11739	7.4	3.8	21.5	52.75	53.8	55.2	83.5	89.27
AD-11740	30.0	11.7	-0.4	53.43	-87.3	-64.8	12.1	38.98
AD-11741	-3.5	-6.1	-2.9	42.39	27.2	47.7	36.1	-32.56
AD-11742	-1.0	-1.9	-6.0	40.35	49.9	56.9	20.5	12.25
AD-11743	0.1	6.6	26.9	37.35	46.3	24.7	71.2	26.08
AD-11744	40.5	24.2	21.0	75.24	42.9	52.0	90.3	32.20
AD-11745	26.3	23.1	23.3	55.43	64.3	74.1	84.5	54.24

TABLE 3-continued

duplex name	Ebola Zaire GFP assay % reduction			Ebola Zaire plaque % reduction vs no siRNA	Ebola Sudan IF assay % reduction			Ebola Sudan plaque % reduction vs no siRNA
	100 nM	10 nM	1 nM		100 nM	10 nM	1 nM	
AD-11746	7.2	0.9	-4.8	89.82	42.3	74.8	8.6	33.49
AD-11747	1.6	-3.5	-5.4		54.9	5.8	25.4	
AD-11748	-4.3	0.5	0.9		30.3	10.9	10.9	
AD-11749	8.4	3.9	1.0		41.4	-3.7	54.3	
AD-11750	4.8	2.1	0.0		44.9	30.4	71.5	
AD-11751	3.3	0.6	-2.2		34.3	-18.6	29.4	
AD-11752	17.7	6.5	3.2	-27.63	19.8	19.6	-0.1	61.08
AD-11753	1.7	1.2	0.9		-62.5	-40.7	28.2	
AD-11754	3.7	1.8	0.7		69.9	91.1	44.9	
AD-11755	37.7	26.6	10.8	-58.93	10.1	0.6	28.8	47.05
AD-11756	-1.0	-0.6	0.3		-39.8	-16.7	-5.7	
AD-11757	0.2	0.0	1.8		-46.4	-12.7	-2.7	
AD-11758	18.9	11.7	7.5	21.83	27.3	12.4	-9.8	70.02
AD-11759	26.1	15.1	18.8	-595.93	-6.9	-3.8	-22.6	55.98
AD-11760	21.9	19.5	6.0	11.60	-61.2	6.6	6.6	22.49
AD-11761	18.1	8.6	6.4	-67.30	39.7	43.4	58.0	71.77
AD-11762	15.4	9.7	4.2	-335.24	38.7	-86.0	12.4	73.68
AD-11763	6.8	7.3	3.7		-3.6	-0.7	8.7	
AD-11764	5.1	5.7	0.9		-34.9	28.7	18.2	
AD-11765	14.8	8.0	4.3	5.98	32.1	27.3	58.4	76.08
AD-11766	2.9	3.5	3.9		13.4	-6.1	-68.1	
AD-11767	24.1	15.1	5.5	7.18	32.9	-27.7	0.2	9.57
AD-11768	19.2	16.5	4.3	30.34	-92.5	-3.0	4.0	55.98
AD-11769	5.6	2.3	2.6		6.7	25.7	43.9	
AD-11770	14.0	7.2	2.2	-16.41	19.1	34.2	35.3	49.60
AD-11771	2.8	2.1	1.3		37.7	19.1	15.6	
AD-11772	4.0	2.2	2.6		70.0	52.3	52.3	
AD-11773	1.3	-0.6	-0.7		52.0	65.8	65.4	
AD-11774	2.0	1.5	0.5		73.1	-18.3	6.5	
AD-11775	2.3	0.3	-0.2		57.8	49.1	5.5	
AD-11776	2.7	0.4	1.6		59.6	48.4	41.0	
AD-11777	3.6	1.3	0.0		10.4	47.2	53.8	
AD-11778	0.9	-0.9	-0.8		40.4	-13.5	-17.7	
AD-11779	6.3	1.7	-0.5		42.1	2.9	-6.9	
AD-11780	5.8	0.1	0.1		29.5	-17.6	20.3	
AD-11781	2.3	1.0	-0.2		8.8	33.5	39.1	
AD-11782	-1.7	-4.0	-4.1		48.1	44.5	56.1	
AD-11783	-3.5	-2.1	-1.5		34.8	16.7	44.2	
AD-11784	5.7	3.7	-1.0		50.4	59.1	59.1	
AD-11785	2.8	1.0	1.5		75.3	88.0	81.7	
AD-11786	14.9	6.5	-2.9	97.46	-7.0	-34.8	-6.5	66.19
AD-11787	-0.3	0.0	-2.9		9.0	-8.6	18.2	
AD-11788	-0.1	0.5	1.3		23.5	11.8	55.0	
AD-11789	5.1	5.7	3.1		57.8	45.6	62.5	
AD-11790	5.7	0.0	-4.0		18.8	-12.3	-1.9	
AD-11791	-1.5	-2.6	-4.6		42.1	-27.4	10.4	
AD-11792	-0.9	1.8	1.5		27.1	30.8	24.6	
AD-11793	2.2	3.7	4.4		6.7	34.8	44.3	
AD-11794	0.4	-0.5	-1.6		59.5	81.9	69.3	
AD-11795	-1.9	-0.8	0.1		78.7	94.4	87.2	
AD-11796	5.0	1.8	0.8		89.6	71.8	71.8	
AD-11797	0.6	0.8	1.3		92.1	87.0	83.7	
AD-11798	-0.2	-0.2	-1.8		67.4	69.7	45.4	
AD-11799	-2.1	-1.9	-1.3		-49.1	-5.6	72.9	
AD-11800	1.3	-1.0	-0.4		64.1	58.5	60.6	
AD-11801	0.1	0.1	0.8		23.7	35.3	54.5	
AD-11802	4.8	0.8	-0.4		-115.9	-81.0	-113.4	
AD-11803	3.7	1.3	-2.1		10.5	22.6	16.3	
AD-11804	-2.0	-1.1	0.0		16.0	19.8	36.1	
AD-11805	-0.5	0.2	-0.4		-6.9	6.5	51.6	
AD-11806	-11.1	-16.4	-13.0		79.5	70.3	29.3	
AD-11807	13.3	-13.2	-11.2	-78.63	65.7	20.0	50.5	-58.37
AD-11808	-6.3	-9.0	9.4		27.5	30.6	30.6	
AD-11809	-0.2	-11.7	-11.7		44.7	61.3	73.0	
AD-11810	-15.2	-17.1	-17.8		80.1	81.6	81.7	
AD-11811	-9.7	-12.1	-11.2		40.4	38.0	23.7	
AD-11812	-8.1	-12.5	-15.0		21.4	12.4	22.7	
AD-11813	29.3	-14.5	-6.8	-258.78	31.4	32.0	43.9	-423.13
AD-11814	-19.9	-18.8	-8.1		49.2	15.1	55.1	
AD-11815	-12.8	-13.4	-14.9		62.5	29.3	36.0	
AD-11816	-14.5	-15.2	-14.6		13.0	-11.9	46.2	
AD-11817	-15.4	-15.2	-11.8		43.3	38.2	42.0	
AD-11818	9.5	-1.9	-6.6	-176.59	69.0	43.6	35.2	39.23

TABLE 3-continued

duplex name	Ebola Zaire GFP assay % reduction			Ebola Zaire plaque % reduction vs no siRNA	Ebola Sudan IF assay % reduction			Ebola Sudan plaque % reduction vs no siRNA
	100 nM	10 nM	1 nM		100 nM	10 nM	1 nM	
AD-11819	-3.5	-1.4	-1.9	-327.48	75.0	50.7	31.2	29.19
AD-11820	7.6	4.4	-2.0		58.5	50.6	50.6	
AD-11321	1.3	-0.5	1.6		42.4	26.4	49.4	
AD-11822	7.7	-4.1	-4.2		82.6	70.2	71.9	
AD-11823	-1.5	-2.3	-0.4		63.9	58.8	37.4	
AD-11824	3.2	1.7	-2.7		33.5	13.3	37.3	
AD-11825	0.1	-0.9	0.7		53.6	42.2	44.8	
AD-11826	3.4	-6.1	-3.8		52.8	35.7	39.9	
AD-11827	-4.2	-2.6	-1.1		54.5	26.3	15.2	
AD-11828	0.8	-1.3	-1.4		9.3	-4.3	19.5	
AD-11829	2.0	-1.6	1.1	-260.69	32.2	16.0	32.0	25.84
AD-11830	-4.3	-7.5	-8.0		44.5	48.3	75.6	
AD-11831	-6.4	-5.2	-4.1		39.3	56.5	67.3	
AD-11832	-3.2	-2.5	-3.0		61.0	76.2	76.2	
AD-11833	-1.4	-2.5	-1.2		80.6	87.2	91.0	
AD-11834	18.8	2.8	-7.7		7.3	-11.5	24.5	
AD-11835	-2.0	-0.2	-4.6		52.7	59.4	61.0	
AD-11836	3.3	0.7	-1.9		70.7	76.0	80.0	
AD-11837	0.3	-2.3	-0.7		84.6	70.1	80.8	
AD-11838	-7.4	-7.1	-8.8		-13.5	9.5	8.9	
AD-11839	-5.6	-2.9	-2.2	35.57	-0.7	45.3	48.4	42.11
AD-11840	-1.7	1.3	-0.2		85.7	61.6	59.6	
AD-11841	-3.8	-1.8	-2.3		60.7	52.5	50.8	
AD-11842	14.4	7.7	0.8		-145.5	-117.2	-46.4	
AD-11843	ND	ND	ND		-68.1	-87.6	-36.1	
AD-11844	0.0	1.4	-0.5		-72.1	-80.2	-80.2	
AD-11845	3.9	2.0	1.1		-34.1	-60.1	-13.9	
AD-11846	27.5	20.0	-7.1		-69.7	-62.0	-63.7	
AD-11847	-5.9	-4.8	-6.2		-93.2	-61.3	-57.4	
AD-11848	-1.3	0.6	-1.7		-89.3	-76.1	-18.3	
AD-11849	-1.0	-1.4	50.6	25.19	-0.9	12.6	18.7	13.40
AD-11850	-4.1	2.9	-6.3		-15.1	-3.7	-34.5	
AD-11851	-5.0	-4.6	-2.2		-13.3	-10.3	-15.2	
AD-11852	-0.4	-0.6	-0.6		7.7	11.6	10.2	
AD-11853	1.6	-0.6	-0.8		24.9	28.1	36.7	
AD-11854	-0.6	-1.3	-1.8		50.9	35.6	21.2	
AD-11855	5.0	3.4	1.6		47.8	24.7	28.6	
AD-11856	2.3	0.9	-0.6		60.4	36.7	36.7	
AD-11857	3.4	0.7	0.1		23.1	37.0	57.6	
AD-11858	-2.3	-2.5	-2.7		59.8	78.3	41.7	
AD-11859	0.6	2.0	-1.0	32.44	64.0	33.3	40.7	54.70
AD-11860	11.4	6.4	0.1		40.2	-5.8	34.3	
AD-11861	3.2	1.7	-0.4		-6.3	-30.9	26.8	
AD-11862	15.3	3.2	-3.1		37.3	6.8	51.0	
AD-11863	8.7	8.8	4.1		26.0	17.7	37.8	
AD-11864	8.9	3.5	-1.4		40.7	27.9	-8.9	
AD-11865	4.3	0.2	0.1		0.3	8.6	14.9	
AD-11870	4.5	5.7	1.9		60.1	43.2	68.8	
AD-11871	13.9	4.6	-0.4		59.8	62.3	72.1	
AD-11872	12.4	9.9	-0.2		75.1	77.9	77.9	
AD-11873	16.2	8.7	2.9	-55.53	81.9	90.3	92.0	79.43
AD-11874	14.0	3.6	-0.6		63.74	41.1	21.2	
AD-11875	37.1	22.5	9.9		92.11	54.4	47.5	
AD-11876	11.8	4.2	1.3		5.85	35.7	43.8	
AD-11878	3.5	-0.5	-1.7		65.6	62.4	60.0	
AD-11879	30.5	21.5	3.8		93.89	-4.3	11.4	
AD-11882	16.0	12.4	5.1		80.92	9.1	4.8	
AD-11883	13.2	6.8	1.4		-37.40	-29.3	12.5	
AD-11884	23.7	11.4	5.5		25.83	41.4	25.2	
AD-11885	1.5	-0.8	-2.9		55.9	53.3	53.8	
AD-11886	-1.2	-3.4	-4.7	-85.11	38.7	47.2	67.9	85.65
AD-11887	16.6	4.6	-2.2		69.8	54.1	54.1	
AD-11888	12.9	3.8	2.3		62.2	66.5	65.7	
AD-11889	12.2	5.9	-1.6		4.6	10.5	40.1	
AD-11890	8.5	6.2	5.0		-41.7	-12.9	20.9	
AD-11891	20.7	9.8	1.0		19.3	23.9	32.0	
AD-11892	-0.1	-2.5	-2.0		38.4	32.1	38.0	
AD-11893	0.1	-2.1	-1.5		27.1	21.1	3.1	
AD-11896	3.5	1.3	-1.6		23.3	-6.7	10.0	
AD-11897	1.6	-1.3	-2.6		26.4	13.7	13.1	
AD-11899	8.3	1.9	0.6	-69.47	16.4	20.9	28.9	-1048.33
AD-11901	2.5	3.1	1.4		-89.3	71.9	78.6	
AD-11902	5.9	0.6	-2.6		-15.2	62.9	79.8	

TABLE 3-continued

duplex name	Ebola Zaire GFP assay % reduction			Ebola Zaire plaque % reduction vs no siRNA	Ebola Sudan IF assay % reduction			Ebola Sudan plaque % reduction vs no siRNA
	100 nM	10 nM	1 nM		100 nM	10 nM	1 nM	
AD-11903	1.1	-1.5	-3.6		64.0	45.9	45.9	
AD-11904	0.7	0.4	-0.2		46.4	16.9	70.6	
AD-11905	6.9	2.1	-0.1		-42.4	42.5	27.9	
AD-11906	-0.4	0.3	-2.5		98.5	93.5	54.3	
AD-11907	4.5	0.5	-1.1		95.8	45.3	95.1	
AD-11908	-0.9	-1.7	0.2		39.0	86.6	84.5	
AD-11909	7.0	2.9	1.1		79.1	-27.1	83.9	
AD-11911	5.4	2.6	-1.9		97.4	98.6	94.0	
AD-11912	-3.4	-4.5	-3.7		99.4	96.2	65.9	
AD-11914	-0.6	-0.2	1.0		91.7	69.5	91.5	
AD-11918	-6.3	-8.2	-6.7		-12.4	9.7	8.0	
AD-11919	-1.0	-0.1	3.2		18.2	-9.2	32.5	
AD-11925	3.5	1.9	-2.7		11.9	13.7	13.7	
AD-11926	11.3	4.9	-3.2	46.95	16.9	37.0	66.5	93.16
AD-11927	-6.8	-8.4	-5.7		45.5	-12.2	-17.2	
AD-11933	-0.2	1.4	3.3		-6.7	32.7	3.5	
AD-11938	0.0	-2.4	-4.1		19.2	-45.8	1.4	
AD-11939	-2.3	-2.0	-3.1		3.0	23.1	24.9	
AD-11941	-8.3	-9.6	-7.5		42.1	4.0	-25.0	
AD-11942	-4.4	-3.5	-2.5		31.6	2.5	1.4	
AD-11943	1.9	0.7	-2.7		-5.6	20.5	15.4	
AD-11944	-1.1	-2.1	-2.0		51.0	46.7	50.6	
AD-11945	-7.7	-8.6	-5.9		77.2	30.5	55.8	
AD-11946	0.6	0.8	-2.4		66.9	65.2	65.4	
AD-11947	-0.6	-2.2	-4.1		66.6	68.5	68.5	
AD-11948	-2.3	-4.0	-4.5		77.1	77.6	88.6	
AD-11949	-6.6	-7.9	-6.4		67.5	33.7	2.9	
AD-11950	-3.9	0.8	7.9		56.0	51.7	44.8	
AD-11951	7.7	0.4	6.0		35.6	45.6	46.1	
AD-11952	1.7	-4.5	-4.7		54.8	65.7	66.0	
AD-11953	-9.1	-7.7	-6.8		51.4	22.8	30.1	
AD-11954	-2.7	8.2	8.2		42.1	26.4	32.1	
AD-11955	9.0	7.5	5.9		54.6	43.4	44.2	
AD-11956	6.3	-6.5	-4.0		28.1	40.0	55.3	
AD-11957	-1.9	-6.4	-2.8		78.2	95.9	69.2	
AD-11958	2.0	1.2	-1.0		99.7	89.5	86.2	
AD-11959	8.7	3.4	-1.2		81.2	74.5	74.5	
AD-11960	2.8	1.7	0.2		71.9	61.4	82.7	
AD-11961	-3.4	-5.5	-4.5		84.3	50.6	68.7	
AD-11962	-1.4	0.0	-0.2		74.1	89.2	63.9	
AD-11963	2.1	-0.3	-0.2		78.8	20.1	50.9	
AD-11964	4.5	3.9	-0.1		61.5	51.0	58.4	
AD-11965	-3.8	-6.3	-4.4		64.6	14.2	89.4	
AD-11966	-0.5	-1.0	-0.5		49.5	-30.0	21.4	
AD-11967	5.0	3.8	1.2		24.7	28.2	24.2	
AD-11968	-0.8	-0.2	-0.3		9.5	6.5	26.5	
AD-11969	3.9	0.6	-2.2		63.5	31.6	56.5	
AD-11970	9.2	5.5	-1.6		47.6	36.9	57.1	
AD-11971	-1.2	0.1	-1.7		63.8	55.3	55.3	
AD-11972	2.7	-0.9	-0.8		66.7	82.1	78.1	
AD-11973	3.2	5.3	-0.3		-137.3	-57.5	-69.7	
AD-11974	1.7	2.3	2.6		-15.1	-41.3	-55.7	
AD-11975	1.8	1.4	1.9		-63.6	-63.1	-54.6	
AD-11976	2.3	2.7	2.7		-8.3	10.1	29.0	
AD-11977	4.5	3.7	2.7		-168.8	-142.8	-116.4	
AD-11978	2.1	1.2	0.8		-66.5	-91.0	-120.2	
AD-11979	-1.7	-2.0	-2.2		-140.3	-128.4	-107.4	
AD-11980	-1.3	-0.8	-0.6		-21.5	-10.2	-14.2	
AD-11981	9.1	6.7	4.6		-181.1	-162.1	-164.3	
AD-11982	-0.2	-0.4	-1.4		-189.5	-148.6	-159.6	
AD-11983	-2.7	-3.4	-3.7		-131.3	-111.7	-115.3	
AD-11984	-1.9	-0.5	0.1		-36.1	-12.9	-10.8	
AD-11985	5.5	6.4	2.8		-139.4	-160.3	-216.6	
AD-11986	6.1	6.0	5.8		-111.7	-143.8	-134.7	
AD-11987	0.9	1.5	2.0		-162.1	-148.5	-132.2	
AD-11988	7.4	8.5	8.9		-196.2	-201.7	-178.2	
AD-11989	1.9	2.0	1.6		-62.4	-57.6	-61.2	
AD-11990	0.8	-0.2	-0.8		-51.3	-69.3	-76.8	
AD-11991	-1.9	-2.4	-2.3		-84.5	-65.2	-51.0	
AD-11992	0.6	1.3	1.6		-83.1	-60.4	-57.1	
AD-11993	6.6	5.6	3.7		-29.9	-33.8	-35.4	
AD-11994	0.3	0.1	-0.6		-60.1	-59.5	-59.4	
AD-11995	-2.3	-2.1	-1.9		-73.2	-61.7	-58.5	

TABLE 3-continued

duplex name	Ebola Zaire GFP assay % reduction			Ebola Zaire plaque % reduction vs no siRNA	Ebola Sudan IF assay % reduction			Ebola Sudan plaque % reduction vs no siRNA
	100 nM	10 nM	1 nM		100 nM	10 nM	1 nM	
AD-11996	-0.7	0.3	0.4		-4.6	-14.0	-21.4	
AD-11997	4.1	4.3	-0.1		64.7	26.4	52.9	
AD-11998	0.3	0.0	-0.7		69.1	61.0	52.1	
AD-11999	-0.5	-0.1	0.1		31.9	37.3	39.5	
AD-12000	-0.8	0.2	0.8		51.7	54.7	60.6	
AD-12001	3.7	3.3	2.9		54.7	49.4	47.3	
AD-12002	3.8	2.9	2.2		34.0	49.3	62.0	
AD-12003	-0.3	-0.4	-0.2		26.3	29.4	32.9	
AD-12004	0.6	0.8	1.4		37.7	36.6	36.9	
AD-12005	4.9	4.7	3.9		19.6	9.3	7.5	
AD-12006	0.8	0.5	0.5		-22.4	-23.6	-19.4	
AD-12007	-1.3	-0.7	-0.3		-10.4	-13.6	-21.7	
AD-12008	0.7	1.4	1.4		31.6	31.4	33.4	
AD-12009	0.8	2.5	-2.2		36.9	25.4	31.0	
AD-12010	-1.2	-1.5	-1.6		35.6	23.9	15.8	
AD-12011	-1.2	-0.3	0.2		50.4	46.0	41.7	
AD-12012	0.7	0.7	0.5		47.1	49.2	51.5	
AD-12013	1.3	1.1	-0.5		31.5	25.2	21.8	
AD-12014	-1.8	-3.0	-2.7		21.5	26.1	26.3	
AD-12015	-4.2	-4.1	-3.8		36.5	35.7	34.2	
AD-12016	-1.9	-1.5	-1.2		38.7	41.0	41.9	
AD-12017	-0.4	-1.0	-2.2		6.4	12.0	17.5	
AD-12018	-1.0	-2.6	-2.9		-19.0	-8.5	-7.1	
AD-12019	-3.1	-2.7	-2.3		-3.4	-3.2	0.0	
AD-12020	-2.1	-1.4	-1.0		2.2	10.2	12.7	
AD-12021	4.9	3.3	2.1		-34.2	-9.2	-30.6	
AD-12022	3.9	3.1	2.1		-139.2	-184.9	-191.4	
AD-12023	0.4	0.5	0.8		10.7	11.7	-2.8	
AD-12024	0.2	1.4	2.3		-8.4	6.0	18.1	
AD-12025	6.9	4.8	3.2		18.0	-1.4	-25.4	
AD-12026	-0.3	-0.1	0.2		-79.8	-64.2	-44.8	
AD-12027	-0.8	-1.0	-0.5		-28.2	-43.7	-57.1	
AD-12028	-1.2	-0.2	0.8		-8.1	-0.6	6.2	
AD-12029	6.9	5.6	3.8		-65.2	-79.3	-106.3	
AD-12030	-0.1	-0.8	-2.0		-70.4	-93.2	-119.7	
AD-12031	0.2	0.1	-0.1		-32.4	-34.4	-34.9	
AD-12032	0.0	0.1	0.6		-86.6	-68.5	-43.4	
AD-12033	-4.7	-8.2	-7.6		-115.7	-149.7	-58.5	
AD-12034	-6.0	-5.7	-5.7		-217.6	-210.8	-215.6	
AD-12035	-7.7	-7.4	-7.6		-99.6	-102.4	-107.1	
AD-12036	-6.6	-6.1	-5.1		-111.1	-91.0	-89.8	
AD-12037	-7.8	-7.8	-7.7		45.1	-5.3	-44.2	
AD-12462	-8.4	-8.6	-9.4		-150.2	-135.8	-130.8	
AD-12463	-8.3	-8.4	-8.6		-70.1	-70.9	-75.8	
AD-12464	-6.3	-6.6	-6.5		-38.9	-19.7	-2.6	
AD-12465	-2.8	-3.8	-5.3		-52.8	-75.2	-92.9	
AD-12466	-1.9	-3.1	-3.5		-68.5	-103.9	-113.9	
AD-12467	-5.9	-5.8	-5.6		-118.6	-121.8	-117.7	
AD-12468	-4.9	-4.2	-3.3		-14.2	-8.4	-6.8	
AD-12469	-4.8	-6.0	-6.6		-29.1	-6.1	-10.2	
AD-12470	-0.9	-1.4	-1.9		-39.5	-19.2	-1.8	
AD-12471	-4.0	-4.0	-4.0		-15.4	-16.3	-17.3	
AD-12472	-3.2	-3.4	-2.6		25.9	24.0	27.4	
AD-12473	-3.5	-3.8	-4.1		-42.5	-31.5	-26.4	
AD-12474	-5.0	-5.3	-5.8		-7.1	-3.0	-1.8	
AD-12475	-7.1	-7.6	-7.4		-10.2	-4.3	-0.7	
AD-12476	-5.2	-4.7	-3.5		24.3	26.9	32.2	
AD-12477	-2.8	-3.8	-4.4		-41.4	-45.4	-37.6	
AD-12478	-7.3	-7.6	-7.8		-59.7	-63.8	-67.5	
AD-12479	-7.9	-8.3	-8.1		-54.9	-39.3	-8.6	
AD-12480	-2.7	-2.4	-1.4		9.3	18.0	12.4	
AD-12481	-5.7	-5.5	-7.5		-131.4	-152.8	-48.2	
AD-12482	-4.4	-4.5	-4.1		-159.1	-127.8	-140.7	
AD-12483	-4.8	-4.8	-4.5		-66.0	-64.8	-60.3	
AD-12484	-4.2	-3.9	-3.8		17.9	27.6	29.1	
AD-12485	-2.7	-4.0	-5.4		-170.2	-176.6	-186.0	
AD-12486	-8.5	-8.6	-8.6		-200.1	-194.6	-169.0	
AD-12487	-8.1	-8.0	-7.8		-117.5	-106.8	-86.5	
AD-12488	-8.2	-7.9	-7.8		-17.3	-13.1	0.8	
AD-12489	-4.2	-5.6	-7.1		-128.0	-152.3	-185.8	
AD-12490	-8.6	-8.9	-8.8		-241.7	-258.1	-251.4	
AD-12491	-8.9	-9.0	-8.4		-181.6	-184.9	-188.0	
AD-12492	-8.6	-7.7	-6.7		-23.3	-22.4	-27.3	

TABLE 3-continued

duplex name	Ebola Zaire GFP assay % reduction			Ebola Zaire plaque % reduction vs no siRNA	Ebola Sudan IF assay % reduction			Ebola Sudan plaque % reduction vs no siRNA
	100 nM	10 nM	1 nM		100 nM	10 nM	1 nM	
AD-12493	-8.2	-6.7	-4.5		-93.8	-54.8	6.7	
AD-12494	-7.6	-8.0	-8.2		-62.2	-63.5	-54.7	
AD-12495	-6.6	-5.5	-4.9		16.6	0.4	-10.0	
AD-12496	-4.1	-4.2	-3.8		40.2	41.7	39.5	
AD-12497	-5.1	-5.2	-6.3		-67.9	-65.2	-67.3	
AD-12498	-5.8	-6.0	-6.4		-68.6	-84.1	-94.1	
AD-12499	-6.6	-6.3	-6.3		-93.4	-80.2	-61.0	
AD-12500	-6.0	-5.4	-4.7		9.7	10.2	11.3	
AD-12501	-4.8	-5.0	-5.8		-87.1	-108.5	-110.6	
AD-12502	-6.8	-7.2	-6.7		-126.5	-125.5	-119.1	
AD-12503	-10.4	-9.6	-9.9		-85.2	-76.4	-86.7	
AD-12504	-4.0	-3.6	-2.5		-59.7	-52.7	-34.7	
AD-12505	18.1	-0.9	-5.5	42.18	3.5	-116.5	-34.2	-282.78
AD-12506	-2.1	-2.7	-4.2		-29.8	-22.3	-11.9	
AD-12507	-1.7	7.2	15.0	44.47	29.1	33.0	32.3	
AD-12508	-4.2	-3.2	-1.9		40.7	33.9	30.5	
AD-12509	-9.2	-8.9	-8.2		-54.3	-67.7	-80.3	
AD-12510	-7.9	-7.6	-7.6		-127.6	-117.5	-124.2	
AD-12511	-7.1	-6.9	-6.8		-28.4	-18.6	-6.7	
AD-12512	-5.2	-4.1	-3.3		-0.4	-1.7	-1.5	
AD-12513	-6.9	-7.4	-8.8		-65.5	-80.0	-100.6	
AD-12514	-10.8	-11.0	-10.6		-83.8	-85.0	-92.0	
AD-12515	-10.7	-9.6	-9.3		-19.3	-41.2	-32.7	
AD-12516	-5.8	-5.3	-4.2		-5.9	0.9	15.9	
AD-12517	-8.6	-8.7	-7.1		29.2	53.0	49.4	
AD-12518	-8.3	-8.0	-8.0		65.7	57.8	53.6	
AD-12519	-7.2	-7.2	-7.6		51.2	51.3	53.9	
AD-12520	-7.0	-6.3	-5.4		67.7	73.2	78.8	
AD-12521	-6.4	-8.0	-9.1		19.8	27.6	35.5	
AD-12522	-9.9	-9.0	-9.1		50.3	43.0	34.3	
AD-12523	-9.0	-8.7	-8.9		28.3	31.9	38.1	
AD-12524	-7.0	-7.4	-7.0		63.7	67.1	65.9	
AD-12525	-1.6	-3.8	-6.0		16.1	21.8	21.2	

TABLE 4

In vitro Plaque assay controls (all values average of 3-4 experiments and expressed as % inhibition relative to no siRNA treatment)			
AD-1955 (luc)	31.37	10.97	
AD-5179 (GFP)	15.15	18.65	
LS L#1	77.23	N/A	

TABLE 4-continued

In vitro Plaque assay controls (all values average of 3-4 experiments and expressed as % inhibition relative to no siRNA treatment)			
LS NP#1	73.94	N/A	
LS VP35#1	67.27	N/A	

TABLE 5

parent duplex	Exo + endo light duplex	Target	sense	anti-sense	seq id no	sense 5'-3'	seq id no	antisense 5'-3'
AD-11594	AD-3542	NP	A-30768	A-30769	995	cAGuAGGAcAcAuGAuGGudTsdT	996	ACcAUcAUGUGUCCuACUGdTsdT
AD-11596	AD-3543	NP	A-30770	A-30771	997	GuAGGAcAcAuGAuGGuGAdTsdT	998	UcACcAUcAUGUGUCCuACdTsdT
AD-11597	AD-3544	NP	A-30772	A-30773	999	uAGGAcAcAuGAuGGuGAudTsdT	1000	AUcACcAUcAUGUGUCCuAdTsdT
AD-11598	AD-3545	NP	A-30774	A-30775	1001	GAuGGuGAuuuuccGuuuGdTsdT	1002	cAAACGGAAAAUcACcAUcCdTsdT
AD-11600	AD-3546	NP	A-30776	A-30777	1003	uGGuGAuuuuccGuuuGAudTsdT	1004	AUcAAACGGAAAAUcACcAdTsdT
AD-11603	AD-3547	NP	A-30778	A-30779	1005	cuGAGAAGcAAcuccAAcAdTsdT	1006	UGUUGGAGUUGCUUCUcAgdTsdT

TABLE 5-continued

parent duplex	Exo + endo light duplex	Target	sense	anti-sense	seq id no	sense 5'-3'	seq id no	antisense 5'-3'
AD-11604	AD-3548	NP	A-30780	A-30781	1007	uGAGAAGcAAcuccAAcAAAdTsdT	1008	UUGUUGAGUUGCUUCUcAdTsdT
AD-11606	AD-3549	NP	A-30782	A-30783	1009	AGAAGcAAcuccAAcAAuAdTsdT	1010	uAUUGUUGAGUUGCUUCUdTsdT
AD-11609	AD-3550	VP35	A-30784	A-30785	1011	AAGuGAuGAAGAuAAGAAAdTsdT	1012	UUCUuAAUCUUCAUcACUUDTsdT
AD-11610	AD-3551	VP35	A-30786	A-30787	1013	AGuGAuGAAGAuAAGAAAdTsdT	1014	UUUCUuAAUCUUCAUcACUdTsdT
AD-11611	AD-3552	VP40	A-30788	A-30789	1015	cuGccuGcuGcAAcAuGGAdTsdT	1016	UCCAUUGUGcAGcAGGcAGdTsdT
AD-11613	AD-3553	GP	A-30790	A-30791	1017	GGcuGAAAACuGcuAcAAudTsdT	1018	AUUGuAGcAGUUUUCAGCCdTsdT
AD-11614	AD-3554	GP	A-30792	A-30793	1051	GcuGAAAACuGcuAcAAucdTsdT	1052	GAUUGuAGcAGUUUUCAGCdTsdT
AD-11615	AD-3555	GP	A-30794	A-30795	1053	cuGAAAACuGcuAcAAucdTsdT	1054	AGAUGuAGcAGUUUUCAGdTsdT
AD-11618	AD-3556	GP	A-30796	A-30797	1055	AAAACuGcuAcAAucuuGAdTsdT	1056	UcAAGAUUGuAGcAGUUUdTsdT
AD-11621	AD-3557	GP	A-30798	A-30799	1057	AcuGcuAcAAucuuGAAudTsdT	1058	AUUUcAAGAUUGuAGcAGUdTsdT
AD-11622	AD-3558	VP30	A-30800	A-30801	1059	AGcAAAuccAAcGGcuGAudTsdT	1060	AUcAGCCGUUGGAUUUGCUdTsdT
AD-11624	AD-3559	VP30	A-30802	A-30803	1061	cAAAuccAAcGGcuGAuGAdTsdT	1062	UcAUcAGCCGUUGGAUUUGdTsdT
AD-11627	AD-3560	L	A-30804	A-30805	1063	AuGcAuGucAGuGAuuAuudTsdT	1064	AAuAAUcACUGAcAUGcAUdTsdT
AD-11628	AD-3561	L	A-30806	A-30807	1065	uGcAuGucAGuGAuuAuAdTsdT	1066	uAAuAAUcACUGAcAUGcAdTsdT
AD-11629	AD-3562	L	A-30808	A-30809	1067	GcAuGucAGuGAuuAuAdTsdT	1068	AuAAuAAUcACUGAcAUGCdTsdT
AD-11630	AD-3563	L	A-30810	A-30811	1069	cAuGucAGuGAuuAuAdTsdT	1070	uAuAAuAAUcACUGAcAUGdTsdT
AD-11631	AD-3564	L	A-30812	A-30813	1071	AuGucAGuGAuuAuAAAdTsdT	1072	UuAuAAuAAUcACUGAcAUdTsdT

TABLE 6

modified duplex	In vitro plasmid screen IC50 (nM)	In vitro plaque assay against Ebola-Zaire (% inhibition relative to no siRNA)	In vitro plaque assay against Ebola-Sudan (% inhibition relative to no siRNA)
AD-3542	1.60	84%	74%
AD-3543	0.00	74%	77%
AD-3544	0.61	65%	13%
AD-3545	0.00	82%	61%
AD-3546	1.35	88%	71%
AD-3547	0.00	81%	26%
AD-3548	7.82	68%	73%
AD-3549	0.00	68%	74%
AD-3550	0.17	77%	22%
AD-3551	0.08	2%	75%

TABLE 6-continued

modified duplex	In vitro plasmid screen IC50 (nM)	In vitro plaque assay against Ebola-Zaire (% inhibition relative to no siRNA)	In vitro plaque assay against Ebola-Sudan (% inhibition relative to no siRNA)
AD-3552	0.00	78%	70%
AD-3553	0.19	85%	26%
AD-3554	0.77	84%	15%
AD-3555	0.88	-5%	19%
AD-3556	0.00	73%	74%
AD-3557	0.21	100%	28%
AD-3558	0.00	-7%	68%
AD-3559	0.00	-252%	-7%
AD-3560	0.50	16%	-35%
AD-3561	0.00	-105%	19%

TABLE 6-continued

modified duplex	In vitro plasmid screen IC50 (nM)	In vitro plaque assay against Ebola-Zaire (% inhibition relative to no siRNA)	In vitro plaque assay against Ebola-Sudan (% inhibition relative to no siRNA)
AD-3562	0.00	-75%	63%
AD-3563	5.42	73%	-17%
AD-3564	0.97	-86%	19%
AD-3621	0.00	80%	48%
AD-3622	16.00	61%	47%
AD-3623	0.00	98%	37%
AD-3624	0.00	84%	24%
AD-3625	0.00	-7%	-5%
AD-3626	0.00	53%	31%

TABLE 7

IFN induction normalized to positive control siRNA		TNF induction normalized to positive control siRNA	
duplex	% of control	duplex	% of control
AD-11546	131.6	AD-11546	30.1
AD-11558	98.8	AD-11558	57.2
AD-11570	0	AD-11570	0
AD-11588	147	AD-11588	0.0
AD-11590	0.0	AD-11590	130.5
AD-11594	17.8	AD-11594	124.8
AD-11597	22.0	AD-11597	242.9
AD-11598	25.7	AD-11598	180.3
AD-11599	136	AD-11599	183.9
AD-11600	13.5	AD-11600	141.0
AD-11603	69.0	AD-11603	81.4
AD-11606	33.2	AD-11606	79.8
AD-11609	126.9	AD-11609	46.3
AD-11610	138.2	AD-11610	48.5
AD-11611	43.2	AD-11611	41.0
AD-11613	167.0	AD-11613	52.6
AD-11614	162.1	AD-11614	48.1
AD-11615	171.2	AD-11615	60.0
AD-11618	137.3	AD-11618	54.6
AD-11621	0.0	AD-11621	325.8
AD-11622	37.2	AD-11622	28.5
AD-11623	58.0	AD-11623	26.0
AD-11624	63.2	AD-11624	29.2
AD-11627	6.4	AD-11627	125.0
AD-11628	0.0	AD-11628	101.1
AD-11630	0.0	AD-11630	170.3
AD-11631	0.0	AD-11631	156.0
AD-11644	0	AD-11644	0.0
AD-11650	0.0	AD-11650	0.0
AD-11659	0.0	AD-11659	69.0
AD-11673	9.2	AD-11673	0.0
AD-11678	0.0	AD-11678	0.0
AD-11683	0.0	AD-11683	19.6
AD-11684	0.0	AD-11684	24.3
AD-11691	0	AD-11691	0.0
AD-11695	0.0	AD-11695	5.2
AD-11698	5.3	AD-11698	14.0
AD-11706	0.0	AD-11706	0.0
AD-11707	0	AD-11707	0.0
AD-11710	0	AD-11710	0.0
AD-11721	0.0	AD-11721	0.0
AD-11725	0.0	AD-11725	0.0
AD-11732	0.0	AD-11732	0.0
AD-11743	16.1	AD-11743	0.0
AD-11756	0.0	AD-11756	0.0
AD-11757	0.0	AD-11757	0.0
AD-11758	0.0	AD-11758	0.0
AD-11759	0.0	AD-11759	0.0
AD-11773	0.0	AD-11773	0.0
AD-11780	0.0	AD-11780	7.9
AD-11789	0.0	AD-11789	5.7
AD-11804	0.0	AD-11804	0.0

TABLE 7-continued

IFN induction normalized to positive control siRNA		TNF induction normalized to positive control siRNA	
duplex	% of control	duplex	% of control
AD-11811	0.0	AD-11811	0.0
AD-11814	0.0	AD-11814	0.0
AD-11816	0.0	AD-11816	0.0
AD-11822	9.8	AD-11822	0.0
AD-11823	0.0	AD-11823	0.0
AD-11832	0.0	AD-11832	0.0
AD-11836	0.0	AD-11836	0.0
AD-11939	0.0	AD-11939	0.0
AD-11976	0.0	AD-11976	0.0
AD-11982	6.9	AD-11982	15.1
AD-11990	12.2	AD-11990	0.0
AD-11992	7.3	AD-11992	0.0
AD-12007	0.0	AD-12007	0.0
AD-12013	0.0	AD-12013	0.0
AD-12019	24.6	AD-12019	5.0
AD-12024	0.0	AD-12024	19.8
AD-12035	0.0	AD-12035	8.5
AD-12475	0.0	AD-12475	0.0
AD-12484	21.0	AD-12484	9.5
AD-12491	0.0	AD-12491	13.1
AD-12500	12.6	AD-12500	53.3
AD-12502	101.6	AD-12502	55.5
AD-3542	0.0	AD-3542	0.0
AD-3543	0.0	AD-3543	0.0
AD-3544	10.1	AD-3544	0.0
AD-3545	11.8	AD-3545	0.0
AD-3546	0.0	AD-3546	0.0
AD-3547	0.0	AD-3547	31.0
AD-3548	0.0	AD-3548	5.9
AD-3549	0.0	AD-3549	10.8
AD-3550	7.4	AD-3550	0.0
AD-3551	0.0	AD-3551	0.0
AD-3552	0.0	AD-3552	0.0
AD-3553	0.0	AD-3553	0.0
AD-3554	0.0	AD-3554	11.5
AD-3555	0.0	AD-3555	6.2
AD-3556	0.0	AD-3556	9.1
AD-3557	0.0	AD-3557	0.0
AD-3558	0.0	AD-3558	5.8
AD-3559	0.0	AD-3559	5.3
AD-3560	0.0	AD-3560	0.0
AD-3561	0.0	AD-3561	0.0
AD-3562	0.0	AD-3562	0.0
AD-3563	0.0	AD-3563	0.0
AD-3564	0.0	AD-3564	0.0
AD-3621	0.0	AD-3621	0.0
AD-3622	0.0	AD-3622	0.0
AD-3623	5.7	AD-3623	0.0
AD-3624	10.0	AD-3624	0.0
AD-3625	0.0	AD-3625	0.0
AD-3626	0.0	AD-3626	0.0

TABLE 8

duplex name	In vitro plasmid screen single dose (% silencing)	In vitro plasmid screen IC50 (nM)
AD-11542	9%	
AD-11543	-3%	
AD-11544	-1%	
AD-11545	13%	
AD-11546	68%	0.72
AD-11547	28%	
AD-11548	48%	
AD-11549	43%	
AD-11550	-5%	
AD-11551	8%	
AD-11552	-17%	
AD-11553	-6%	

TABLE 8-continued

duplex name	In vitro plasmid screen single dose (% silencing)	In vitro plasmid screen IC50 (nM)
AD-11554	15%	0.46
AD-11555	2%	
AD-11556	6%	
AD-11557	-6%	
AD-11558	70%	
AD-11559	28%	
AD-11560	5%	
AD-11561	30%	
AD-11562	24%	
AD-11563	6%	
AD-11564	0%	0.95
AD-11565	-4%	
AD-11566	4%	
AD-11567	-2%	
AD-11568	0%	
AD-11569	12%	
AD-11570	73%	
AD-11571	-2%	
AD-11572	-3%	
AD-11573	2%	
AD-11574	15%	0.57
AD-11575	-2%	
AD-11576	-9%	
AD-11577	-9%	
AD-11578	77%	
AD-11579	37%	
AD-11580	35%	
AD-11581	60%	
AD-11582	21%	
AD-11583	47%	
AD-11584	0%	0.35
AD-11585	-1%	
AD-11586	36%	
AD-11587	66%	
AD-11588	47%	
AD-11589	77%	
AD-11590	83%	
AD-11591	65%	
AD-11592	62%	
AD-11593	55%	
AD-11594	85%	0.24
AD-11595	72%	
AD-11596	84%	
AD-11597	85%	
AD-11598	87%	
AD-11599	91%	
AD-11600	89%	
AD-11601	84%	
AD-11602	71%	
AD-11603	80%	
AD-11604	81%	6.38
AD-11605	75%	
AD-11606	78%	
AD-11607	53%	
AD-11608	60%	
AD-11609	75%	
AD-11610	74%	
AD-11611	61%	
AD-11612	-5%	
AD-11613	84%	
AD-11614	85%	0.24
AD-11615	79%	
AD-11616	66%	
AD-11617	59%	
AD-11618	78%	
AD-11619	57%	
AD-11620	64%	
AD-11621	74%	
AD-11622	70%	
AD-11623	67%	
AD-11624	75%	0.15
AD-11625	11%	
AD-11626	51%	
AD-11627	71%	

TABLE 8-continued

duplex name	In vitro plasmid screen single dose (% silencing)	In vitro plasmid screen IC50 (nM)
AD-11628	68%	0.33
AD-11629	75%	0.18
AD-11630	73%	0.24
AD-11631	69%	0.31
AD-11632	53%	1.78
AD-11633	63%	
AD-11634	65%	
AD-11635	29%	
AD-11636	43%	
AD-11637	-5%	
AD-11638	6%	
AD-11639	2%	
AD-11640	38%	
AD-11641	35%	0.7
AD-11642	55%	
AD-11643	33%	
AD-11644	36%	
AD-11645	45%	
AD-11646	37%	
AD-11647	41%	
AD-11648	61%	
AD-11649	35%	
AD-11650	84%	0.038
AD-11651	13%	
AD-11652	64%	
AD-11653	61%	
AD-11654	6%	
AD-11655	59%	
AD-11656	38%	
AD-11657	39%	
AD-11658	59%	
AD-11659	82%	
AD-11660	39%	0.117
AD-11661	-5%	
AD-11662	-1%	
AD-11663	14%	
AD-11664	19%	
AD-11665	7%	
AD-11666	-4%	
AD-11667	-14%	
AD-11668	63%	
AD-11669	28%	
AD-11670	23%	0.149
AD-11671	23%	
AD-11672	15%	
AD-11673	79%	
AD-11674	67%	
AD-11675	46%	
AD-11676	20%	
AD-11677	34%	
AD-11678	79%	
AD-11679	51%	
AD-11680	24%	0.056
AD-11681	72%	
AD-11682	73%	
AD-11683	88%	
AD-11684	80%	
AD-11685	33%	
AD-11686	72%	
AD-11687	32%	
AD-11688	15%	
AD-11689	58%	
AD-11690	26%	0.46
AD-11691	60%	
AD-11694	54%	
AD-11695	81%	
AD-11696	32%	
AD-11698	73%	
AD-11700	-9%	
AD-11704	35%	
AD-11705	39%	
AD-11706	67%	
AD-11707	2%	0.56
AD-11708	-10%	

TABLE 8-continued

duplex name	In vitro plasmid screen single dose (% silencing)	In vitro plasmid screen IC50 (nM)
AD-11710	65%	4.57
AD-11711	-3%	
AD-11712	17%	
AD-11713	0%	
AD-11714	42%	
AD-11715	41%	
AD-11716	18%	
AD-11717	32%	
AD-11718	-3%	
AD-11719	36%	
AD-11720	41%	
AD-11721	68%	1.35
AD-11722	31%	
AD-11723	49%	
AD-11724	27%	
AD-11725	67%	0.34
AD-11726	12%	
AD-11727	3%	
AD-11728	5%	
AD-11729	12%	
AD-11730	6%	
AD-11731	63%	49.4
AD-11732	76%	2.88
AD-11733	60%	8.76
AD-11734	44%	
AD-11735	17%	
AD-11736	44%	
AD-11737	14%	
AD-11738	-9%	
AD-11739	23%	
AD-11740	1%	
AD-11741	9%	
AD-11742	40%	
AD-11743	77%	0.11
AD-11744	24%	
AD-11745	27%	
AD-11746	-9%	
AD-11747	16%	
AD-11748	8%	
AD-11749		
AD-11750	33%	
AD-11751	19%	
AD-11752	61%	6.92
AD-11753	13%	
AD-11754	53%	
AD-11755		
AD-11756	61%	1.21
AD-11757	63%	1.57
AD-11758	28%	
AD-11759	66%	1.61
AD-11760	64%	3.4
AD-11761	59%	
AD-11762	54%	
AD-11763	47%	
AD-11764	21%	
AD-11765	-1%	
AD-11766	-1%	
AD-11767	67%	4.4
AD-11768	52%	
AD-11769	21%	
AD-11770	55%	
AD-11771	36%	
AD-11772	41%	
AD-11773	76%	0.37
AD-11774	35%	
AD-11775	49%	
AD-11776	50%	
AD-11777	-5%	
AD-11778	18%	
AD-11779	15%	
AD-11780	62%	0.76
AD-11781	14%	
AD-11782	38%	
AD-11783	46%	

TABLE 8-continued

duplex name	In vitro plasmid screen single dose (% silencing)	In vitro plasmid screen IC50 (nM)
AD-11784	23%	
AD-11785	-11%	
AD-11786	42%	
AD-11787	48%	
AD-11788	19%	
AD-11789	64%	0.38
AD-11790	26%	
AD-11791	22%	
AD-11792	-8%	
AD-11793	26%	
AD-11794	57%	
AD-11795		
AD-11796	59%	
AD-11797	11%	
AD-11798	11%	
AD-11799	35%	
AD-11800	2%	
AD-11801	-6%	
AD-11802	0%	
AD-11803	5%	
AD-11804	88%	0.281
AD-11805	5%	
AD-11806	9%	
AD-11807	6%	
AD-11808	50%	
AD-11809	24%	
AD-11810	-1%	
AD-11811	66%	1.56
AD-11812	1%	
AD-11813	17%	
AD-11814	65%	0.43
AD-11815	-1%	
AD-11816	65%	0.99
AD-11817	67%	2.98
AD-11818	44%	
AD-11819	64%	
AD-11820	36%	
AD-11821	64%	
AD-11822	62%	1.44
AD-11823	69%	0.32
AD-11824	38%	
AD-11825	18%	
AD-11826	23%	
AD-11827	2%	
AD-11828	51%	
AD-11829		
AD-11830	46%	
AD-11831	20%	
AD-11832	71%	0.94
AD-11833	3%	
AD-11834	23%	
AD-11835	-8%	
AD-11836	65%	1.46
AD-11837	26%	
AD-11838	-16%	
AD-11839	22%	
AD-11840	-5%	
AD-11841	60%	
AD-11342	19%	
AD-11843	57%	
AD-11844	18%	
AD-11845	8%	
AD-11846	48%	
AD-11847	57%	
AD-11848	2%	
AD-11849	66%	0.32
AD-11850	25%	
AD-11851	-12%	
AD-11852	27%	
AD-11853	28%	
AD-11854	55%	
AD-11855	43%	
AD-11856	-20%	
AD-11857	61%	3.28

TABLE 8-continued

duplex name	In vitro plasmid screen single dose (% silencing)	In vitro plasmid screen IC50 (nM)
AD-11858	6%	
AD-11859	4%	
AD-11860	79%	
AD-11861	31%	
AD-11862	48%	
AD-11863	85%	0.39
AD-11864	42%	
AD-11865	37%	
AD-11870	34%	
AD-11871	79%	0.63
AD-11872	70%	
AD-11873	70%	
AD-11874	39%	
AD-11875	39%	
AD-11876	34%	
AD-11878	-1%	
AD-11879	50%	
AD-11882	-6%	
AD-11883	11%	
AD-11884	7%	
AD-11885	-3%	
AD-11886	-5%	
AD-11887	18%	
AD-11888	41%	
AD-11889	1%	
AD-11890	44%	
AD-11891	20%	
AD-11892	37%	
AD-11893	29%	
AD-11896	1%	
AD-11897	41%	
AD-11899	12%	
AD-11901	-2%	
AD-11902	40%	
AD-11903	14%	
AD-11904	1%	
AD-11905	33%	
AD-11906	2%	
AD-11907	9%	
AD-11908	5%	
AD-11909	16%	
AD-11911	37%	
AD-11912	19%	
AD-11914	19%	
AD-11918	1%	
AD-11919	-1%	
AD-11925	-5%	
AD-11926	60%	
AD-11927	-11%	
AD-11933	30%	
AD-11938	6%	
AD-11939	71%	0.48
AD-11941	47%	
AD-11942	33%	
AD-11943	48%	
AD-11944	51%	
AD-11945	69%	
AD-11946	39%	
AD-11947	39%	
AD-11948	83%	0.68
AD-11949	41%	
AD-11950	73%	
AD-11951	55%	
AD-11952	81%	1.42
AD-11953	52%	
AD-11954	55%	
AD-11955	79%	0.63
AD-11956	37%	
AD-11957	39%	
AD-11958	34%	
AD-11959	36%	
AD-11960	19%	
AD-11961	-4%	
AD-11962	-6%	

TABLE 8-continued

duplex name	In vitro plasmid screen single dose (% silencing)	In vitro plasmid screen IC50 (nM)
AD-11963	3%	
AD-11964	13%	
AD-11965	28%	
AD-11966	-4%	
AD-11967	15%	
AD-11968	4%	
AD-11969	0%	
AD-11970	-7%	
AD-11971	3%	
AD-11972	58%	
AD-11973	4%	
AD-11974	38%	
AD-11975	-11%	
AD-11976	63%	0.21
AD-11977	2%	
AD-11978	56%	
AD-11979	5%	
AD-11980	5%	
AD-11981	19%	
AD-11982	65%	0.14
AD-11983	52%	
AD-11984	50%	
AD-11985	50%	
AD-11986	6%	
AD-11987	-7%	
AD-11988	58%	
AD-11989	27%	
AD-11990	72%	0.24
AD-11991	29%	
AD-11992	76%	0.94
AD-11993	46%	
AD-11994	21%	
AD-11995	-3%	
AD-11996	53%	
AD-11997	0%	
AD-11998	3%	
AD-11999	19%	
AD-12000	41%	
AD-12001	3%	
AD-12002	37%	
AD-12003	17%	
AD-12004	5%	
AD-12005	6%	
AD-12006	69%	
AD-12007	81%	0.268
AD-12008	35%	
AD-12009	22%	
AD-12010	34%	
AD-12011	10%	
AD-12012	25%	
AD-12013	75%	0.60
AD-12014	29%	
AD-12015	4%	
AD-12016	2%	
AD-12017	5%	
AD-12018	7%	
AD-12019	79%	0.904
AD-12020	6%	
AD-12021	0%	
AD-12022	2%	
AD-12023	16%	
AD-12024	87%	0.075
AD-12025	4%	
AD-12026	5%	
AD-12027	66%	
AD-12028	24%	
AD-12029	-8%	
AD-12030	46%	
AD-12031	48%	
AD-12032	-10%	
AD-12033	64%	6.74
AD-12034	45%	
AD-12035	70%	0.11
AD-12036	43%	

TABLE 8-continued

duplex name	In vitro plasmid screen single dose (% silencing)	In vitro plasmid screen IC50 (nM)
AD-12037		
AD-12462	35%	
AD-12463	39%	
AD-12464	47%	
AD-12465	34%	
AD-12466	35%	
AD-12467	25%	
AD-12468	-9%	
AD-12469	-3%	
AD-12470	50%	
AD-12471	12%	
AD-12472	9%	
AD-12473	52%	
AD-12474	1%	
AD-12475	62%	0.13
AD-12476	19%	
AD-12477	-12%	
AD-12478	5%	
AD-12479	21%	
AD-12480	13%	
AD-12481	22%	
AD-12482	65%	
AD-12483	78%	
AD-12484	90%	0.023
AD-12485	76%	
AD-12486	13%	
AD-12487	60%	
AD-12488	54%	
AD-12489	11%	
AD-12490	72%	
AD-12491	86%	0.047
AD-12492	41%	
AD-12493	26%	
AD-12494	12%	
AD-12495	69%	
AD-12496	44%	
AD-12497	2%	
AD-12498	14%	
AD-12499	63%	
AD-12500	86%	0.057
AD-12501	57%	
AD-12502	88%	0.048
AD-12503	-2%	
AD-12504	8%	
AD-12505	29%	
AD-12506	31%	
AD-12507	48%	
AD-12508	47%	
AD-12509	2%	
AD-12510	-21%	
AD-12511	28%	
AD-12512	43%	
AD-12513	-22%	
AD-12514	38%	
AD-12515	-9%	
AD-12516	58%	
AD-12517	18%	
AD-12518	-8%	
AD-12519	-5%	
AD-12520	62%	1.12
AD-12521	-12%	
AD-12522	53%	
AD-12523	55%	
AD-12524	60%	
AD-12525	32%	

TABLE 9

		WBC	Platelets	Lymphocyte #
Animal #1	day 0	7.1	328	2
(AD-11570	day 3	6.9	308	1.9
treatment)	day 5	6.2	394	3.4

TABLE 9-continued

		WBC	Platelets	Lymphocyte #
Animal #2	day 0	3.6	299	1.5
(AD-11570	day 3	10.9	254	1.5
treatment)	day 5	12.9	281	1.8
	day 8	21.2	444	4.3
Animal #3	day 0	3.2	218	2.2
(AD-11570	day 3	10.9	202	1.9
treatment)	day 5	6.4	266	2.4
	day 8	18.5	306	3.6
Animal #4	day 0	9.7	398	7.3
(untreated)	day 3	8.4	448	4.4
	day 5	6.2	263	1.9
	day 8	2.8	143	1.5

NP Chimeric sequence (SEQ ID NO: 1043):
GGTACCTCGAGGAGGAAGATTAATAATTTTC

TCTCATTGAAATTTATATCGGAATTTAAATTG

AAATTGTTACTGTAATCACACCTGGTTTGTTC

AGAGCCACATCACAAGATAGAGAACAACCTA

GGTCTCCGAAGGGAGCAAGGCATCAGTGTGCT

CAGTTGAAAATCCCTTGTCAACACCTAGGTCT

TATCACATCACAAGTTCCACCTCAGACTCTGCA

GGGTGATCCAACAACCTTAATAGAAACATTAT

TGTTAAAGGACAGCATTAGTTACAGTCAAACA

AGCAAGATTGAGAATTAACCTTGGTTTGAAC

TTGAACACTTAGGGGATTGAAGATTCAACAACC

CTAAAGCTTGGGGTAAACATTGGAATAGTT

AAAAGACAAATTGCTCGGGTTTACCTGAGAGCC

TACAACATGGATAAACGGGTGAGAGGTTTCATT

GGCGCCGAGTCTCACTGAATCTGACATGGATTA

CCACAAGATCTTGACAGCAGGTCTGTCGTTTC

AACAGGGGATTGTTTCGGCAAAGAGTCATCCAG

TGTATCAAGTAAACAATCTTGAAGAAATTGTC

CAACTTATCATACAGGCCCTTGAAGCAGGTGTT

GATTTTCAAGAGAGTGCAGGACAGTTTCTTCT

CATGCTTTGTCTTCATCATGCGTACCAGGAGGA

TTACAAACTTTTCTTGGAAAGTGGCGCAGTCA

AGTATTTGGAAGGCACGGGTTCCGTTTGAAG

TCAAGAAGCGTGATGGAGTGAAGCGCCTTGAG

GAATTGCTGCCAGCAGTATCTAGTGGAAAAAAC

ATTAAGAGAACACTTGCTGCCATGCCGGAAGA

GGAGACAAGTGAAGCTAATGCCGTCAGTTTCT

CTCCTTTGCAAGTCTATTTCTACCCAACTTG

TCGTTGGAGAAAAGGCTTGCTTGAGAAGGTTCT

-continued

AAAGGCAAATTCAGTACATGCAGAGCAAGGA
 CTGATACAATATCCAACAGCTTGCAATCAGTA
 GGACACATGATGGTGATTTTCCGTTTGATGCG
 AACAAATTTTCTGATCAAATTTCTCCTAATACA
 CCAAGGGATGCACATGGTTGCCGGGCATGATG
 CCAACGATGCTGTGATTTCAAATTCAGTGGCTC
 AAGCTCGTTTTTCAGGCTTATTGATTGTCAAA
 ACAGTACTTGATCATATCTACAAAAGACAGAA
 CGAGGAGTTCGTCTCCATCTCTTGCAAGGAC
 CGCCAAGGTAAAAAATGAGGTGAACTCCTTTAA
 GGTGCACTCAGCTCCTGGCCAAGCATGGAG
 AGTATGCTCCTTTCCGCCGACTTTTGAACCTTT
 CTGGAGTAAATAATCTTGAGCATGGTCTTTTC
 CCTCAACTATCGGCAATTGCACTCGGAGTCGCC
 ACAGCACACGGGAGTACCCTCGCAGGAGTAA
 TGTGAGGAAACAGTATCAACAACCTCAGAGGGC
 TGCCACTGAGGCTGAGAAGCAACTCCAACAAT
 ACGCAGAGTCTCGCGAAGTACCATCTTGAGC
 TTGATGATCAGGAAAAGAAAATCTTATGAAC
 TTCCATCAGAAAAGAACGAAATCAGCTTCCAG
 CAAACAAACGCTATGGTAACTCTAAGAAAAGA
 GCGCCTGGCCAAGCTGACAGAAGCTATCACTGC
 TGCGTCACTGCCAAAACAAGTGGACATTACG
 ATGATGATGACGACATTCATTTCCCGGGCCGA
 TCTATGATGACGACAATCTGGCCATCAAGAT
 GATGATCCGACTGACTCACAGGATACGACCATT
 CCCGATGGTGTGTTGACCCGTATGATGGAAG
 CTACGGCGAATATCCTGACTACGAGGATTGCGC
 TGAAGGTGCACCAGATGACTTGGTCTTATTCG
 ATCTAGACGAGGACGACGAGGACACTAAGCCAG
 TGCCTAATAGATCGACCAAGGGTGGACAACAG
 AAGAACAGTCAAAAGGGCCAGCATATAGAGGGC
 AGACAGATCCGACCTTGACGAGGAGCAAAAAG
 GTGCCGAGTTGAGAACAAATCCACCACGCCAG
 TGCGCCACTCACGGACAATGACAGAAGAAATG
 AACCTCCGGCTCAACCAGCCCTCGCATGCTGA
 CACCAATTAACGAAGAGGCAGACCCACTGGAC
 GATGCCGACGACGAGAGTCTCACATCCCTGCCC
 TTGGAGTCAGATGATGAAGAGCAGGACAGGGA

-continued

CGGAACCTCCAACCGCACACCCACTGTCGCCCC
 ACCGGCTCCCGTATACAGAGATCACTCTGAAA
 AGAAAGAACTCCCGCAAGACGAGCAACAAGATC
 AGGACCACACTCAAGAGGCCAGGAACCAGGAC
 AGTGACAACACCCAGTCAGAACACTCTTTTGAG
 GAGATGTATCGCCACATTCTAAGATCACAGGG
 GCCATTTGATGCTGTTTTGTATTATCACCTAAT
 GAGTGATGAGCCTGTAGTTTTTTCAGTACCAGTG
 ATGGCAAAGAGTACACGTATCCAGACTCCCTTG
 AAGAGGAATATCCACCATGGCTCACTGAAAAA
 GAGGCTATGAATGAAGAGAATAGATTTGTTACA
 TTGGATGGTCAACAATTTTATTGGCCGGTGAT
 GAATCACAAGAATAAATTCATGGCAATCCTGCA
 ACATCATCAGTGAATGAGCATGGAACAATGGG
 ATGATTCAACCGACAATAGCTAACATTAAGTA
 GTCAAGGAACGAAACAGGAAGAATTTTGAT
 GTCTAAGGTGTAATTATTATCACAATAAAAGT
 GATTCTTATTTTTGAATTTAAAGCTAGCTTAT
 TATTACTAGCCGTTTTTCAAAGTTCAATTTGAG
 TCTTAATGCAAATAGGCGTTAAGCCACAGTTA
 TAGCCATAATTGTAACCTCAATATTCTAAGTAGC
 GATTTATCTAAATTAATTAACATTATGCTTTT
 ATAACCTACCTACTAGCCTGCCAACATTTACA
 CGATCGTTTTTATAATTAAGAAAAAGCGGCCG
 CAGAGCTC
 GP Chimeric sequence (SEQ ID NO: 1044):
 GGTACCCTCGAGGATGAAGATTAAGCCGACAGT
 GAGCGTAATCTTCATCTCTCTTAGATTATTTG
 TTTTCCAGAGTAGGGGTCGTCAGGTCCTTTTCA
 ATCGTGTAACCAAAATAAACTCCACTAGAAGG
 ATATTGTGGGGCAACAACAATGGGCGTTCTT
 AGCCTACTCCAATTGCCTCGTGATCGATTCAA
 GAGGACATCATTTCTTTCTTGGGTAATTATCCT
 TTTCCAAAGAACATTTTCCATCCCCTTGGAG
 TCATCCACAATAGCACATTACAGGTTAGTGAGA
 TTGACCAGCTAGTCTGCAAGGATCATACTGAT
 ATGCCATCTGCAACTAAAAGATGGGGCTTCAGG
 TCCGGTGTCCACCAAAGGTGGTCAATTATGA
 AGCTGGTGAATGGGCTGAAAAGTGTACAACTCT

-continued

TGAAATCAAAAACCGGACGGGAGCGAATGCT
 TACCCGCGAGCGCCAGACGGGATTCGGGGCTTCC
 CCCGGTGCCGGTATGTGCACAAAGTATCAGGA
 ACGGGACCGTGTGCCGGAGACTTTGCCTTCCAT
 AAAGAGGGTGCTTTCTTCCTGTATGATCGACT
 TGCTTCCACAGTTATCTACCGAGGAACGACTTT
 CGCTGAAGGTGTCGTTGCATTTCTGATACTGC
 CCCAAGCTAAGAAGGACTTCTTCAGCTCACACC
 CCTTGAGAGAGCCGGTCAATGCACCGGAGGAC
 CCGTCTAGTGGCTACTATTCTACCACAATTAGA
 TATCAGGCTACCGGTTTTTGGAAACCAATGAGAC
 AGAGTACTTGTTCGAGGTTGACAATTTGACCTA
 CGTCCAACTTGAATCAAGATTACACCCACAGT
 TTCTGCTCCAGCTGAATGAGACAATATATACAA
 GTGGGAAAAGGAGCAATACCACGGGAAAACCTA
 ATTTGGAAGGTCAACCCCGAAATTGATACAACA
 ATCGGGGAGTGGGCTTCTGGGAAACTAAAA
 AACCTCACTAGAAAAATTCGAGTGAAGATTG
 TCTTTCACAGTTTTATCGCTCAACGAGACAGA
 CATCAGTGGTCAGAGTCCGGCGCGAAGTTCTTC
 CGGAAGAATCTCCGACCGGGCCACTGAAGACC
 ACAAATCATGGCTTCAGAAAATTCCTCTGCAA
 TGGTTCAAGTGCACAGTCAAGGAAGGGAACA
 ACATTGCCGTCTCAGAATTGACAGAGAAGTCGA
 AGAGCGAGTCCCAATCCCTCACAACCAAAACC
 AGGTCCGACACACAGACCCATAATACACCCGT
 GTATAAACTTGACATCTCTGAGGCAACTCAAG
 TTGAACAACATCACCGCAGAACAGACAACGACA
 GCACAGCCTCCGACACTCCCTCTGCCACGACC
 GCAGCCGGACCCCCAAAAGCAGAGAACACCAAC
 ACGAGCAAGAGCACTGACTTCTTGACCCCGC
 CACCACAACAAGTCCCCAAAACACAGCGAGAC
 CGCTGGCAACAACAACACTCATCACAAGATA
 CCGGAGAAGAGAGTGCCAGCAGCGGAAGCTAG
 GCTTAATTACCAATACTATTGCTGGAGTCGCA
 GGACTGATCACAGGCGGGAGAAGAACTCGAAGA
 GAAGCAATTGTCAATGCTCAACCCAAATGCAA
 CCCTAATTTACATTACTGGACTACTCAGGATGA
 AGGTGCTGCAATCGACTGGCCTGGATACCAT

-continued

ATTCGGGCCAGCAGCCGAGGGAATTACATAG
 AGGGGCTAATGCACAATCAAGATGGTTTAATC
 TGTGGGTTGAGACAGCTGGCCAACGAGACGACT
 CAAGCTCTTCAACTGTTCTGAGAGCCACAAC
 GGAGCTGCGGACATATACCATACTCAACCGTAA
 GGCAATTGATTTCTTGCTGCAGCGATGGGGCG
 GCACATGCCACATTCTGGGACCGGACTGCTGTA
 TCGAACCACATGATTGGACCAAGAACATAACA
 GACAAAATTGATCAGATTATTCTGATTTTGT
 GATAAAACCTTCCGGACCGGGGACAATGA
 CAATTGGTGGACAGGATGGAGACAATGGATACC
 GGCAGGTATTGGAGTTACAGGCGTTATAATTG
 CAGTTATCGCTTTATTCTGTATATGCAAAATTG
 TCTTTTAGTTTTTCTTCAGATTGCTTCATGGA
 AAAGCTCAGCCTCAAATCAATGAAACCAGGATT
 TAATTATATGGATTACTTGAATCTAAGATTAC
 TTGACAAATGATAATATAATACTGGAGCTTT
 AAACATAGCCAATGTGATTCTAACTCCTTAA
 ACTCACAGTTAATCATAAACAAGGTTTGACATC
 AATCTAGTTATCTCTTGAGAATGATAAACTT
 GATGAAGATTAAGAAAAGCGGCCGAGAGCTC

[0231] The L gene was generated as 2 fragments (L-ABC, SEQ ID NO:1049; and L-DEFG, SEQ ID NO:1050).

L fragment 1: SEQ ID NO: 1049 (L-ABC):
 GCGCGCCTCGAGGAGGAAGATTAGAAAACT

GCTTATTGGGTCTTCCGTGTTTAGATGAAG
 CAGTTGAAATTTCTCTTTGATATTAATGGC
 TACCCAACTACACAATACCCAGACGCTAGTT
 ATCATACCAATTGTATTGGACCAATGTGACCT
 AGTCACTAGAGCTTGCGGGTTATATCATCAT
 ACTCCCTTAATCCGCAACTACGCAACTGTAAAC
 TCCCGAAACATATCTACCGTTTGAAATACGTG
 TAACTGTTACCAAGTTCTTGAGTGATGTACCAG
 TGGCGACATTGCCCATAGATTTTATAGTCCCA
 GTTCTTCTCAAGGCACTGTGAGGCAATGGATT
 TGTCTGTTGAGCCGCGGTGCCAACAGTTCTA
 GATGAAATCATTAAGTACACAATGCAAGATGCT
 CTCTTCTGAAATATTATCTCAAAAATGTGGG
 TGCTCAAGAAGACTGTGTTGATGAACACTTTCA

-continued

AGAGAAAATCTTATCTTCAATTCAAGGCAATG
 ATTTTACATCAAATGTTTTCTGGTATGATCT
 GGCTATTTTAACTCGAAGGGGTAGATTAAATC
 GAGGAACTCTAGATCAACATGTTTTGTTTATG
 ATGATTTAATAGACATCTTAGGCTATGGGGAC
 TTGTTTTTTGGAAGATCCCAATTCAATGTTAC
 CACTGAACACACAAGGAATCCCCATGCTGCT
 ATGGACTGGTATCAGGCATCAGTATTCAAAGAA
 GCGGTTCAAGGGCATAACACATGTTTCTGT
 TTTACTGCCGACGCTCTGATAATGTGCAAAGAT
 TTAATTACATGTCGATTCAACACAACCTCTAAT
 CTCAAAAATAGCAGAGATTGAGGATCCAGTTTG
 TTCTGATTATCCCAATTTTAAGATTGTGTCTA
 TGCTTACCAGAGCGGAGATTACTTACTCTCCAT
 ATTAGGGTCTGATGGGTATAAAATTATTAAGT
 TCCTCGAACCATTGTGCTTGGCCAAAATTCAT
 TATGCTCAAAGTACACTGAACGAAAAGGGCGG
 TTTTAACACAAATGCATTTAGCTGTAAATCACA
 CCCTAGAAGAAATTACAGAAATCGGTGCTACTA
 AAGCCTTCACAGGCTCAAAAGATCCGTGAATTC
 CATAGAACATTGATAAGGCTGGAGATGACGCC
 ACAACACTTTGTGAGCTATTTTCCATTCAAAAA
 CACTGGGGGCATCCTGTGCTACATAGTGAAAC
 AGCAATCCAAAAGTTAAAAACATGCTACGGT
 GCTAAAAGCATTACGCCCTATAGTGATTTTCG
 AGACATCTGTGTTTTTAAATATAGTATTGCCAA
 ACATTATTTTGATAGTCAAGGATCTTGGTACA
 GTGTTACTTCAGACCGATGTTTAACGCCGGGAT
 TGAATTCTTATATCAAAGAAATCAATTCCCT
 CCGTTGCAATGATTAAAGAACTACTATGGGAAT
 TTTACCACCTTGACCACCTCCACTTTTCTCA
 ACCAAAATTATTAGTGACTTAAGTATTTTATA
 AAAGACAGAGCTACCGCAGTAGAAAGGACATG
 CTGGGATGAGTATTCGAGCCTAATGTTCTAGGA
 TATAATCCACCTCACAAATTTAGTACTAAACG
 TGTACCGGAACAATTTTtagagCAAGAAACTT
 TTCTATTGAGAATGTTCTTTCATACGCCCAAG
 AACTTAGGTTCTACTACCACAATATCGGAACTT
 TTCTTTCTCATTGAAAGAGAAAGAGTTGAATG

-continued

TAGGTAGAACCTTCGGAATTTGCCTTATCCGA
 CTCGCAATGTTCAAACACTTTGTGAAGCTCTG
 TTAGCTGATGCTTGTCTAAAGCATTTCCTAGCA
 ATATGATGGTAGTTACGGAACGTGAGCAAAAA
 GAAAGCTTATTGCATCAAGCATCATGGCACCAC
 ACAAGTGATGATTTTGGTGAACATGCCACAGT
 TAGAGGGAGTACTTTGTAACTGATTTAGAGAAA
 TACAATCTTGCATTTAGATATGAGTTTACAGC
 ACCTTTTATAGAATATTGCAACCGTTGCTATGG
 TGTTAAGAATGTTTTAATGGATGCATTATA
 CAATCCACAGTTTATATGCATGTCAGTGATTA
 TTATAATCCACCACATAACCTCACACTGGAGA
 ATCGAGACAACCCCCCGAAGGGCCTAGTTCAT
 ACAGGGGTCTATGGGAGGGATTGAAGGACTG
 CAACAAAACCTCTGACAAGTATTTTATGTGCTC
 AAATTTCTTTAGTTGAAATTAAGACTGGTTTT
 AAGTTACGCTCAGCTGTGATGGGTGACAATCAG
 TGCATTACTGTTTTATCAGTCTTCCCCCTAGA
 GACTGACGCAGACGGCAGGAACAGAGCGCCGAA
 GACAATGCAGCGAGGGTGGCCGCCAGCCTAGC
 AAAAGTTACAAGTGCCTGTGGAATCTTTTAA
 ACCTGATGAGACTTTCGTACACTCAGGTTTTA
 TCTATTTTGAAAAACAATATTTGAATGGGGT
 CCAATTGCCTCAGTCCCTTAAACGGCTACAA
 GAATGGCACCATTGTCTGATGCAATTTTGTATG
 ATCTTCAAGGGACCTGGCTAGTATAGGCACT
 GCTTTTGAGCGATCAACTCCGAAACTAGACATA
 TCTTTCCTTGCAGGATAACCGCAGCTTTCAT
 ACGTTTTTTTCGGTGAGAATCTTGCAATATCAT
 CATCTCGGGTTCAATAAAGTTTTTGACCTTGG
 ACAGTTAACACTCGGCAACCTCTGGATTTCCGA
 ACAATATCATTGGCACTAGCGGTACCGCAGGT
 GCTTGGAGGGTTATCCTTCTTGAATCCTGAGAA
 ATGTTTCTACCGAATCTAGGAGATCCAGTTA
 CCTCAGGCTTATTCAGTAAAACTTATCTCCG
 AATAGAGACCTATTGAGCTCCACCGCGGTGGC
 GGCCGCTCTAGCCGGGGCGGATCCCCGGGCTG

-continued

CAGGAATTCGATATCAAGCTTATCGATACCGT
CGACCTCGAGGGGGGCGCCGTACCTTACATCGC
GTTAATTAAGTAGTGGATCGATCCCCAATTCG
L fragment 2: SEQ ID NO: 1050 (L-DEFG):
CGCGACGTAATACGACTCACTATAGGGCGAATT
GGGCGCGCCCCGCTCAGAATGATTGAGATGG
ATGATTTATTCTTACCTTTAATTGCGAAGAACC
CTGGGAATTGTAGCGCAATTGACTTTGTGTAA
ATCCTAGCGGATTAAATGTCCCTGGGTCGCAAG
ACTTAACCTTCATTTCTGCGCCAGATTGTACGC
AGGACCATCACCCCTAAGTGCAGAAAAACAACTT
ATTAATACCTTATTTCATGCGTCAGCTGACTC
GAAGACGAAATGGTTTGTAAATGGCTATTATCA
TCAACTCCTGTTATGAGTCGTTTTCGCGCCGA
TATCTTTTTCACGACGCGGAGCGGGAAGCGATT
GCAAATTCAGGATACCTGGAAGGAACACGCC
ATTATTAGCCTCTAAGATCATCAACAATAATAC
AGAGACACCGGTTTGGACAGACTGAGGAAAA
TAACATTGCAAAGGTGGAGCCTATGGTTTAGTT
ATCTTGATCATTGTGATAATATCCTGGCGGAG
CTTTAACCCAAATAACTTGACAGTTGATTTAG
CACAGATTCTGAGGGAATATTCATGGGCTCAT
ATTTTAGAGGAAGACCTCTTATTGGAGCCACA
CTCCCATGTATGATTGAGCAATTCAAAGTGTT
TGGCTGAAACCCCTACGAACAATGTCGCGAGTGT
TCAAATGCAAAGCAACCGTGGGAACCAT
CGTGTGAGTGGCAGTCAAGAAACATATTGTTAG
TGCATGGCCGAACGCATCCGAATAAGCTGGA
CTTCGGGGATGGAATCCCATACATTGGATCAAG
GACAGAAGATAAGATAGGTGAGCCGCTATTA
AGCCGAGGTGTCCTTCGCGAGCCTTAAGAGAGG
CCATTGAATTGGCGTCCCGTTTAAACATGGGTA
ACTAAGGCGAGTTCGAACAGTGACTTGCTAATAA
AACCATTTTGGGAAGCAGAGTAAATTTAAGT
GTTCAAGAAATACTTCAAATGACCCCTTCACAT
TACTCAGGAAATATTGTGATCGGTATAACGA
TCAAACAGTCCCTCATCTTTTCATGGCCAATCGT
ATGAGTAATTGAGCAACGCGCTTGATGGTATC
TACAAACACTTTAGGTGAGTTTTCAGGAGGTGG

-continued

CCAGTCTGCACGCGACAGCAATATTATTTTCC
AGAATTTATAAATTATGCAGTTGCACTGTTCGA
TATTAAATTAGAAAACACTGAGGCTACAGATA
TCCAATATAATCGTGCTCACCTTCATCTAACTA
AGTGTGACACCCGGGAAGTACCAGCTCAGTAT
TTAACAACACAACCACGCTAAATCTAGATTAA
CAAGATACCGAGAAAAAGCAATTGATTTATGAC
AGTAATCCTCTAAAAGGAGGACTCAATTGCAAC
TTATCGATTGACAGTCCTTTTTCAGGTA
ACGGCTGACATTATAGAAGATGATCTTATTCGA
CTGCCTCACTTATCTGGATGGGAGCTAGCCAA
GACCATCATGCAATCAATTATTCAGATAGCAA
CAATTCATCTACAGACCAATTAGCAGTGGAG
AAACAAGACATTCCTACCCATTTCTTAACCTA
TCCCAAGATAGGACTTCTGTACAGTTTGGGG
CCTTTGTAAGTTATTATCTTGGCAATACAATTC
TTTGACGAAAAAGATCGGACTTGACAATTTT
TTATATTACTAATACTCAAAATCATAATCTAC
CACATCGCTCATTCGAATACTTAAGCCAACA
TTCAAACATGCAAGCGTTATGTACAGGTTAATG
AGTATTGATCCTCATTTTCTATTACATAGG
CGGTGCTGCAGTGACAGAGGACTCTCAGATGCG
GCCAGGTTATTTTGGAGACGTCATTTTCATC
TTTTCTTACATTTGTAAAAGAAATGGATAATTAA
TCGCGGAACAATTGTCCCTTTATGGATAGTAT
ATCCGCTAGAGGTCAAACCCCAACCTGTGAA
TAATTTTCTCTATCAGATCGTAGAAGTGTGG
TGATGATTCATCAAGACAACAGGCTTTTAAAA
CTACCATAAGTGATCATGTACATCCTCACGAC
AATCTTGTGTACCATGTAAGAGTACAGCCAGCA
ATTTCTTCCATGCATCATTGGCGTACTGGAGG
AGCAGACACAGAAACAGCAACCGAAATACTTG
GCAAGAGACTCTTCAACTGGATCAAGCACAAA
CAACAGTGATGGTATATTGAGAGAAGTCAAGAA
CAAACACCAGAGATCCACATGATGGCACTGA
ACGGAATCTAGTCTTACAAATGAGCCATGAAT
AAAAAGAAGCACAATTCACAAGAAAAACGCG
ACCAGGTCCTGCTGTCAGTCCTTTCTAAGTGA
CTCTGCTTGTGGTACAGCAATCCAAACTAA

-continued

ATTTTCGATCGATCGAGACACAATGTGAAATTTTC
 AGGATCATAACTCGGCATCCAAGAGGGAAGGT
 CATCAAATAATCTCAACCGTCTAGTCCTACCTT
 TCTTTACATTATCTCAAGGGACACGCCAATTA
 ACGTCATCCAATGAGTCACAAACCCAAGACGAG
 ATATCAAAGTACTTACGGCAATTGAGATCCGT
 CATTGATACTACCATAATTGTCGCTTCACCGGT
 ATAGTCTCGTCCATGCATTACAACTTGATGA
 GGTCTTTTGGGAAATAGAGAGTTTCAAGTCGGC
 TGTGACGCTAGCAGAGGGAGAAGGTGCTGGTG
 CCTTACTATTGATTCAAATACGGCGTTAAGAA
 GTTATTTTTCAACACGCTAGCTACTGAGTCCA
 GTATAGAGTCAGAAATAGTATCAGGAATGACTA
 CTCCTAGGATGCTTCTACCTGTTATGTCAAAA
 TTCCATAATGACCAAATTAGATTATTCTTAACA
 ACTCAGCAAGCCAAATAACAGACATAACAAAT
 CCTACTTGGTTTAAAGACCAAAGAGCAAGGCTA
 CCTAAGCAAGTCGAGGTTATAACCATGGATGC
 AGAGACAACAGAGAATATAACAGATCGAAATTG
 TACGAAGCTGTATATAAATTGATCTTACACCA
 TATTGATCCTAGCGTATTGAAAGCAGTGGTCCT
 TAAAGTCTTTCTAAGTGATACTGAGGGTATGT
 TATGGCTAAATGATAATTTACCCGTTTTTTGC
 CACTGGTTATTTAATTAAGCCAATAACGTCAA
 GTGCTAGATCTAGTGAGTGGTATCTTTGTCTGA
 CGAACTTCTTCACTACACGTAAGATGCCA
 CACCAAACCATCTCAGTTGTAACAGGTAATAC
 TTACGGCATTGCAACTGCAAATTCACGAAGC
 CCATACTGGCTAAGTCATTTAACTCAGTATGCT
 GACTGTGAGTTACATTTAAGTTATATCCGCCT
 TGGTTTTCCATCATTAGAGAAATACTATACCAC
 AGGTATAACCTCGTCGATTCAAAAAGAGTCC
 ACTAGTCTCTATCACTCAGCACTTAGCACATCT
 TAGAGCAGAGATTCGAGAATTAATAATGATT
 ATAATCAACAGCGACAAAGTCGACCCAGACTTA
 TCATTTTATTCGTACTGCAAAAGGACGGATAA
 CTAAACTAGTCAATGATTATTTAAATTTCTTTC
 TTATTGTGCAAGCATTAAAAATAATGGGACA
 TGGCAAGCTGAGTTTAAGAAATTACAGAGTTGA

-continued

TTAGTGTGTGCAATAGGTTCTACCATATTAGA
 GATTGCAATTGTGAAGAACGTTTCTTAGTTCAA
 ACCTTATATTTACATAGAATGCAGGATTCTGA
 AGTTAAGCTTATCGAAAGGCTGACAGGCTTCTG
 AGTTTATTTCCGGATGGTCTCTACAGGTTTGA
 TTGAATTACCGTGCATAGTATCCTGATACTTGC
 AAAGGTTGGTTATTAAACATACAGATTATAAAA
 AAGCGGCCGCAGAGCTCCAGCGGTGGGGCCGCG
 GGCGTCTAGCCCGGGCGGATCCCTGCAGGAAT
 TCGATATCAAGCTTATCGATACCGTCGACCTCG
 AGGGCCCATGCAGGCCGCCAGGTACCTTAGT
 TAATTAACAGCTTTTGTTCCTTTAGTAGGGTT
 AATTGACGCGCTC
 VP24 (SEQ ID NO: 1045):
 GCGCGCCTCGAGGATGAAGATTAATGCGGAGG
 TCTGATAAGAATAAACCTTATTATTAGATTA
 GGCCCCAAGAGGCATTCTTCATCTCCTTTTAGC
 AAAGTACTATTTTCAAGGTAGTCCAATTAGTGG
 CACGTCTTTTAGCTGTATATCAGTCGCCCTAG
 GCTAGGGTTTATAGTTGTCTCTAAGCTAAATT
 GGTCTTGAAGTACTGCTACTCGCAGAATCCTACC
 GGAATAGACTAATTGAAGTACCCGTTTAA
 ATTTAGTGCATAAATCTGGGCTAACACCAACGAG
 GTCAACTCCATTGGCTGAAAAGAAGCTTACCT
 ACAACGAACATCACTTTGAGCGCCCTCACAATT
 AAAAAATAGGAACGTCGTTCCAACAATCGAGC
 GCAAGGTTTCAATATTATACGGGTCCATTAATT
 TCAACAAAATATTGATACTCCAGACACCAAGC
 AAGACCTGAGAAAAAACCTATGGCTAAAGCTACG
 GGACGATACAATCTAATATCGCCCAAAAGGA
 CCTGGAGAAAGGGTTGTCTTAAGCGACCTCTG
 TAACTTCTTAGTTAGCCAACTATTAGGGGT
 GGAAGGTTTATTGGGCTGGTATTGAGTTTGATG
 TGAACCAAAAGGGTATTACCTTATGCATAGA
 CTGAAAATAATGACTTTGCCCCGTCATGGTCA
 ATGACAAGGAATCTCTTCTCATTATTTTCA
 AAATCCGAATTCCACAATTGAATCACCGCTGTG
 GGCATTGAGAGTCATCCTTGACGAGGGATAC
 AGGACCAGCTGATTGACCAGTCTTTGATTGAAC
 CCTTAGCAGGAGCCCTTGGTCTGATCTCTGAT

-continued

TGGCTGCTAACAACCAACACTAACCATTTC AAC
ATGCGAACACAACGTGTCAAGGAACAATTGAG
CCTAAAAATGCTGTCTGTGATTCGATCCAATAT
TCTCAAGTTTATTAAACAAATTGGATGCTCTAC
ATGTCTGTAACACACGGAATTGTTGAGCAGTA
TTGAAATTGGAACTCAAAATCATACAATCATC
ATAACTCGGACTAATATGGGTTATCTTGTGAG
CTCCAAGAACCCGACAAATCTGCGATGGATAT
ACGACACCCCTGGGCCGGCGAAATTTTCCTTACT
ACATGAATCGACACTTAAAGCATTACACAAG
GATCCTCGACACGAATGCAAAGTTTGATTCTTG
AATTTAATAGCTCTCTTGCTATCTAACTAAGG
TAGAAAAAATTGTACGATAGGGCTAACATTATG
CTGACTCAATAGTTATCTTGACATCTCTGCTT
TCATAATCAGATATATAAGCATAATAAATAAAT
ACTCATATTTCTTGATAATTTGTTTAAACACA
GATAAATCCTCACTGTAAGCCAGCTTCCAAGTT
GACACCCCTTACAAAACCGAGCTCAGAATCC
CTCAACAAGAGATTCCAAGACAACATCATAGA
ATTGCTTTATTATATGAATAAGCATTTTATCA
CCAGAAATCCTATATACTAAATGGTTAATTGTA
ACTGAACCCGCGAGGTACATGTGTTAGGTTTC
ACAGATTCTATATATTACTAACTCTAGAGCCCA
AATTAACACGGTATAAGTAGATTAAAGAAAAA
GCCTGAGGAAGATTAAAGAAAAAGCGGCCGATT
AATTAA
VP30 (SEQ ID NO: 1046):
GGCGCGCCTCGAGGATGAAGATTAAAGAAAAAG
TAATCTTTTCGATTATCTTTAATCTTCATCCTT
GATTCTACAATCATGACAGTTGTCTTTAGTGAC
AAGGGAAGAAGCCTTTTATTAAAGTTGTAAT
AATCAGATCTGCGAACCGGTAGAGTTTAGTTGC
AACCTAACACACATAAAGCATTGGTCAAAAAG
TCAATAGAAATTTAAACAGTGAGTGGAGACAAC
TTTTAAATGGAAGCTTCGTGAGCGCGGAGAT
CAAGGAATTCACGTGCCGACCAGCAAGGGATGG
ACACGACCACCATGTTTCGAGCACGATCATCAT
CCAGAGAGAATTATCGAGGTGAGTACCGTCAAT
CAAGGAGCGCCTCACAAGTGCGCGTTTCCTACT

-continued

GTATTCATAAGAAGAGAGTTGAACCATTAAACA
GTTCTCCAGCACCTAAAGACATATGTCCGAC
CTTGAAAAAAGGATTTTGTGTGACAGTAGTTT
TTGCAAAAAAGATCACCAGCTTGAAAGCCTAA
CCGACCGGAATTACTCTACTAATCGCCCGTA
AGACTTGTGGATCAGTTGATTCATCGCTTAAT
ATAACTGCACCCAAGGACTCGCGCTTAGCAAAT
CCAACGGCTGATGATTCCAGCAAGAGGAAGG
TCCAAAAAATTACTAGTCGAGACTGCTCAAGAC
GGCAGAACTGCGGCGAGACAAGACATCAGAA
CCATAGAGGATTCAAATTAAGAGCATTGTTGA
CTCTATGTGCTGTGATGACGAGGAAATCTCA
AAATCCCAGCTGAGTCTTTTATGTGAGACACAC
CTAAGGCGCGAGGGGCTTGGGCAAGATCAGGC
AGAACCCTTCTCGAAGTATATCAACGATTACA
CAGTGATAAAGGAGGCAGTTTTGAAGCTGCAC
TATGGCAACAATGGGACCGACAATCCCTAATTA
TGTTTACTACTGCATTCTTGAATATTGCTCTC
CAGTTACCGTGTGAAAGTTCTGCTGCTGTTGTT
TCAGGCCTACGCTTACTTGCCCCCAAGCGT
TAATGAAGAAGCTTCAACCAACCGGGGACATG
CTCATGGTCTGATGAGGGTACCCCTTAATAAG
GCTGACTAAAACACTATATAACCTTCTACTTGA
TCACAATACTCCGTATACCTATCATCATATAT
TTAATCAAGACGATATCCTTTAAACTTATTCA
GTACTATAATCACTCTCGTTTCAAATTAATAA
GATGTGCATGATTGCCCTAATATATGAAGAGGT
ATGATACAACCCTAACAGTGATCAAAGAAAAT
CATAATCTCGTATCGCTCGTAATATAACCTGCC
AAGCATACTCCCTAGAAGCGTTGAATCTTGTA
CACAAATAATGTTTTACTCTACAGGAGGTAGCA
ACGATCCATCCCATCAAAAAATAAGTATTTCA
TGACTTACTAATGATCTCTTAAAAATATTAAGAA
AAAGCGGCCGATTAAATTAA
VP35 (SEQ ID NO: 1047):
GGTACCGCGATCGCGATGAAGATTAAACCTTC
ATCATCCTTACGTCAATTGAATTCTCTAGCAC
TCGAAGCTTATTGTCTTCAATGTAAAAAGAAAAG
CTGGTCTAACAAGATGACAAC TAGAACAAAGG
GCAGGGGCCATACTGCGGCCACGACTCAAAACG

-continued

ACAGAATGCCAGGCCCTGAGCTTTCGGGCTGG
ATCTCTGAGCAGCTAATGACCGGCAAAATACCG
CTAACCGACATCTTCTGTGATATTGAGAACAA
TCCAGGATTATGCTACGCATCCCAATGCAACA
AACGAAGCCAAACCCGAAGACGCGCAACAGTC
AAACCCAAACGAGCCCAATTTGCAATCATAGTT
TTGAGGAGGTAGTACAAACATTGGCTTCATTG
GCTACAGCTGTGCGTCGGCAAACCATCGCATCA
GAATCATTAGAACACGCATTACGAGTCTTGA
GAATGGTCTAAAGCCAGTTTATGATATGGCAAA
AACAATATCATCCCTGAATCGCAGCTGTGCTG
AGATGGTTGCAAAATATGATCTTCTGGTGATGA
CAACCGGTCTGGGCAACAGCAACCGCTGCGGCA
ACTGAGGCTTATTGGGCCGAACATGGTCAACCA
CCACCAGGCCCATCATTTGTACGAGGATGGTGC
GATTCGGGGTAAATTGAAAGATCCGAACGGGAC
CGTCCCTCAAAGTGTTAGGGAGGCATTCAACA
ATCTAAACAGTACCACCTTCACTAACTGAGGAAA
ATTTCTGGGCGACCTTACATTTCTGGCAAAGGAT
TTGAGAAACATTATGTATGATCACTTGCCCTGGT
TTTGGAAGTCTTTCCACCAATTAGTACAAGT
GATTTGTAAATTGGGAAAAGATAGCAACTCATT
GGACATCATTCATGCTGAGTTCAGGCCAGCC
TGGCTGAAGGAGACTCTCCTCAATGTGCCCTAA
TTCAAATTACAAAAAGAGTTCCAATCTTCAA
GATGCTGCTCCACCTGTCATCCACATCCGCTCT
CGAGGTGACATTCCTCCGAGCTTGCCAGAAAAAG
CTTGCGTCCAGTCCCACCATCGCCCAAGATTGA
TCGAGGTTGGGTATGTGTTTTTTCAGCTTCAAG
ATGGTAAAACACTTGGAATCAAATTTGAGCCA
ATGTAAGCTCATTTTGCATGGGCGAATAATA
GCAGAGGCTTCAACTGCTGAACATAGGGTACG
TTACATTAATGATACACTTGTGAGTATCAGCC
CTGGATAATATAAGTCAATCCTAATCAATTGAT
AATATTGTTTATATCTCGCTAGCAGCTTAAAA
TATAAATGTAATAGGAGCTATATCTCTGACAGT
ATTATAATCAATTGTTATTAAAGTAACCCAAAC
CAAAAGTGATGAAGATTAAGAAAAAGCGGCCGC
AGAGCTC

-continued

VP40 (SEQ ID NO: 1048):
GGTACCTCGAGGATGAAGATTAAGAAAAACCTA
CCTCGGCTGAGAGAGTGTTTTTTCATTAACTT
TCATCTTGTAACGTTGAGCAAAATGTTAAAA
ATATGAGGCGGGTTATATTGCCTACTGCTCCT
CCTGAATATATGGAGGCCATATACCTGTGAGG
TCAAATTCACAAATTGCTAGAGGTGGCAACAG
CAATACAGGCTTCTTGACACCGGAGTCAGTCAA
TGGGGACACTCCATCGAATCCACTCAGGCCAA
TTGCCGATGACACCATCGACCATGCCAGCCACA
CACCAGGCGAGTGTGTCATCAGCATTCACTCTT
GAAGCTATGGTGAATGTATATCGGGCCCCAAA
GTGCTAATGAAGCAAAATCCCTATTTGGTTGCC
TCTAGGTGTCGCTGATCAAAAGACCTACAGCTT
TGACTCAACTACGGCCGCAATTATGCTCGCAT
CTTATACGATCACCATTTCGGCAAGGCAACCA
ACCCCTCGTTAGAGTGAATCGACTGGGTCTT
GGAATCCCGGATCATCCCTCAGGCTCCTGCGA
ATTGGAACACAGGCTTTCCTCCAGGAGTTCGT
TCTTCCGCCAGTCCAACCTACCCAGTATTTTAC
CTTTGATTTGACAGCACTCAAACCTGATCACC
AACCCTGCCTGCTGCAACATGGACCGATGACA
CTCCAACAGGATCAAATGGAGCGTTGCGTCCA
GGAATTTTCAATTCATCCAAACTTCGCCCCATT
CTTTTACCAACAAAAGTGGGAAGAAGGGGAA
CAGTGCCGATCTAACATCTCCGAGAAAATCCA
AGCAATAATGACTTCACTCCAGGACTTTAAGA
TCGTGCCAATTGATCCAGCCAAGAGTATCATTG
GGATCGAGGTGCCAGAACTCTGGTCCACAAG
CTGACCGGTAAGAAGGTGACTTCTAAAAATGGA
CAACCAATCATCCCTGTTCTTTTGCCAAAGTA
CATTGGGTGGACCCGGTGGCTCCAGGAGACCT
CACCATGGTAATCACACAGGATTGTGACACGT
GTCATTCTCTGCAAGTCTTCCAGCTGTGATTG
AGAAGTAATTGCAATAATTGACTCAGATCCAG
TTTTATAGAATCTTCTCAGGGATAGCAACTCAA
TCGACTTTTAGGACCGTCCATTAGAGGAGACA
CTTTTAATTGAAAAATGTACTAATCGGGTCAAG
GACCATTGCTTTTTTCTCTCCTAAATGTAGA

-continued
 ACTTAACAAAGACTCATAATATACTTGTTTT
 AAAGGATTGATTGATGAAAGAACATGCATAAG
 CGATCCATACTTCGCCCTACTATAATCAATACG
 GTGATTCAAATGTTAATCTTTCTCATTGCACA
 TACTTTTTGCCCTTATCCTCAAATTGCCTGCAT

-continued
 GCTTACATCTGAGGATAGCCAGTGTGACTTGG
 ATTGGAAATGTGGAGAAAAATCGGGACCCATT
 TCTAGGTTGTTTACAATCCAAGTACAGACATT
 GCCCTTCTAATTAAGAAAAAGCGCCGCAGAG
 CTC

[0232] Other embodiments are in the claims.

SEQUENCE LISTING

The patent application contains a lengthy "Sequence Listing" section. A copy of the "Sequence Listing" is available in electronic form from the USPTO web site (<http://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US20160264972A1>). An electronic copy of the "Sequence Listing" will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

We claim:

1. A double-stranded ribonucleic acid (dsRNA) for inhibiting the expression of a human Ebola genome in a cell, wherein said dsRNA comprises at least two sequences that are complementary to each other and wherein a sense strand comprises a first sequence and an antisense strand comprises a second sequence comprising a region of complementarity which is substantially complementary to at least a part of a mRNA encoding Ebola, and wherein said region of complementarity is less than 30 nucleotides in length.

2. The dsRNA of claim 1, wherein said first sequence is selected from the group consisting of the sense sequences of Table 2 and said second sequence is selected from the group consisting of the antisense sequences of Table 2.

3. The dsRNA of claim 1, wherein said dsRNA comprises at least one modified nucleotide.

4. The dsRNA of claim 2, wherein said dsRNA comprises at least one modified nucleotide.

5. The dsRNA of claims 3, wherein said modified nucleotide is chosen from the group of: a 2'-O-methyl modified nucleotide, a nucleotide comprising a 5'-phosphorothioate group, and a terminal nucleotide linked to a cholesteryl derivative or dodecanoic acid bisdecylamide group.

6. The dsRNA of claim 3, wherein said modified nucleotide is chosen from the group of: a 2'-deoxy-2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a locked nucleotide, an abasic nucleotide, 2'-amino-modified nucleotide, 2'-alkyl-modified nucleotide, morpholino nucleotide, a phosphoramidate, and a non-natural base comprising nucleotide.

7. The dsRNA of claims 3, wherein said first sequence is selected from the group consisting of the sense sequences of Table 2 and said second sequence is selected from the group consisting of the antisense sequences of Table 2.

8. The dsRNA of claim 6, wherein said first sequence is selected from the group consisting of the sense sequences of Table 2 and said second sequence is selected from the group consisting of the antisense sequences of Table 2.

9. A cell comprising the dsRNA of claim 1.

10. A pharmaceutical composition for inhibiting the expression of a gene from an Ebola virus in an organism, comprising a dsRNA of claim 1 and a pharmaceutically acceptable carrier.

11. The pharmaceutical composition of claim 10, wherein said first sequence of said dsRNA is selected from the group consisting of the sense sequences of Table 2 and said second sequence is selected from the group consisting of the antisense sequences of Table 2.

12. The pharmaceutical composition of claim 10, wherein said first sequence of said dsRNA consists of the sequence of SEQ ID NO:1029, and said second sequence consists of the sequence of SEQ ID NO:1030.

13. A method for inhibiting the expression of a gene from an Ebola virus in a cell, the method comprising:

(a) introducing into the cell a double-stranded ribonucleic acid (dsRNA) of claim 1; and

(b) maintaining the cell produced in step (a) for a time sufficient to obtain degradation of the mRNA transcript of a gene from the Ebola virus, thereby inhibiting expression of a gene from the Ebola virus in the cell.

14. A method of treating, preventing or managing pathological processes mediated by Ebola expression comprising administering to a patient in need of such treatment, prevention or management a therapeutically or prophylactically effective amount of a dsRNA of claim 1.

15. A vector for inhibiting the expression of a gene from the Ebola virus in a cell, said vector comprising a regulatory sequence operably linked to a nucleotide sequence that encodes the dsRNA of claim 1.

16. A cell comprising the vector of claim 15.

17. The dsRNA of claim 1, wherein the dsRNA targets the VP35 of Ebola.

18. The dsRNA of claim 1, wherein the dsRNA has a sense strand consisting of the sequence of SEQ ID NO:1029, and antisense strand consisting of the sequence of SEQ ID NO:1030.

19. The dsRNA of claim 1, wherein said dsRNA, upon contact with a cell infected with Ebola virus, inhibits expression of a gene from the virus by at least 40%.

20. The dsRNA of claim 1, wherein said region of complementarity is 15-30 nucleotides in length.

* * * * *